

Swapan Dey

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5347966/swapan-dey-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

50
papers

688
citations

16
h-index

24
g-index

57
ext. papers

804
ext. citations

3.1
avg, IF

4.13
L-index

#	Paper	IF	Citations
50	d-Mannitol based surfactants for cosmetic and food applications and hydrogels to produce stabilized Ag nanoparticles. <i>New Journal of Chemistry</i> , 2022 , 46, 6193-6200	3.6	1
49	Strategy to Design a Flexible and Macromolecular Sensor to Bind Cd Ions: A Complete Photophysical Analysis and Bio-Imaging Study. <i>ACS Omega</i> , 2021 , 6, 27936-27945	3.9	0
48	Application of the Meerwein reaction of 1,4-benzoquinone to a metal-free synthesis of benzofuropyridine analogues. <i>Beilstein Journal of Organic Chemistry</i> , 2021 , 17, 977-982	2.5	4
47	Dicyanamide-intertwined assembly of two new Zn complexes based on NO-type pro-ligand: Synthesis, crystal networks, spectroscopic insights, and selective nitroaromatic turn-off fluorescence sensing. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2021 , 254, 119612	4.4	12
46	Tandem Nenitzescu Reaction/Nucleophilic Aromatic Substitution to Form Novel Pyrido Fused Indole Frameworks. <i>European Journal of Organic Chemistry</i> , 2021 , 2021, 4865-4875	3.2	2
45	Novel Class of Isoxazole-Based Gelators for the Separation of Bisphenol A from Water and Cleanup of Oil Spills. <i>ACS Omega</i> , 2020 , 5, 8613-8618	3.9	2
44	Dual core clickate fluorophores for selective recognition of Cu ²⁺ and Ni ²⁺ along with live cell imaging. <i>Inorganica Chimica Acta</i> , 2020 , 509, 119655	2.7	1
43	Template-Assisted Regioselective Identification of Metal Ions on Coumarin-Furan Conjugated Chemosensors: AIEE Effect and Photo-Switching pH Indicator by ICT. <i>European Journal of Inorganic Chemistry</i> , 2020 , 2020, 3771-3777	2.3	1
42	Unusual absence of FRET in triazole bridged coumarin-hydroxyquinoline, an active sensor for Hg ²⁺ detection. <i>Photochemical and Photobiological Sciences</i> , 2020 , 19, 1211-1221	4.2	5
41	Synthesis, characterizations and single crystal structure of di-nuclear azido-bridged Cd(II) coordination polymer with Schiff base precursor (H ₂ LpentOMe): DFT, fluorescence, solvatochromism and in vitro antimicrobial assay. <i>Inorganica Chimica Acta</i> , 2019 , 496, 119069	2.7	10
40	Photo-physical aspects of BODIPY-coumarin conjugated sensor and detection of Al ³⁺ in MCF-7 cell. <i>Supramolecular Chemistry</i> , 2019 , 31, 695-702	1.8	2
39	Detection of Hg ²⁺ ion using highly selective fluorescent chemosensor in real water sample and in-vitro cell study upon breast adenocarcinoma (MCF-7). <i>Supramolecular Chemistry</i> , 2019 , 31, 382-390	1.8	3
38	Polyurethane network using 1-naphthylamine embedded epoxy-based polymer: ferric ion selective fluorescent probe. <i>Polymer Bulletin</i> , 2019 , 76, 205-213	2.4	3
37	Recognition of Al ³⁺ through the off-on mechanism as a proficient driving force for the hydrolysis of BODIPY conjugated Schiff base and its application in bio-imaging. <i>Inorganica Chimica Acta</i> , 2019 , 498, 119157	2.7	8
36	Heterometallic Zn(II)-K(I) complex with salen-type Schiff-base ligand: Synthesis, crystal structure, solid-state photoluminescent property and theoretical study. <i>Journal of Molecular Structure</i> , 2019 , 1185, 112-120	3.4	13
35	Syntheses, crystal structures and photo physical aspects of azido-bridged tetranuclear cadmium (II) complexes: DFT/TD-DFT, thermal, antibacterial and anti-biofilm properties. <i>Journal of Molecular Structure</i> , 2019 , 1179, 694-708	3.4	15
34	Epoxy-based polymer incorporating 1-naphthylamine and sebacic acid moieties: A selective fluorescent sensor for ferric ions. <i>Journal of Molecular Structure</i> , 2019 , 1180, 406-410	3.4	3

33	Highly selective and safe <i>in vitro</i> detection of biologically important Uric acid in living cells by a new fluorescent turn-on probe along with quantum chemical calculation. <i>Sensors and Actuators B: Chemical</i> , 2018 , 264, 208-215	8.5	6
32	Synthesis, characterization, physical and thermodynamic properties of a novel anionic surfactant derived from .. <i>RSC Advances</i> , 2018 , 8, 24485-24499	3.7	36
31	Water switched aggregation/disaggregation strategies of a coumarin- <i>l</i> -aphthalene conjugated sensor and its selectivity towards Cu ²⁺ and Ag ⁺ ions along with cell imaging studies on human osteosarcoma cells (U-2 OS). <i>New Journal of Chemistry</i> , 2018 , 42, 10983-10988	3.6	21
30	Nitrato, Pseudohalo-Linked Zn(II)/Cd(II) Schiff-Base Complexes with 1,3-Diimine Spacer Group: Syntheses, Crystal Structures, DFT, TD-DFT and Fluorescence Studies. <i>ChemistrySelect</i> , 2018 , 3, 12371-12382	1.8	9
29	Intracellular detection of hazardous Cd through a fluorescence imaging technique by using a nontoxic coumarin based sensor. <i>Dalton Transactions</i> , 2017 , 46, 2524-2531	4.3	33
28	Detection of Hg ²⁺ and Cs ⁺ with a Rhodamine-Based Sensor and Ethoxy-Substituted Dihydroimidazole Ring Formation Associated with the Reduction of Hg ²⁺ to Hg. <i>ChemistrySelect</i> , 2017 , 2, 1106-1110	1.8	14
27	Synthesis of Rhodamine-Based Chemosensor for Fe ³⁺ Selective Detection with off-on Mechanism and its Biological Application in DL-Tumor Cells. <i>ChemistrySelect</i> , 2017 , 2, 2969-2974	1.8	9
26	A bis-hydrazone derivative of 2,5-furandicarboxaldehyde with perfect hetero-atomic cavity for selective sensing of Hg(II) and its intracellular detection in living HeLa S3 cell. <i>Sensors and Actuators B: Chemical</i> , 2017 , 243, 1181-1190	8.5	12
25	Characterizations of surfactant synthesized from palm oil and its application in enhanced oil recovery. <i>Journal of the Taiwan Institute of Chemical Engineers</i> , 2017 , 81, 343-355	5.3	63
24	A Non-Perilous Coumarin-Based Ratiometric Probe for <i>In Vitro</i> Detection of Cu through Cell Imaging Technique. <i>ChemistrySelect</i> , 2017 , 2, 8270-8277	1.8	5
23	Indole-based distinctive chemosensors for <i>naked-eye</i> detection of CN and HSO ₄ ⁻ associated with hydrogen-bonded complex and their DFT study. <i>Supramolecular Chemistry</i> , 2016 , 28, 239-248	1.8	16
22	A real time colorimetric <i>two in one</i> kit for tracking ppb levels of uric acid and Hg ²⁺ in live HeLa S3 cells and Hg ²⁺ induced keto-enol tautomerism. <i>RSC Advances</i> , 2016 , 6, 62990-62998	3.7	13
21	Lead ion induced chemodosimeter approach of a tripodal hydroxyl-quinoline based phospho-ester through P-O bond cleavage. <i>Dalton Transactions</i> , 2016 , 45, 9187-92	4.3	14
20	Colorimetric <i>turn on</i> TBET sensors for <i>in the spot</i> visual detection of HSO ₄ ⁻ in semi-aqueous medium: Experimental and quantum chemical (DFT) studies. <i>Sensors and Actuators B: Chemical</i> , 2015 , 221, 849-856	8.5	13
19	Directed molecular recognition: furfurylamine appended ditopic receptor for succinic acid. <i>Supramolecular Chemistry</i> , 2014 , 26, 769-776	1.8	4
18	A series of ditopic receptors for succinic acid binding. <i>RSC Advances</i> , 2014 , 4, 51486-51495	3.7	6
17	Steric inhibition of hydrogen bonding in molecular recognition of dicarboxylic acids: di-topic receptors containing a nitro group designed to behave like monotopic receptors. <i>RSC Advances</i> , 2014 , 4, 49663-49671	3.7	1
16	Naphthyridine based fluorescent receptors for the recognition of uric acid. <i>RSC Advances</i> , 2014 , 4, 428-433	4.3	29

15	Recognition of a dicarboxylic acid with dipicolyl urea in solution and in solid phases: intramolecular hydrogen bond inhibiting both pyridine nitrogens from binding carboxyl groups. <i>Tetrahedron</i> , 2008 , 64, 6426-6433	2.4	13
14	Design and synthesis of a unique ditopic macrocyclic fluorescent receptor containing furan ring as a spacer for the recognition of dicarboxylic acids. <i>Tetrahedron</i> , 2008 , 64, 6358-6363	2.4	40
13	Tailor-made naphthyridines: Self-assembling multiple hydrogen-bonded supramolecular architectures from dimer to helix. <i>Journal of Molecular Structure</i> , 2007 , 846, 97-107	3.4	31
12	Microwave-Expedited One-Pot, Two-Component, Solvent-Free Synthesis of Functionalized Pyrimidines. <i>Australian Journal of Chemistry</i> , 2007 , 60, 120	1.2	33
11	Specificity in Crystal Engineering: Polymerisation and Dimerisation Motif Design in Pyridine System. <i>Letters in Organic Chemistry</i> , 2007 , 4, 448-451	0.6	
10	Recognition of Dicarboxylic Acid by 6,6'-Dipivaloylamino-3,3'-bipyridine and the Supramolecular Solid State Locking of the Carboxyls in the anti Form. <i>Supramolecular Chemistry</i> , 2006 , 18, 571-574	1.8	18
9	Directed molecular recognition: design and synthesis of neutral receptors for biotin to bind both its functional groups. <i>Journal of Organic Chemistry</i> , 2006 , 71, 7280-7	4.2	19
8	2-Amino-6-(hydroxymethyl)pyridine. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2006 , 62, o3225-o3227		1
7	A Simple and Convenient Manganese Dioxide Oxidation of Benzyl Halides to Aromatic Aldehydes under Neutral Condition. <i>Chemistry Letters</i> , 2005 , 34, 194-195	1.7	19
6	Directed H-bonding inhibition in molecular recognition: an NMR case study of the H-bonding of a dicarboxylic acid with a new mixed diamide receptor having one adjacent pyridine-N-oxide. <i>Tetrahedron Letters</i> , 2005 , 46, 1315-1318	2	13
5	Dynamic molecular recognition in the generation of a new crystal-engineering motif: a unique case study of a dicarboxylic acid with a ditopic receptor favouring a polymeric over a dimeric hydrogen-bonded supramolecular complex. <i>Tetrahedron Letters</i> , 2005 , 46, 7187-7191	2	23
4	A convenient palladium catalyzed synthesis of symmetric biaryls, biheterocycles and biaryl chiral diamides. <i>Tetrahedron</i> , 2005 , 61, 4289-4295	2.4	17
3	N-[6-(Hydroxymethyl)pyridin-2-yl]-2,2-dimethylpropanamide. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2005 , 61, o105-o107		2
2	Multiple hydrogen bonds and tautomerism in naphthyridine derivatives. <i>New Journal of Chemistry</i> , 2004 , 28, 700-707	3.6	40
1	Side Chain Bromination of Mono and Dimethyl Heteroaromatic and Aromatic Compounds by Solid Phase N-Bromosuccinimide Reaction without Radical Initiator under Microwave. <i>Chemistry Letters</i> , 2004 , 33, 916-917	1.7	30