

Valeria C Sandrim

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5348053/valeria-c-sandrim-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

146
papers

2,589
citations

27
h-index

44
g-index

157
ext. papers

2,901
ext. citations

3.4
avg, IF

4.88
L-index

#	Paper	IF	Citations
146	Analysis of the associations of the T-786C and Glu298Asp polymorphisms of the eNOS gene as risk factors in the rupture of intracranial aneurysms. <i>Meta Gene</i> , 2022 , 31, 101003	0.7	0
145	Potential role of uric acid to activate NLRP3 inflammasome triggering endothelial dysfunction in preeclampsia. <i>Clinical Immunology Communications</i> , 2022 , 2, 69-75		0
144	Interaction among extracellular nicotinamide phosphoribosyltransferase, toll-like receptor-4, and inflammatory cytokines in pre-eclampsia.. <i>American Journal of Reproductive Immunology</i> , 2022 , 87, e13514 ^{3,8}	3.8	14
143	Comprehensive analyses of DNA methylation of the TIMP3 promoter in placentas from early-onset and late-onset preeclampsia. <i>Placenta</i> , 2021 , 117, 118-121	3.4	0
142	NLRP3 Activation and Its Relationship to Endothelial Dysfunction and Oxidative Stress: Implications for Preeclampsia and Pharmacological Interventions. <i>Cells</i> , 2021 , 10,	7.9	7
141	Different profiles of circulating arginase 2 in subtypes of preeclampsia pregnant women. <i>Clinical Biochemistry</i> , 2021 , 92, 25-33	3.5	0
140	Effects of arginase genetic polymorphisms on nitric oxide formation in healthy pregnancy and in preeclampsia. <i>Nitric Oxide - Biology and Chemistry</i> , 2021 , 109-110, 20-25	5	1
139	single-nucleotide polymorphism rs1319501 and visfatin/NAMPT affect nitric oxide formation, sFlt-1 and antihypertensive therapy response in preeclampsia. <i>Pharmacogenomics</i> , 2021 , 22, 451-464	2.6	3
138	Circulating levels of hydrogen sulphide negatively correlate to nitrite levels in gestational hypertensive and preeclamptic pregnant women. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2021 , 48, 1224-1230	3	3
137	Circulating Total Cell-Free DNA Levels Are Increased in Hypertensive Disorders of Pregnancy and Associated with Prohypertensive Factors and Adverse Clinical Outcomes. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	2
136	Effects of vitamin D-induced supernatant of placental explants from preeclamptic women on oxidative stress and nitric oxide bioavailability in human umbilical vein endothelial cells. <i>Brazilian Journal of Medical and Biological Research</i> , 2021 , 54, e11073	2.8	0
135	Monocytes from preeclamptic women previously treated with silibinin attenuate oxidative stress in human endothelial cells. <i>Hypertension in Pregnancy</i> , 2021 , 40, 124-132	2	1
134	Resveratrol and grape juice: Effects on redox status and nitric oxide production of endothelial cells in in vitro preeclampsia model. <i>Pregnancy Hypertension</i> , 2021 , 23, 205-210	2.6	4
133	Potential roles of visfatin/NAMPT on endothelial dysfunction in preeclampsia and pathways underlying cardiac and vascular remodeling. <i>Journal of Cellular Physiology</i> , 2021 ,	7	1
132	Circulating MicroRNAs in the Second Trimester From Pregnant Women Who Subsequently Developed Preeclampsia: Potential Candidates as Predictive Biomarkers and Pathway Analysis for Target Genes of miR-204-5p. <i>Frontiers in Physiology</i> , 2021 , 12, 678184	4.6	1
131	Letter to the editor regarding: "Resveratrol and endothelial function: A literature review". <i>Pharmacological Research</i> , 2021 , 172, 105799	10.2	
130	Preeclampsia and Gestational Hypertension: Biochemical and Antioxidant Features in Vitro Might Help Understand Different Outcomes.. <i>Revista Brasileira De Ginecologia E Obstetricia</i> , 2021 , 43, 894-903 ^{1.1}		1.1

129	Effects of chronic dietary nitrate supplementation on longevity, vascular function and cancer incidence in rats.. <i>Redox Biology</i> , 2021 , 48, 102209	11.3	0
128	Analysis of SIRT1 Expression in Plasma and in an Model of Preeclampsia. <i>Oxidative Medicine and Cellular Longevity</i> , 2020 , 2020, 4561083	6.7	9
127	Interaction Between NOS3 and HMOX1 on Antihypertensive Drug Responsiveness in Preeclampsia. <i>Revista Brasileira De Ginecologia E Obstetricia</i> , 2020 , 42, 460-467	1.1	1
126	Association of Omnivorous and Vegetarian Diets With Antioxidant Defense Mechanisms in Men. <i>Journal of the American Heart Association</i> , 2020 , 9, e015576	6	6
125	Candidate genes identified by whole-exome sequencing in preeclampsia families: insights into functional annotation and in-silico prediction of deleterious variants. <i>Journal of Hypertension</i> , 2020 , 38, 372-374	1.9	1
124	SIRT1-dependent effects of resveratrol and grape juice in an in vitro model of preeclampsia. <i>Biomedicine and Pharmacotherapy</i> , 2020 , 131, 110659	7.5	6
123	Existe Alteraç em Marcadores Inflamatōos em Estudantes de Medicina apō Participaç em Programa Mente-Corpo?. <i>Revista Brasileira De Educacao Medica</i> , 2019 , 43, 79-86	0.3	0
122	Role of plasma PlGF, PDGF-AA, ANG-1, ANG-2, and the ANG-1/ANG-2 ratio as predictors of preeclampsia in a cohort of pregnant women. <i>Pregnancy Hypertension</i> , 2019 , 16, 105-111	2.6	5
121	Resveratrol improves endothelial cell markers impaired by plasma incubation from women who subsequently develop preeclampsia. <i>Hypertension Research</i> , 2019 , 42, 1166-1174	4.7	10
120	Circulating HO-1 levels are not associated with plasma sFLT-1 and GT HMOX1 polymorphism in preeclampsia. <i>Hypertension in Pregnancy</i> , 2019 , 38, 73-77	2	3
119	Consumption of animal-based and processed food associated with cardiovascular risk factors and subclinical atherosclerosis biomarkers in men. <i>Revista Da Associaō Medica Brasileira</i> , 2019 , 65, 43-50	1.4	4
118	Amino Acid Biosignature in Plasma among Ischemic Stroke Subtypes. <i>BioMed Research International</i> , 2019 , 2019, 8480468	3	10
117	C>T (rs17035945) polymorphism of TIMP-4 protects against preeclampsia. <i>Journal of Obstetrics and Gynaecology</i> , 2019 , 39, 135-137	1.3	2
116	Metabolic Disturbances Identified in Plasma Samples from ST-Segment Elevation Myocardial Infarction Patients. <i>Disease Markers</i> , 2019 , 2019, 7676189	3.2	4
115	NAMPT levels are inversely related to nitric oxide formation and positively related to soluble fms-like tyrosine kinase-1 levels in preeclampsia. <i>Pregnancy Hypertension</i> , 2019 , 18, 137-140	2.6	3
114	Shortening telomere is associated with subclinical atherosclerosis biomarker in omnivorous but not in vegetarian healthy men. <i>Aging</i> , 2019 , 11, 5070-5080	5.6	3
113	Influence of NKG2C gene deletion and CCR5Δ2 in Pre-eclampsia-Approaching the effect of innate immune gene variants in pregnancy. <i>International Journal of Immunogenetics</i> , 2019 , 46, 82-87	2.3	6
112	Haptoglobin levels are influenced by Hp1-Hp2 polymorphism, obesity, inflammation, and hypertension in type 2 diabetes mellitus. <i>Endocrinologia, Diabetes Y Nutriciō</i> , 2019 , 66, 99-107	1.3	5

111	esomeprazole to treat women with preeclampsia: possible implications in the nitric oxide homeostasis. <i>American Journal of Obstetrics and Gynecology</i> , 2019 , 220, 206-207	6.4	2
110	Levels of MMP-9 in patients with intracranial aneurysm: Relation with risk factors, size and clinical presentation. <i>Clinical Biochemistry</i> , 2018 , 55, 63-68	3.5	12
109	Preeclamptic plasma stimulates the expression of miRNAs, leading to a decrease in endothelin-1 production in endothelial cells. <i>Pregnancy Hypertension</i> , 2018 , 12, 75-81	2.6	12
108	Gain-of-function SNPs in NLRP3 and IL1B genes confer protection against obesity and T2D: undiscovered role of inflammasome genetics in metabolic homeostasis?. <i>Endocrine</i> , 2018 , 60, 368-371	4	3
107	Pharmacogenomics of Hypertension and Preeclampsia: Focus on Gene-Gene Interactions. <i>Frontiers in Pharmacology</i> , 2018 , 9, 168	5.6	12
106	Tobacco influence in heavy metals levels in head and neck cancer cases. <i>Environmental Science and Pollution Research</i> , 2018 , 25, 27650-27656	5.1	6
105	Prevalence of Metabolic Syndrome and Framingham Risk Score in Apparently Healthy Vegetarian and Omnivorous Men. <i>Arquivos Brasileiros De Cardiologia</i> , 2018 , 110, 430-437	1.2	5
104	Higher levels of circulating TIMP-4 in preeclampsia is strongly associated with clinical parameters and microRNA. <i>Clinical and Experimental Hypertension</i> , 2018 , 40, 609-612	2.2	6
103	Circulating Tissue Inhibitor of Metalloproteinase-4 levels are not a Predictor of Preeclampsia in the period between 20 and 25 Weeks of Gestation. <i>Revista Brasileira De Ginecologia E Obstetricia</i> , 2018 , 40, 757-762	1.1	1
102	Autophagy-related 16-like 1 gene polymorphism, risk factors for cardiovascular disease and associated carotid intima-media thickness in postmenopausal women. <i>Clinical Biochemistry</i> , 2018 , 61, 12-17	3.5	2
101	Circulating Heme Oxygenase-1: Not a Predictor of Preeclampsia but Highly Expressed in Pregnant Women Who Subsequently Develop Severe Preeclampsia. <i>Oxidative Medicine and Cellular Longevity</i> , 2018 , 2018, 6035868	6.7	5
100	Gene-gene interactions in the NAMPT pathway, plasma visfatin/NAMPT levels, and antihypertensive therapy responsiveness in hypertensive disorders of pregnancy. <i>Pharmacogenomics Journal</i> , 2017 , 17, 427-434	3.5	18
99	Myeloperoxidase in Hypertensive Disorders of Pregnancy and Its Relation With Nitric Oxide. <i>Hypertension</i> , 2017 , 69, 1173-1180	8.5	13
98	Pharmacogenetics in the treatment of pre-eclampsia: current findings, challenges and perspectives. <i>Pharmacogenomics</i> , 2017 , 18, 571-583	2.6	15
97	Myeloperoxidase is not a good biomarker for preeclampsia prediction. <i>Scientific Reports</i> , 2017 , 7, 10257	4.9	6
96	Decoding resistant hypertension signalling pathways. <i>Clinical Science</i> , 2017 , 131, 2813-2834	6.5	7
95	IL-6, TNF- α and IL-10 levels/polymorphisms and their association with type 2 diabetes mellitus and obesity in Brazilian individuals. <i>Archives of Endocrinology and Metabolism</i> , 2017 , 61, 438-446	2.2	52
94	Childhood Obesity, MMP-9 Levels, and Vitamin D. <i>Arquivos Brasileiros De Cardiologia</i> , 2017 , 109, 380-381	1.2	

93	Longitudinal assessment of maternal-fetal Doppler parameters and maternal plasma level of matrix metalloproteinases 2 and 9. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2016 , 29, 3967-70	2	2
92	Plasma levels of increased miR-195-5p correlates with the sFLT-1 levels in preeclampsia. <i>Hypertension in Pregnancy</i> , 2016 , 35, 150-8	2	12
91	Assessment of nitrite oxide and maternal-fetal Doppler parameters during pregnancy. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2016 , 29, 3406-9	2	
90	Antihypertensive therapy in pre-eclampsia: effects of plasma from nonresponsive patients on endothelial gene expression. <i>Pharmacogenomics</i> , 2016 , 17, 1121-1127	2.6	7
89	Circulating microRNA expression profiles in pre-eclampsia: evidence of increased miR-885-5p levels. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016 , 123, 2120-2128	3.7	42
88	Role of adiponectin on antioxidant profile: evaluation during healthy and hypertensive disorders of pregnancy. <i>Blood Pressure</i> , 2016 , 25, 241-3	1.7	8
87	Plasma matrix metalloproteinase-9 levels, MMP-9 gene haplotypes, and cardiovascular risk in obese subjects. <i>Molecular Biology Reports</i> , 2016 , 43, 463-71	2.8	11
86	Lack of association between genetic polymorphism of FTO, AKT1 and AKTIP in childhood overweight and obesity. <i>Jornal De Pediatria (Versão Em Português)</i> , 2016 , 92, 521-527	0.2	2
85	Reduced levels of potential circulating biomarkers of cardiovascular diseases in apparently healthy vegetarian men. <i>Clinica Chimica Acta</i> , 2016 , 461, 110-3	6.2	10
84	Lack of association between genetic polymorphism of FTO, AKT1 and AKTIP in childhood overweight and obesity. <i>Jornal De Pediatria</i> , 2016 , 92, 521-7	2.6	11
83	Plasma from pre-eclamptic patients induces the expression of the anti-angiogenic miR-195-5p in endothelial cells. <i>Journal of Cellular and Molecular Medicine</i> , 2016 , 20, 1198-200	5.6	25
82	Antihypertensive therapy in preeclampsia is not modulated by VEGF polymorphisms. <i>Archives of Gynecology and Obstetrics</i> , 2015 , 291, 799-803	2.5	5
81	Positive correlations between circulating adiponectin and MMP2 in preeclampsia pregnant. <i>Pregnancy Hypertension</i> , 2015 , 5, 205-8	2.6	34
80	Simvastatin does not reduce chemokine production in obesity without comorbidities. <i>Inflammation</i> , 2015 , 38, 1297-301	5.1	2
79	Effect of acetylsalicylic acid on platelet activation and oxidative profile in a set of Brazilian patients with type 2 diabetes mellitus. <i>Blood Coagulation and Fibrinolysis</i> , 2015 , 26, 123-30	1	17
78	NLRP1 L155H Polymorphism is a Risk Factor for Preeclampsia Development. <i>American Journal of Reproductive Immunology</i> , 2015 , 73, 577-81	3.8	25
77	Visfatin levels are decreased in advanced stages of diabetic nephropathy. <i>Renal Failure</i> , 2015 , 37, 1529-30	0.9	1
76	Correlations among antiangiogenic factors and trace elements in hypertensive disorders of pregnancy. <i>Journal of Trace Elements in Medicine and Biology</i> , 2015 , 29, 130-5	4.1	12

75	An update on the pharmacogenetics of treating hypertension. <i>Journal of Human Hypertension</i> , 2015 , 29, 283-91	2.6	26
74	Endocan: a new biomarker associated with inflammation in type 2 diabetes mellitus?. <i>Diabetes/Metabolism Research and Reviews</i> , 2015 , 31, 479-80	7.5	11
73	Role of MMP-2 and MMP-9 in resistance to drug therapy in patients with resistant hypertension. <i>Arquivos Brasileiros De Cardiologia</i> , 2015 , 105, 168-75	1.2	4
72	Association of a Large Panel of Cytokine Gene Polymorphisms with Complications and Comorbidities in Type 2 Diabetes Patients. <i>Journal of Diabetes Research</i> , 2015 , 2015, 605965	3.9	29
71	Effects of NAMPT polymorphisms and haplotypes on circulating visfatin/NAMPT levels in hypertensive disorders of pregnancy. <i>Hypertension Research</i> , 2015 , 38, 361-6	4.7	18
70	Hypertension and Vascular Endothelial Growth Factors 2015 , 695-707		1
69	Evaluation of clinical and inflammatory markers of nonalcoholic fatty liver disease in postmenopausal women with metabolic syndrome. <i>Metabolic Syndrome and Related Disorders</i> , 2014 , 12, 330-8	2.6	6
68	Polymorphisms in endothelial nitric oxide synthase gene in early and late severe preeclampsia. <i>Nitric Oxide - Biology and Chemistry</i> , 2014 , 42, 19-23	5	21
67	Tissue inhibitor of matrix metalloproteinase-1 polymorphism, plasma TIMP-1 levels, and antihypertensive therapy responsiveness in hypertensive disorders of pregnancy. <i>Pharmacogenomics Journal</i> , 2014 , 14, 535-41	3.5	26
66	Reduced circulating miR-196b levels is associated with preeclampsia. <i>Pregnancy Hypertension</i> , 2014 , 4, 11-3	2.6	7
65	Homocysteine and nitrite levels are modulated by MTHFR 677C>T polymorphism in obese women treated with simvastatin. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2014 , 41, 744-7	3	5
64	Polymorphisms of the adiponectin gene in gestational hypertension and pre-eclampsia. <i>Journal of Human Hypertension</i> , 2014 , 28, 128-32	2.6	11
63	Effect of simvastatin treatment on plasma visfatin levels in obese women. <i>Gynecological Endocrinology</i> , 2014 , 30, 577-80	2.4	
62	Correlations between circulating levels of adipokines and anti-angiogenic factors in women with BMI. <i>Hypertension in Pregnancy</i> , 2014 , 33, 72-80	2	11
61	Relationship between adiponectin and nitrite in healthy and preeclampsia pregnancies. <i>Clinica Chimica Acta</i> , 2013 , 423, 112-5	6.2	21
60	Haptoglobin polymorphism affects nitric oxide bioavailability in preeclampsia. <i>Journal of Human Hypertension</i> , 2013 , 27, 349-54	2.6	14
59	Simvastatin treatment increases nitrite levels in obese women: modulation by T(-786)C polymorphism of eNOS. <i>Nitric Oxide - Biology and Chemistry</i> , 2013 , 33, 83-7	5	15
58	Assessment of oxidative status markers and NO bioavailability in hypertensive disorders of pregnancy. <i>Journal of Human Hypertension</i> , 2013 , 27, 345-8	2.6	12

57	Pharmacogenomic approaches that may guide preeclampsia therapy. <i>Pharmacogenomics</i> , 2013 , 14, 591-3.6	10
56	Maternal flow-mediated dilation and nitrite concentration during third trimester of pregnancy and postpartum period. <i>Hypertension in Pregnancy</i> , 2013 , 32, 225-34	2 7
55	Simvastatin therapy decreases MMP-9 levels in obese women. <i>Journal of Clinical Pharmacology</i> , 2013 , 53, 1072-7	2.9 14
54	Functional VEGF haplotypes affect the susceptibility to hypertension. <i>Journal of Human Hypertension</i> , 2013 , 27, 31-7	2.6 10
53	Effects of matrix metalloproteinase (MMP)-2 polymorphisms on responsiveness to antihypertensive therapy of women with hypertensive disorders of pregnancy. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2012 , 111, 262-7	3.1 16
52	Association between matrix metalloproteinase (MMP)-2 polymorphisms and MMP-2 levels in hypertensive disorders of pregnancy. <i>Experimental and Molecular Pathology</i> , 2012 , 92, 217-21	4.4 48
51	Endogenous nitric oxide formation correlates negatively with circulating matrix metalloproteinase (MMP)-2 and MMP-9 levels in black subjects. <i>Molecular and Cellular Biochemistry</i> , 2012 , 360, 393-9	4.2 11
50	Functional polymorphism located in MMP-9 gene promoter is strongly associated with obesity. <i>DNA and Cell Biology</i> , 2012 , 31, 1054-7	3.6 13
49	Evaluation of plasmatic MMP-8, MMP-9, TIMP-1 and MPO levels in obese and lean women. <i>Clinical Biochemistry</i> , 2012 , 45, 412-5	3.5 39
48	Maternal iNOS genetic polymorphisms and hypertensive disorders of pregnancy. <i>Journal of Human Hypertension</i> , 2012 , 26, 547-52	2.6 17
47	Matrix metalloproteinase-9 polymorphisms affect plasma MMP-9 levels and antihypertensive therapy responsiveness in hypertensive disorders of pregnancy. <i>Pharmacogenomics Journal</i> , 2012 , 12, 489-98	3.5 38
46	Endothelial FOS expression and pre-eclampsia. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2012 , 119, 1564-71	3.7 15
45	Epistasis among eNOS, MMP-9 and VEGF maternal genotypes in hypertensive disorders of pregnancy. <i>Hypertension Research</i> , 2012 , 35, 917-21	4.7 25
44	Functional MMP-9 polymorphisms modulate plasma MMP-9 levels in multiple sclerosis patients. <i>Journal of Neuroimmunology</i> , 2012 , 249, 56-9	3.5 15
43	eNOS tag SNP haplotypes in hypertensive disorders of pregnancy. <i>DNA and Cell Biology</i> , 2012 , 31, 1665-306	16
42	Vitamin D receptor polymorphisms in hypertensive disorders of pregnancy. <i>Molecular Biology Reports</i> , 2012 , 39, 10903-6	2.8 18
41	4G/5G polymorphism modulates PAI-1 circulating levels in obese women. <i>Molecular and Cellular Biochemistry</i> , 2012 , 364, 299-301	4.2 7
40	Polymorphisms and haplotypes in candidate genes related to angiogenesis and endothelial dysfunction in preeclampsia. <i>Journal of Pregnancy</i> , 2012 , 2012, 914704	2.5 4

39	Negative correlation between D-dimer and plasminogen activator inhibitor-1 levels is absent in obese women. <i>Blood Coagulation and Fibrinolysis</i> , 2012 , 23, 402-5	1	2
38	Importance of haplotype analysis in association studies considering VEGF promoter polymorphisms. <i>Clinical Biochemistry</i> , 2011 , 44, 747; author reply 748	3.5	2
37	Alterations in cyclic GMP levels in preeclampsia may reflect increased B-type natriuretic peptide levels and not impaired nitric oxide activity. <i>Clinical Biochemistry</i> , 2011 , 44, 1012-4	3.5	14
36	eNOS haplotypes affect the responsiveness to antihypertensive therapy in preeclampsia but not in gestational hypertension. <i>Pharmacogenomics Journal</i> , 2010 , 10, 40-5	3.5	56
35	Effects of eNOS polymorphisms on nitric oxide formation in healthy pregnancy and in pre-eclampsia. <i>Molecular Human Reproduction</i> , 2010 , 16, 506-10	4.4	51
34	Matrix metalloproteinase (MMP)-9 genotypes and haplotypes in preeclampsia and gestational hypertension. <i>Clinica Chimica Acta</i> , 2010 , 411, 874-7	6.2	35
33	Interethnic differences in ADMA concentrations and negative association with nitric oxide formation in preeclampsia. <i>Clinica Chimica Acta</i> , 2010 , 411, 1457-60	6.2	28
32	Increased circulating cell-free hemoglobin levels reduce nitric oxide bioavailability in preeclampsia. <i>Free Radical Biology and Medicine</i> , 2010 , 49, 493-500	7.8	41
31	Interethnic differences in the distribution of clinically relevant vascular endothelial growth factor genetic polymorphisms. <i>DNA and Cell Biology</i> , 2009 , 28, 567-72	3.6	22
30	Positive correlations between serum and plasma matrix metalloproteinase (MMP)-2 or MMP-9 levels in disease conditions. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 888-91	5.9	28
29	T allele of -344 C/T polymorphism in aldosterone synthase gene is not associated with resistant hypertension. <i>Hypertension Research</i> , 2009 , 32, 159-62	4.7	10
28	Matrix metalloproteinase-9 genotypes and haplotypes are associated with multiple sclerosis and with the degree of disability of the disease. <i>Journal of Neuroimmunology</i> , 2009 , 214, 128-31	3.5	23
27	Aldosterone synthase gene polymorphism is not associated with gestational hypertension or preeclampsia. <i>Clinica Chimica Acta</i> , 2009 , 400, 139-41	6.2	10
26	Vascular endothelial growth factor genotypes and haplotypes are associated with pre-eclampsia but not with gestational hypertension. <i>Molecular Human Reproduction</i> , 2009 , 15, 115-20	4.4	53
25	eNOS haplotypes associated with gestational hypertension or preeclampsia. <i>Pharmacogenomics</i> , 2008 , 9, 1467-73	2.6	75
24	Nitric oxide formation is inversely related to serum levels of antiangiogenic factors soluble fms-like tyrosine kinase-1 and soluble endogline in preeclampsia. <i>Hypertension</i> , 2008 , 52, 402-7	8.5	128
23	Haplotypes of vitamin D receptor modulate the circulating levels of lead in exposed subjects. <i>Archives of Toxicology</i> , 2008 , 82, 29-36	5.8	36
22	Evidence of thermostable amylolytic activity from <i>Rhizopus microsporus</i> var. <i>rhizopodiformis</i> using wheat bran and corncob as alternative carbon source. <i>Bioprocess and Biosystems Engineering</i> , 2008 , 31, 329-34	3.7	17

21	Comparative assessment of matrix metalloproteinase (MMP)-2 and MMP-9, and their inhibitors, tissue inhibitors of metalloproteinase (TIMP)-1 and TIMP-2 in preeclampsia and gestational hypertension. <i>Clinical Biochemistry</i> , 2008 , 41, 875-80	3.5	84
20	Haplotype analysis can provide improved clinical information than single genotype analysis. <i>Thrombosis Research</i> , 2007 , 120, 779	8.2	7
19	Influence of eNOS haplotypes on the plasma nitric oxide products concentrations in hypertensive and type 2 diabetes mellitus patients. <i>Nitric Oxide - Biology and Chemistry</i> , 2007 , 16, 348-55	5	75
18	An interethnic comparison of the distribution of vitamin D receptor genotypes and haplotypes. <i>Clinica Chimica Acta</i> , 2007 , 384, 155-9	6.2	21
17	Endothelial nitric oxide synthase polymorphisms and hypertension: improved clinical evidence derived from haplotype analysis. <i>International Journal of Cardiology</i> , 2007 , 116, 116	3.2	6
16	Anti-inflammatory effects of atorvastatin: modulation by the T-786C polymorphism in the endothelial nitric oxide synthase gene. <i>Atherosclerosis</i> , 2007 , 193, 438-44	3.1	69
15	A polymorphism in the delta-aminolevulinic acid dehydratase gene modifies plasma/whole blood lead ratio. <i>Archives of Toxicology</i> , 2006 , 80, 394-8	5.8	35
14	Regarding "eNOS G894T polymorphism as a mild predisposing factor for abdominal aortic aneurysm". <i>Journal of Vascular Surgery</i> , 2006 , 43, 1079	3.5	1
13	Susceptible and protective eNOS haplotypes in hypertensive black and white subjects. <i>Atherosclerosis</i> , 2006 , 186, 428-32	3.1	89
12	Endothelial nitric oxide synthase haplotypes affect the susceptibility to hypertension in patients with type 2 diabetes mellitus. <i>Atherosclerosis</i> , 2006 , 189, 241-6	3.1	71
11	Influence of T-786C polymorphism on the promoter activity of eNOS. <i>Clinica Chimica Acta</i> , 2006 , 367, 208	6.2	3
10	Considerations about functional implications of Glu298Asp polymorphism of endothelial nitric oxide synthase gene. <i>Clinica Chimica Acta</i> , 2006 , 367, 207	6.2	
9	Ethnicity affects the distribution of delta-aminolevulinic acid dehydratase (ALAD) genetic variants. <i>Clinica Chimica Acta</i> , 2006 , 367, 192-5	6.2	18
8	eNOS genotype-dependent correlation between whole blood lead and plasma nitric oxide products concentrations. <i>Nitric Oxide - Biology and Chemistry</i> , 2006 , 14, 58-64	5	27
7	Endothelial nitric oxide synthase genotype and haplotype are not associated with diabetic retinopathy in diabetes type 2 patients. <i>Nitric Oxide - Biology and Chemistry</i> , 2006 , 15, 417-22	5	25
6	FARMACOGENÉTICA CARDIOVASCULAR. <i>Medicina</i> , 2006 , 39, 535	0.1	
5	Endothelial nitric oxide synthase haplotypes are related to blood pressure elevation, but not to resistance to antihypertensive drug therapy. <i>Journal of Hypertension</i> , 2006 , 24, 2393-7	1.9	47
4	Screening of filamentous fungi for production of enzymes of biotechnological interest. <i>Brazilian Journal of Microbiology</i> , 2006 , 37, 474-480	2.2	64

3	Consistent interethnic differences in the distribution of clinically relevant endothelial nitric oxide synthase genetic polymorphisms. <i>Nitric Oxide - Biology and Chemistry</i> , 2005 , 12, 177-82	5	136
2	Purification and biochemical characterization of two xylanases produced by <i>Aspergillus caespitosus</i> and their potential for kraft pulp bleaching. <i>Process Biochemistry</i> , 2005 , 40, 1823-1828	4.8	77
1	Influence of temperature on the properties of the xylanolytic enzymes of the thermotolerant fungus <i>Aspergillus phoenicis</i> . <i>Journal of Industrial Microbiology and Biotechnology</i> , 2004 , 31, 88-93	4.2	22