

Wilma van de Berg

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5351298/publications.pdf>

Version: 2024-02-01

81
papers

4,747
citations

94433

37
h-index

106344

65
g-index

85
all docs

85
docs citations

85
times ranked

7746
citing authors

#	ARTICLE	IF	CITATIONS
1	Lewy pathology in Parkinson's disease consists of crowded organelles and lipid membranes. <i>Nature Neuroscience</i> , 2019, 22, 1099-1109.	14.8	604
2	Therapeutic potential of autophagy-enhancing agents in Parkinson's disease. <i>Molecular Neurodegeneration</i> , 2017, 12, 11.	10.8	211
3	Oligomeric and phosphorylated alpha-synuclein as potential CSF biomarkers for Parkinson's disease. <i>Molecular Neurodegeneration</i> , 2016, 11, 7.	10.8	198
4	Prenatal stress and neonatal rat brain development. <i>Neuroscience</i> , 2006, 137, 145-155.	2.3	173
5	Microglial phenotypes and toll-like receptor 2 in the substantia nigra and hippocampus of incidental Lewy body disease cases and Parkinson's disease patients. <i>Acta Neuropathologica Communications</i> , 2014, 2, 90.	5.2	140
6	Longterm quiescent cells in the aged human subventricular neurogenic system specifically express GFAP. <i>Aging Cell</i> , 2010, 9, 313-326.	6.7	126
7	Lysosomal Dysfunction and α -Synuclein Aggregation in Parkinson's Disease: Diagnostic Links. <i>Movement Disorders</i> , 2016, 31, 791-801.	3.9	125
8	Stage-dependent nigral neuronal loss in incidental Lewy body and Parkinson's disease. <i>Movement Disorders</i> , 2014, 29, 1244-1251.	3.9	122
9	Generation and characterization of novel conformation-specific monoclonal antibodies for α -synuclein pathology. <i>Neurobiology of Disease</i> , 2015, 79, 81-99.	4.4	116
10	Evidence for Immune Response, Axonal Dysfunction and Reduced Endocytosis in the Substantia Nigra in Early Stage Parkinson's Disease. <i>PLoS ONE</i> , 2015, 10, e0128651.	2.5	114
11	Topographic Mapping between Basal Forebrain Cholinergic Neurons and the Medial Prefrontal Cortex in Mice. <i>Journal of Neuroscience</i> , 2014, 34, 16234-16246.	3.6	112
12	Distribution and Load of Amyloid- β Pathology in Parkinson Disease and Dementia with Lewy Bodies. <i>Journal of Neuropathology and Experimental Neurology</i> , 2016, 75, 936-945.	1.7	109
13	The proliferative capacity of the subventricular zone is maintained in the parkinsonian brain. <i>Brain</i> , 2011, 134, 3249-3263.	7.6	103
14	LRP10 genetic variants in familial Parkinson's disease and dementia with Lewy bodies: a genome-wide linkage and sequencing study. <i>Lancet Neurology</i> , The, 2018, 17, 597-608.	10.2	101
15	Characterization of Brain Lysosomal Activities in GBA-Related and Sporadic Parkinson's Disease and Dementia with Lewy Bodies. <i>Molecular Neurobiology</i> , 2019, 56, 1344-1355.	4.0	97
16	Changes in endolysosomal enzyme activities in cerebrospinal fluid of patients with Parkinson's disease. <i>Movement Disorders</i> , 2013, 28, 747-754.	3.9	88
17	CSF α -Synuclein Does Not Discriminate Dementia with Lewy Bodies from Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2010, 22, 87-95.	2.6	87
18	Low-density lipoprotein receptor-knockout mice display impaired spatial memory associated with a decreased synaptic density in the hippocampus. <i>Neurobiology of Disease</i> , 2004, 16, 212-219.	4.4	84

#	ARTICLE	IF	CITATIONS
19	Î±-Synuclein species as potential cerebrospinal fluid biomarkers for dementia with lewy bodies. <i>Movement Disorders</i> , 2018, 33, 1724-1733.	3.9	79
20	Increased Amoeboid Microglial Density in the Olfactory Bulb of Parkinson's and Alzheimer's Patients. <i>Brain Pathology</i> , 2014, 24, 152-165.	4.1	70
21	CSF or Serum Neurofilament Light Added to Î±-Synuclein Panel Discriminates Parkinson's From Controls. <i>Movement Disorders</i> , 2020, 35, 288-295.	3.9	69
22	Diagnostic cerebrospinal fluid biomarkers for Parkinson's disease: A pathogenetically based approach. <i>Neurobiology of Disease</i> , 2010, 39, 229-241.	4.4	67
23	Reduced Î±-Synuclein levels in cerebrospinal fluid in Parkinson's disease are unrelated to clinical and imaging measures of disease severity. <i>European Journal of Neurology</i> , 2014, 21, 388-394.	3.3	67
24	Prefrontal cortical ChAT-VIP interneurons provide local excitation by cholinergic synaptic transmission and control attention. <i>Nature Communications</i> , 2019, 10, 5280.	12.8	65
25	The subcellular arrangement of alpha-synuclein proteoforms in the Parkinson's disease brain as revealed by multicolor STED microscopy. <i>Acta Neuropathologica</i> , 2021, 142, 423-448.	7.7	65
26	Transcriptomic signatures of brain regional vulnerability to Parkinson's disease. <i>Communications Biology</i> , 2020, 3, 101.	4.4	58
27	Impact of perinatal asphyxia on the GABAergic and locomotor system. <i>Neuroscience</i> , 2003, 117, 83-96.	2.3	57
28	Clusterin Levels in Plasma Predict Cognitive Decline and Progression to Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2015, 46, 1103-1110.	2.6	55
29	Patterns of alpha-synuclein pathology in incidental cases and clinical subtypes of Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , 2012, 18, S28-S30.	2.2	54
30	The Proteome of the Locus Ceruleus in Parkinson's Disease: Relevance to Pathogenesis. <i>Brain Pathology</i> , 2012, 22, 485-498.	4.1	53
31	Increased levels of CSF total but not oligomeric or phosphorylated forms of alpha-synuclein in patients diagnosed with probable Alzheimer's disease. <i>Scientific Reports</i> , 2017, 7, 40263.	3.3	51
32	A comprehensive analysis of SNCA-related genetic risk in sporadic parkinson disease. <i>Annals of Neurology</i> , 2018, 84, 117-129.	5.3	50
33	Perinatal Asphyxia Induced Neuronal Loss by Apoptosis in the Neonatal Rat Striatum: A Combined TUNEL and Stereological Study. <i>Experimental Neurology</i> , 2002, 174, 29-36.	4.1	47
34	The coarse-grained plaque: a divergent A β plaque-type in early-onset Alzheimer's disease. <i>Acta Neuropathologica</i> , 2020, 140, 811-830.	7.7	45
35	A delayed increase in hippocampal proliferation following global asphyxia in the neonatal rat. <i>Developmental Brain Research</i> , 2003, 142, 67-76.	1.7	44
36	Loss of Functional Connectivity in Patients with Parkinson Disease and Visual Hallucinations. <i>Radiology</i> , 2017, 285, 896-903.	7.3	44

#	ARTICLE	IF	CITATIONS
37	Perinatal asphyxia results in changes in presynaptic bouton number in striatum and cerebral cortex—a stereological and behavioral analysis. <i>Journal of Chemical Neuroanatomy</i> , 2000, 20, 71-82.	2.1	43
38	Path mediation analysis reveals GBA impacts Lewy body disease status by increasing α -synuclein levels. <i>Neurobiology of Disease</i> , 2019, 121, 205-213.	4.4	43
39	A Large-Scale Full GBA1 Gene Screening in Parkinson's Disease in the Netherlands. <i>Movement Disorders</i> , 2020, 35, 1667-1674.	3.9	41
40	LRRK2 levels and phosphorylation in Parkinson's disease brain and cases with restricted Lewy bodies. <i>Movement Disorders</i> , 2017, 32, 423-432.	3.9	39
41	Pedunculopontine Cholinergic Cell Loss in Hallucinating Parkinson Disease Patients but Not in Dementia With Lewy Bodies Patients. <i>Journal of Neuropathology and Experimental Neurology</i> , 2013, 72, 1162-1170.	1.7	38
42	Axonal degeneration as substrate of fractional anisotropy abnormalities in multiple sclerosis cortex. <i>Brain</i> , 2019, 142, 1921-1937.	7.6	38
43	No alterations of hippocampal neuronal number and synaptic bouton number in a transgenic mouse model expressing the β -cleaved C-terminal APP fragment. <i>Neurobiology of Disease</i> , 2003, 12, 110-120.	4.4	37
44	A non-cholinergic neuronal loss in the pedunculopontine nucleus of toxin-evoked Parkinsonian rats. <i>Experimental Neurology</i> , 2013, 248, 213-223.	4.1	36
45	Damaged fiber tracts of the nucleus basalis of Meynert in Parkinson's disease patients with visual hallucinations. <i>Scientific Reports</i> , 2017, 7, 10112.	3.3	36
46	Differential insular cortex subregional vulnerability to α -synuclein pathology in Parkinson's disease and dementia with Lewy bodies. <i>Neuropathology and Applied Neurobiology</i> , 2019, 45, 262-277.	3.2	36
47	Differential insular cortex sub-regional atrophy in neurodegenerative diseases: a systematic review and meta-analysis. <i>Brain Imaging and Behavior</i> , 2020, 14, 2799-2816.	2.1	36
48	Cognitive correlates of visual hallucinations in non-demented Parkinson's disease patients. <i>Parkinsonism and Related Disorders</i> , 2013, 19, 795-799.	2.2	33
49	CSF total and oligomeric α -Synuclein along with TNF- α as risk biomarkers for Parkinson's disease: a study in LRRK2 mutation carriers. <i>Translational Neurodegeneration</i> , 2020, 9, 15.	8.0	32
50	An update on the genetics of dementia with Lewy bodies. <i>Parkinsonism and Related Disorders</i> , 2017, 43, 1-8.	2.2	31
51	Regional differences in gene expression and promoter usage in aged human brains. <i>Neurobiology of Aging</i> , 2013, 34, 1825-1836.	3.1	30
52	Transcriptome and proteome profiling of neural stem cells from the human subventricular zone in Parkinson's disease. <i>Acta Neuropathologica Communications</i> , 2019, 7, 84.	5.2	28
53	7T MRI allows detection of disturbed cortical lamination of the medial temporal lobe in patients with Alzheimer's disease. <i>NeuroImage: Clinical</i> , 2019, 21, 101665.	2.7	28
54	Neuropathological correlates of parkinsonian disorders in a large Dutch autopsy series. <i>Acta Neuropathologica Communications</i> , 2020, 8, 39.	5.2	28

#	ARTICLE	IF	CITATIONS
55	Morphometric Changes in the Cortical Microvascular Network in Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2010, 22, 811-818.	2.6	26
56	Cerebrospinal fluid and plasma clusterin levels in Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , 2013, 19, 1079-1083.	2.2	26
57	Normal Aging Brain Collection Amsterdam (NABCA): A comprehensive collection of postmortem high-field imaging, neuropathological and morphometric datasets of non-neurological controls. <i>NeuroImage: Clinical</i> , 2019, 22, 101698.	2.7	25
58	Use of cryostat sections from snap-frozen nervous tissue for combining stereological estimates with histological, cellular, or molecular analyses on adjacent sections. <i>Journal of Chemical Neuroanatomy</i> , 2000, 20, 21-29.	2.1	23
59	Cerebral Corpora amylacea are dense membranous labyrinths containing structurally preserved cell organelles. <i>Scientific Reports</i> , 2018, 8, 18046.	3.3	21
60	c-Jun/AP-1 (N) directed antibodies cross-react with α -apoptosis-specific protein which marks an autophagic process during neuronal apoptosis. <i>Neuroscience</i> , 2000, 96, 445-446.	2.3	20
61	Imaging of post-mortem human brain tissue using electron and X-ray microscopy. <i>Current Opinion in Structural Biology</i> , 2019, 58, 138-148.	5.7	20
62	Anterior insular network disconnection and cognitive impairment in Parkinson's disease. <i>NeuroImage: Clinical</i> , 2020, 28, 102364.	2.7	20
63	Post-Mortem MRI and Histopathology in Neurologic Disease: A Translational Approach. <i>Neuroscience Bulletin</i> , 2019, 35, 229-243.	2.9	18
64	Can post-mortem MRI be used as a proxy for in vivo? A case study. <i>Brain Communications</i> , 2019, 1, fcz030.	3.3	17
65	Relationship between β -amyloid and structural network topology in decedents without dementia. <i>Neurology</i> , 2020, 95, e532-e544.	1.1	17
66	Contactin-1 Is Reduced in Cerebrospinal Fluid of Parkinson's Disease Patients and Is Present within Lewy Bodies. <i>Biomolecules</i> , 2020, 10, 1177.	4.0	14
67	Cortical axonal loss is associated with both gray matter demyelination and white matter tract pathology in progressive multiple sclerosis: Evidence from a combined MRI-histopathology study. <i>Multiple Sclerosis Journal</i> , 2021, 27, 380-390.	3.0	13
68	Developmental apoptosis in the spinal cord white matter in neonatal rats. <i>Glia</i> , 2002, 37, 89-91.	4.9	12
69	A lathe system for micrometre-sized cylindrical sample preparation at room and cryogenic temperatures. <i>Journal of Synchrotron Radiation</i> , 2020, 27, 472-476.	2.4	12
70	Neuropathological and genetic characteristics of a post-mortem series of cases with dementia with Lewy bodies clinically suspected of Creutzfeldt-Jakob's disease. <i>Parkinsonism and Related Disorders</i> , 2019, 63, 162-168.	2.2	11
71	The adult human subventricular zone: partial ependymal coverage and proliferative capacity of cerebrospinal fluid. <i>Brain Communications</i> , 2020, 2, fcaa150.	3.3	10
72	Imaging hippocampal subregions with in vivo MRI: advances and limitations. <i>Nature Reviews Neuroscience</i> , 2012, 13, 70-70.	10.2	9

#	ARTICLE	IF	CITATIONS
73	Nitric Oxide Production in the Striatum and Cerebellum of a Rat Model of Preterm Global Perinatal Asphyxia. <i>Neurotoxicity Research</i> , 2017, 31, 400-409.	2.7	8
74	Origin of β -mannosidase activity in CSF. <i>International Journal of Biochemistry and Cell Biology</i> , 2017, 87, 34-37.	2.8	7
75	CSF Biomarkers Reflecting Protein Pathology and Axonal Degeneration Are Associated with Memory, Attentional, and Executive Functioning in Early-Stage Parkinson's Disease. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8519.	4.1	7
76	Clinical and Pathological Phenotypes of LRP10 Variant Carriers with Dementia. <i>Journal of Alzheimer's Disease</i> , 2020, 76, 1161-1170.	2.6	7
77	Dementia With Lewy Bodies. <i>Alzheimer Disease and Associated Disorders</i> , 2020, 34, 178-182.	1.3	5
78	Cingulate networks associated with gray matter loss in Parkinson's disease show high expression of cholinergic genes in the healthy brain. <i>European Journal of Neuroscience</i> , 2021, 53, 3727-3739.	2.6	5
79	High-Intensity Interval Cycle Ergometer Training in Parkinson's Disease: Protocol for Identifying Individual Response Patterns Using a Single-Subject Research Design. <i>Frontiers in Neurology</i> , 2020, 11, 569880.	2.4	4
80	Reply: Quantitative evaluation of the human subventricular zone. <i>Brain</i> , 2012, 135, e222-e222.	7.6	2
81	Alterations in Sub-Axonal Architecture Between Normal Aging and Parkinson's Diseased Human Brains Using Label-Free Cryogenic X-ray Nanotomography. <i>Frontiers in Neuroscience</i> , 2020, 14, 570019.	2.8	2