

Agnes Jamin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5368006/publications.pdf>

Version: 2024-02-01

19
papers

966
citations

623734

14
h-index

794594

19
g-index

20
all docs

20
docs citations

20
times ranked

1380
citing authors

#	ARTICLE	IF	CITATIONS
1	New insights in the pathogenesis of immunoglobulin A vasculitis (Henoch-Schönlein purpura). <i>Autoimmunity Reviews</i> , 2017, 16, 1246-1253.	5.8	228
2	Transglutaminase is essential for IgA nephropathy development acting through IgA receptors. <i>Journal of Experimental Medicine</i> , 2012, 209, 793-806.	8.5	145
3	Maternal stress during late gestation has moderate but long-lasting effects on the immune system of the piglets. <i>Veterinary Immunology and Immunopathology</i> , 2009, 131, 17-24.	1.2	70
4	Biomarkers of IgA vasculitis nephritis in children. <i>PLoS ONE</i> , 2017, 12, e0188718.	2.5	63
5	Dietary Protein Excess during Neonatal Life Alters Colonic Microbiota and Mucosal Response to Inflammatory Mediators Later in Life in Female Pigs. <i>Journal of Nutrition</i> , 2013, 143, 1225-1232.	2.9	53
6	Classical swine fever virus induces activation of plasmacytoid and conventional dendritic cells in tonsil, blood, and spleen of infected pigs. <i>Veterinary Research</i> , 2008, 39, 07.	3.0	52
7	Autoantibodies against podocytic UCHL1 are associated with idiopathic nephrotic syndrome relapses and induce proteinuria in mice. <i>Journal of Autoimmunity</i> , 2018, 89, 149-161.	6.5	48
8	Characterization of conventional and plasmacytoid dendritic cells in swine secondary lymphoid organs and blood. <i>Veterinary Immunology and Immunopathology</i> , 2006, 114, 224-237.	1.2	47
9	The Level of Protein in Milk Formula Modifies Ileal Sensitivity to LPS Later in Life in a Piglet Model. <i>PLoS ONE</i> , 2011, 6, e19594.	2.5	46
10	IgA1 Protease Treatment Reverses Mesangial Deposits and Hematuria in a Model of IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2016, 27, 2622-2629.	6.1	44
11	Value of biomarkers for predicting immunoglobulin A vasculitis nephritis outcome in an adult prospective cohort. <i>Nephrology Dialysis Transplantation</i> , 2017, 33, 1579-1590.	0.7	37
12	Fatal Effects of a Neonatal High-Protein Diet in Low-Birth-Weight Piglets Used as a Model of Intrauterine Growth Restriction. <i>Neonatology</i> , 2010, 97, 321-328.	2.0	33
13	Idiopathic nephrotic syndrome: the EBV hypothesis. <i>Pediatric Research</i> , 2017, 81, 233-239.	2.3	31
14	A high-protein neonatal formula induces a temporary reduction of adiposity and changes later adipocyte physiology. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2011, 300, R387-R397.	1.8	18
15	Recruitment of CXCR3+ T cells into injured tissues in adult IgA vasculitis patients correlates with disease activity. <i>Journal of Autoimmunity</i> , 2019, 99, 73-80.	6.5	16
16	A high-protein formula increases colonic peptide transporter 1 activity during neonatal life in low-birth-weight piglets and disturbs barrier function later in life. <i>British Journal of Nutrition</i> , 2014, 112, 1073-1080.	2.3	11
17	Accelerated Growth Rate Induced by Neonatal High-Protein Milk Formula Is Not Supported by Increased Tissue Protein Synthesis in Low-Birth-Weight Piglets. <i>Journal of Nutrition and Metabolism</i> , 2012, 2012, 1-9.	1.8	8
18	Toll-like receptor 3 expression and function in childhood idiopathic nephrotic syndrome. <i>Clinical and Experimental Immunology</i> , 2015, 182, 332-345.	2.6	8

#	ARTICLE	IF	CITATIONS
19	Neonatal high protein intake enhances neonatal growth without significant adverse renal effects in spontaneous IUGR piglets. <i>Physiological Reports</i> , 2017, 5, e13296.	1.7	8