
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5368693/publications.pdf Version: 2024-02-01



VASUARI SAUO

#	Article	IF	CITATIONS
1	Perfluorooctane Sulfonate (PFOS) and Related Perfluorinated Compounds in Human Maternal and Cord Blood Samples: Assessment of PFOS Exposure in a Susceptible Population during Pregnancy. Environmental Health Perspectives, 2004, 112, 1204-1207.	2.8	394
2	Correlations between Prenatal Exposure to Perfluorinated Chemicals and Reduced Fetal Growth. Environmental Health Perspectives, 2009, 117, 660-667.	2.8	250
3	Exposure to house dust phthalates in relation to asthma and allergies in both children and adults. Science of the Total Environment, 2014, 485-486, 153-163.	3.9	124
4	Effects of Prenatal Exposure to Polychlorinated Biphenyls and Dioxins on Mental and Motor Development in Japanese Children at 6 Months of Age. Environmental Health Perspectives, 2006, 114, 773-778.	2.8	118
5	Prenatal exposure to perfluorinated chemicals and relationship with allergies and infectious diseases in infants. Environmental Research, 2012, 112, 118-125.	3.7	116
6	Cohort Profile: The Hokkaido Study on Environment and Children's Health in Japan. International Journal of Epidemiology, 2011, 40, 611-618.	0.9	109
7	Relationships of Occupational Stress to Insomnia and Short Sleep in Japanese Workers. Sleep, 2005, 28, 728-735.	0.6	96
8	Relationship between selected indoor volatile organic compounds, so-called microbial VOC, and the prevalence of mucous membrane symptoms in single family homes. Science of the Total Environment, 2010, 408, 2208-2215.	3.9	93
9	Short Sleep Duration and Poor Sleep Quality Increase the Risk of Diabetes in Japanese Workers With No Family History of Diabetes. Diabetes Care, 2012, 35, 313-318.	4.3	93
10	Relationship between sick building syndrome and indoor environmental factors in newly built Japanese dwellings. International Archives of Occupational and Environmental Health, 2009, 82, 583-593.	1.1	85
11	Helicobacter Pylori Infection is a Significant Risk for Modified Lipid Profile in Japanese Male Subjects. Journal of Atherosclerosis and Thrombosis, 2010, 17, 1041-1048.	0.9	83
12	Relationship between indoor chemical concentrations and subjective symptoms associated with sick building syndrome in newly built houses in Japan. International Archives of Occupational and Environmental Health, 2010, 83, 225-235.	1.1	77
13	The relationship between exposure to microbial volatile organic compound and allergy prevalence in single-family homes. Science of the Total Environment, 2012, 423, 18-26.	3.9	69
14	A longitudinal study of aldehydes and volatile organic compounds associated with subjective symptoms related to sick building syndrome in new dwellings in Japan. Science of the Total Environment, 2012, 417-418, 61-67.	3.9	67
15	Twentyâ€fourâ€hour shift work, depressive symptoms, and job dissatisfaction among Japanese firefighters. American Journal of Industrial Medicine, 2008, 51, 380-391.	1.0	63
16	Naloxone lowers cerebrospinal fluid levels of excitatory amino acids after thoracoabdominal aortic surgery. Journal of Vascular Surgery, 2004, 40, 681-690.	0.6	60
17	Effects of prenatal exposure to dioxin-like compounds on allergies and infections during infancy. Environmental Research, 2011, 111, 551-558.	3.7	59
18	Biomonitoring of mercury, cadmium, and lead exposure in Japanese children: a cross-sectional study. Environmental Health and Preventive Medicine, 2015, 20, 18-27.	1.4	52

#	Article	IF	CITATIONS
19	Relationship of .BETA.2-Microglobulin to Arterial Stiffness in Japanese Subjects. Hypertension Research, 2005, 28, 505-511.	1.5	48
20	Caffeine intake, CYP1A2 polymorphism and the risk of recurrent pregnancy loss. Molecular Human Reproduction, 2005, 11, 357-360.	1.3	48
21	A Newly Estimated Glomerular Filtration Rate Is Independently Associated with Arterial Stiffness in Japanese Patients. Hypertension Research, 2008, 31, 193-201.	1.5	48
22	Job stress and depressive symptoms among Japanese fire fighters. American Journal of Industrial Medicine, 2007, 50, 470-480.	1.0	47
23	Hypothermic circulatory arrest is not a risk factor for neurologic morbidity in aortic surgery: A propensity score analysis. Journal of Thoracic and Cardiovascular Surgery, 2005, 130, 712-718.	0.4	45
24	Adverse Birth Outcomes Associated with Maternal Smoking and Polymorphisms in the N-Nitrosamine-Metabolizing Enzyme Genes NQO1 and CYP2E1. American Journal of Epidemiology, 2007, 167, 719-726.	1.6	44
25	Single nucleotide polymorphisms in the promoter region of the interleukin-6 gene and the risk of recurrent pregnancy loss in japanese women. Fertility and Sterility, 2004, 81, 374-378.	0.5	41
26	Concentrations of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and dioxin-like polychlorinated biphenyls in blood and breast milk collected from 60 mothers in Sapporo City, Japan. Chemosphere, 2008, 72, 1152-1158.	4.2	41
27	The relationship of gamma-glutamyltransferase to C-reactive protein and arterial stiffness. Nutrition, Metabolism and Cardiovascular Diseases, 2008, 18, 211-219.	1.1	37
28	Relation between Self-Reported Sleep Duration and Arterial Stiffness: A Cross-Sectional Study of Middle-Aged Japanese Civil Servants. Sleep, 2011, 34, 1681-1686.	0.6	37
29	Demographic, behavioral, dietary, and socioeconomic characteristics related to persistent organic pollutants and mercury levels in pregnant women in Japan. Chemosphere, 2015, 133, 13-21.	4.2	37
30	Association of blood mercury levels during pregnancy with infant birth size by blood selenium levels in the Japan Environment and Children's Study: A prospective birth cohort. Environment International, 2019, 125, 418-429.	4.8	36
31	Genetic Factors in Fetal Growth Restriction and Miscarriage. Seminars in Thrombosis and Hemostasis, 2005, 31, 334-345.	1.5	35
32	Gender differences in insomnia and the role of paid work and family responsibilities. Social Psychiatry and Psychiatric Epidemiology, 2012, 47, 651-662.	1.6	34
33	Effects of work burden, job strain and support on depressive symptoms and burnout among Japanese physicians. International Journal of Occupational Medicine and Environmental Health, 2014, 27, 980-992.	0.6	34
34	Relationship of Helicobacter pylori Infection to Arterial Stiffness in Japanese Subjects. Hypertension Research, 2005, 28, 283-292.	1.5	32
35	Relationship between two alternative occupational stress models and arterial stiffness: a cross-sectional study among Japanese workers. International Archives of Occupational and Environmental Health, 2009, 82, 175-183.	1.1	32
36	Blood persistent organochlorine pesticides in pregnant women in relation to physical and environmental variables in The Hokkaido Study on Environment and Children's Health. Science of the Total Environment, 2012, 426, 73-82.	3.9	32

#	Article	IF	CITATIONS
37	Epidemic of charcoal burning suicide in Japan. British Journal of Psychiatry, 2014, 204, 274-282.	1.7	32
38	Short Sleep Duration Increases the Risk of Chronic Kidney Disease in Shift Workers. Journal of Occupational and Environmental Medicine, 2014, 56, 1243-1248.	0.9	32
39	Relation of dampness to sick building syndrome in Japanese public apartment houses. Environmental Health and Preventive Medicine, 2009, 14, 26-35.	1.4	31
40	Relationships of C-reactive protein, uric acid, and glomerular filtration rate to arterial stiffness in Japanese subjects. Journal of Human Hypertension, 2005, 19, 907-913.	1.0	30
41	Job stress and burnout among urban and rural hospital physicians in Japan. Australian Journal of Rural Health, 2013, 21, 225-231.	0.7	30
42	Functional maternal catechol-O-methyltransferase polymorphism and fetal growth restriction. Pharmacogenetics and Genomics, 2006, 16, 775-781.	0.7	29
43	A Novel Role of the NRF2 Transcription Factor in the Regulation of Arsenite-Mediated Keratin 16 Gene Expression in Human Keratinocytes. Environmental Health Perspectives, 2008, 116, 873-879.	2.8	29
44	Social support and its interrelationships with demand–control model factors on presenteeism and absenteeism in Japanese civil servants. International Archives of Occupational and Environmental Health, 2017, 90, 539-553.	1.1	29
45	Metabolic Syndrome, C-Reactive Protein and Increased Arterial Stiffness in Japanese Subjects. Hypertension Research, 2006, 29, 589-596.	1.5	28
46	Effect of the Interaction Between Employment Level and Psychosocial Work Environment on Insomnia in Male Japanese Public Service Workers. International Journal of Behavioral Medicine, 2013, 20, 355-364.	0.8	28
47	Synergistic interaction between job control and social support at work on depression, burnout, and insomnia among Japanese civil servants. International Archives of Occupational and Environmental Health, 2015, 88, 143-152.	1.1	28
48	Sex-specific differences in effect of prenatal exposure to dioxin-like compounds on neurodevelopment in Japanese children: Sapporo cohort study. Environmental Research, 2017, 159, 222-231.	3.7	28
49	Serological Studies of Neurologic Helminthic Infections in Rural Areas of Southwest Cameroon: Toxocariasis, Cysticercosis and Paragonimiasis. PLoS Neglected Tropical Diseases, 2010, 4, e732.	1.3	28
50	Genetic association of aromatic hydrocarbon receptor (AHR) and cytochrome P450, family 1, subfamily A, polypeptide 1 (CYP1A1) polymorphisms with dioxin blood concentrations among pregnant Japanese women. Toxicology Letters, 2013, 219, 269-278.	0.4	27
51	Effects of the Interaction between Interleukin-6-634C/G Polymorphism and Smoking on Serum C-Reactive Protein Concentrations. Hypertension Research, 2007, 30, 593-599.	1.5	25
52	Relationships of job demand, job control, and social support on intention to leave and depressive symptoms in Japanese nurses. Industrial Health, 2016, 54, 32-41.	0.4	25
53	Impacts of Diabetes and an SGLT2 Inhibitor on the Clomerular Number and Volume in db/db Mice, as Estimated by Synchrotron Radiation Micro-CT at SPring-8. EBioMedicine, 2018, 36, 329-346.	2.7	25
54	Pregnancy-associated plasma protein-A polymorphism and the risk of recurrent pregnancy loss. Journal of Reproductive Immunology, 2006, 70, 99-108.	0.8	24

#	Article	IF	CITATIONS
55	Inflammation as a cardiovascular risk factor and pulse wave velocity as a marker of early-stage atherosclerosis in the Japanese population. Environmental Health and Preventive Medicine, 2009, 14, 159-164.	1.4	24
56	The Risk of Developing Diabetes in Association With Long Working Hours Differs by Shift Work Schedules. Journal of Epidemiology, 2016, 26, 481-487.	1.1	24
57	Association between duration of daily visual display terminal work and insomnia among local government clerks in Japan. American Journal of Industrial Medicine, 2008, 51, 148-156.	1.0	23
58	Effects of social relationships on mortality of the elderly: How do the influences change with the passage of time?. Archives of Gerontology and Geriatrics, 2008, 47, 327-339.	1.4	23
59	Dampness, food habits, and sick building syndrome symptoms in elementary school pupils. Environmental Health and Preventive Medicine, 2010, 15, 276-284.	1.4	23
60	Enhanced Transmural Dispersion of Repolarization in Patients with J Wave Syndromes. Journal of Cardiovascular Electrophysiology, 2012, 23, 1109-1114.	0.8	23
61	Late gadolinium enhancement on cardiac magnetic resonance represents the depolarizing and repolarizing electrically damaged foci causing malignant ventricular arrhythmia in hypertrophic cardiomyopathy. Heart Rhythm, 2015, 12, 1276-1284.	0.3	23
62	Effects of low-level prenatal exposure to dioxins on cognitive development in Japanese children at 42 months. Science of the Total Environment, 2018, 618, 1423-1430.	3.9	23
63	A polymorphism in the CYP17 gene and intrauterine fetal growth restriction. Molecular Human Reproduction, 2004, 10, 49-53.	1.3	22
64	Risk Factors and the Incidence of Coronary Artery Disease in Young Middleaged Japanese Men: Results from a 10Year Cohort Study. Internal Medicine, 2006, 45, 235-239.	0.3	22
65	Concentrations of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and dioxin-like polychlorinated biphenyls in blood collected from 195 pregnant women in Sapporo City, Japan. Chemosphere, 2007, 69, 1228-1237.	4.2	22
66	Relation between alcohol consumption and arterial stiffness: A cross-sectional study of middle-aged Japanese women and men. Alcohol, 2013, 47, 643-649.	0.8	22
67	Singleâ€Nucleotide Polymorphisms in the <i>SEPTIN12</i> Gene May Be a Genetic Risk Factor for Japanese Patients With Sertoli Cell–Only Syndrome. Journal of Andrology, 2012, 33, 483-487.	2.0	21
68	Clinicopathologic risk factors for recurrence of ovarian endometrioma following laparoscopic cystectomy. Acta Obstetricia Et Gynecologica Scandinavica, 2013, 92, 278-284.	1.3	21
69	Brachial-ankle pulse wave velocity is an independent predictor of incident hypertension in Japanese normotensive male subjects. Environmental Health and Preventive Medicine, 2011, 16, 217-223.	1.4	20
70	Prognostic factors in 105 Japanese cases of mycosis fungoides and Sézary syndrome: Clusterin expression as a novel prognostic factor. Journal of Dermatological Science, 2013, 71, 160-166.	1.0	20
71	Influence of genetic polymorphisms of styrene-metabolizing enzymes and smoking habits on levels of urinary metabolites after occupational exposure to styrene. Toxicology Letters, 2005, 160, 84-91.	0.4	19
72	Association between lifestyle habits and bone mineral density in Japanese juveniles. Environmental Health and Preventive Medicine, 2010, 15, 222-228.	1.4	19

#	Article	IF	CITATIONS
73	Post-Traumatic Stress Disorder and Job Stress among Firefighters of Urban Japan. Prehospital and Disaster Medicine, 2012, 27, 59-63.	0.7	19
74	Visit-to-visit variability and seasonal variation in blood pressure: Combination of Antihypertensive Therapy in the Elderly, Multicenter Investigation (CAMUI) Trial subanalysis. Clinical and Experimental Hypertension, 2015, 37, 411-419.	0.5	19
75	Relations of Occupational Stress to Occupational Class in Japanese Civil Servants-Analysis by Two Occupational Stress Models. Industrial Health, 2007, 45, 247-255.	0.4	18
76	The Effects of a Stress Inoculation Training Program for Civil Servants in Japan: a Pilot Study of a Non-randomized Controlled Trial. Industrial Health, 2009, 47, 173-182.	0.4	18
77	Time trends in method-specific suicide rates in Japan, 1990–2011. Epidemiology and Psychiatric Sciences, 2016, 25, 58-68.	1.8	18
78	Interleukin-4 Gene Polymorphism is not Involved in the Risk of Recurrent Pregnancy Loss. American Journal of Reproductive Immunology, 2004, 52, 143-146.	1.2	14
79	Congener-specific analysis of non-dioxin-like polychlorinated biphenyls in blood collected from 195 pregnant women in Sapporo City, Japan. Chemosphere, 2008, 73, 923-931.	4.2	14
80	Doseâ€dependent associations between prenatal caffeine consumption and small for gestational age, preterm birth, and reduced birthweight in the Japan Environment and Children's Study. Paediatric and Perinatal Epidemiology, 2019, 33, 185-194.	0.8	14
81	Cat and Dog Ownership in Early Life and Infant Development: A Prospective Birth Cohort Study of Japan Environment and Children's Study. International Journal of Environmental Research and Public Health, 2020, 17, 205.	1.2	14
82	Population Attributable Fractions of Modifiable Risk Factors for Nonsyndromic Orofacial Clefts: A Prospective Cohort Study From the Japan Environment and Children's Study. Journal of Epidemiology, 2021, 31, 272-279.	1.1	14
83	Potential impact of renin–angiotensin system inhibitors and calcium channel blockers on plasma high-molecular-weight adiponectin levels in hemodialysis patients. Hypertension Research, 2011, 34, 592-598.	1.5	13
84	Repolarization Characteristics in Early Repolarization and Brugada Syndromes: Insight into an Overlapping Mechanism of Lethal Arrhythmias. Journal of Cardiovascular Electrophysiology, 2014, 25, 1376-1384.	0.8	13
85	A prospective cohort study of insomnia and chronic kidney disease in Japanese workers. Sleep and Breathing, 2018, 22, 257-265.	0.9	13
86	Job Stress Factors Affect Workplace Resignation and Burnout among Japanese Rural Physicians. Tohoku Journal of Experimental Medicine, 2018, 245, 167-177.	0.5	13
87	Geography of suicide in Japan: spatial patterning and rural–urban differences. Social Psychiatry and Psychiatric Epidemiology, 2021, 56, 731-746.	1.6	13
88	бH pyloriï;¼²ï;¼² seropositivity and cytokine gene polymorphisms. World Journal of Gastroenterology, 2007, 13, 4445.	1.4	13
89	Relationship of socioeconomic status to C-reactive protein and arterial stiffness in urban Japanese civil servants. Social Science and Medicine, 2008, 67, 971-981.	1.8	12
90	Association of premorbid personality with behavioral and psychological symptoms in dementia with Lewy bodies: Comparison with Alzheimer's disease patients. Psychiatry and Clinical Neurosciences, 2017, 71, 409-416.	1.0	12

#	Article	IF	CITATIONS
91	Single-nucleotide polymorphisms in HORMAD1 may be a risk factor for azoospermia caused by meiotic arrest in Japanese patients. Asian Journal of Andrology, 2012, 14, 580-583.	0.8	12
92	Nationwide Study of Sick House Syndrome:. Japanese Journal of Hygiene, 2010, 65, 447-458.	0.6	11
93	Combination of antihypertensive therapy in the elderly, multicenter investigation (CAMUI) trial. Journal of Hypertension, 2013, 31, 1245-1255.	0.3	11
94	Spatial and temporal evolution of the epidemic of charcoal-burning suicide in Japan. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 857-868.	1.6	11
95	Single nucleotide polymorphism in the UBR2 gene may be a genetic risk factor for Japanese patients with azoospermia by meiotic arrest. Journal of Assisted Reproduction and Genetics, 2011, 28, 743-746.	1.2	10
96	Severity of low pre-pregnancy body mass index and perinatal outcomes: the Japan Environment and Children's Study. BMC Pregnancy and Childbirth, 2022, 22, 121.	0.9	10
97	Lipoprotein Metabolism, Insulin Resistance, and Adipocytokine Levels in Japanese Female Adolescents With a Normal Body Mass Index and High Body Fat Mass. Circulation Journal, 2009, 73, 534-539.	0.7	9
98	Single nucleotide polymorphisms in the SEPTIN12 gene may be associated with azoospermia by meiotic arrest in Japanese men. Journal of Assisted Reproduction and Genetics, 2012, 29, 47-51.	1.2	9
99	Relationships between Alcohol Intake and Ischemic Stroke Severity in Sex Stratified Analysis for Japanese Acute Stroke Patients. Journal of Stroke and Cerebrovascular Diseases, 2019, 28, 1604-1617.	0.7	9
100	Association between preâ€pregnancy body mass index and gestational weight gain and perinatal outcomes in pregnant women diagnosed with gestational diabetes mellitus: The Japan Environment and Children's Study. Journal of Diabetes Investigation, 2022, 13, 889-899.	1.1	9
101	Single-nucleotide polymorphisms in the human <i>RAD21L</i> gene may be a genetic risk factor for Japanese patients with azoospermia caused by meiotic arrest and Sertoli cell-only syndrome. Human Fertility, 2017, 20, 217-220.	0.7	8
102	Late gadolinium enhancement of cardiac magnetic resonance imaging indicates abnormalities of time-domain T-wave alternans in hypertrophic cardiomyopathy with ventricular tachycardia. Heart Rhythm, 2015, 12, 1747-1755.	0.3	7
103	Relationships between roadâ€distance to primary care facilities and ischemic heart disease and stroke mortality in Hokkaido, Japan: A Bayesian hierarchical approach to ecological count data. Journal of General and Family Medicine, 2018, 19, 4-8.	0.3	7
104	Prenatal alcohol exposure and adverse fetal growth restriction: findings from the Japan Environment and Children's Study. Pediatric Research, 2022, 92, 291-298.	1.1	7
105	Relations of mold, stove, and fragrance products on childhood wheezing and asthma: A prospective cohort study from the Japan Environment and Children's Study. Indoor Air, 2022, 32, .	2.0	7
106	The relationship between prenatal psychological stress and placental abruption in Japan, The Japan Environment and Children's Study (JECS). PLoS ONE, 2019, 14, e0219379.	1.1	6
107	Student Loans and Psychological Distress: A Cross-sectional Study of Young Adults in Japan. Journal of Epidemiology, 2020, 30, 436-441.	1.1	6
108	Work stress and oral conditions: a systematic review of observational studies. BMJ Open, 2021, 11, e046532.	0.8	6

#	Article	IF	CITATIONS
109	Indoor airborne mold spores in newly built dwellings. Environmental Health and Preventive Medicine, 2005, 10, 157-161.	1.4	5
110	Insight into specific pro-arrhythmic triggers in Brugada and early repolarization syndromes: results of long-term follow-up. Heart and Vessels, 2016, 31, 2035-2044.	0.5	5
111	Identifying a risk score for childhood obesity based on predictors identified in pregnant women and 1-year-old infants: An analysis of the data of the Hokkaido Study on Environment and Children's Health. Clinical Pediatric Endocrinology, 2019, 28, 81-89.	0.4	4
112	Accidental hypothermia: Factors related to a prolonged hospital stay – A nationwide observational study in Japan. American Journal of Emergency Medicine, 2021, 47, 169-175.	0.7	4
113	Association between frailty and mortality among patients with accidental hypothermia: a nationwide observational study in Japan. BMC Geriatrics, 2021, 21, 507.	1.1	4
114	Cross-sectional associations between oral diseases and work productivity loss among regular employees in Japan. Industrial Health, 2022, 61, 3-13.	0.4	4
115	Comparative study of simple semiquantitative dust mite allergen tests. Environmental Health and Preventive Medicine, 2007, 12, 187-192.	1.4	3
116	Combination of Antihypertensive Therapy in The Elderly, Multicenter Investigation (CAMUI) Trial. International Heart Journal, 2012, 53, 244-248.	0.5	3
117	An analysis of secular trends in method-specific suicides in Japan, 1950–1975. Population Health Metrics, 2017, 15, 14.	1.3	3
118	Assessment of suitable antihypertensive therapies: Combination with highâ€dose amlodipine/irbesartan vs triple combination with amlodipine/irbesartan/indapamide (ASAHlâ€Al study). Journal of Clinical Hypertension, 2020, 22, 1577-1584.	1.0	3
119	Lower Respiratory Tract Infections and Orofacial Clefts: A Prospective Cohort Study From the Japan Environment and Children's Study. Journal of Epidemiology, 2021, , .	1.1	3
120	An association study of the single-nucleotide polymorphism c190C>T (Arg64Cys) in the human testis-specific histone variant, H3t, of Japanese patients with Sertoli cell-only syndrome. Asian Journal of Andrology, 2018, 20, 527.	0.8	3
121	Higher Density of Primary Care Facilities Is Inversely Associated with Ischemic Heart Disease Mortality, but Not with Stroke Mortality: A Japanese Secondary Medical Service Area Level Ecological Count Data. Tohoku Journal of Experimental Medicine, 2020, 251, 217-224.	0.5	3
122	Relationship between Alcohol Intake and Stroke Severity in Japanese Patients: a Sex- and Subtype-Stratified Analysis. Journal of Stroke and Cerebrovascular Diseases, 2022, 31, 106513.	0.7	3
123	Prevalence of Asthma, Atopic Dermatitis, and Rhinitis and MVOC Exposure in Single Family Homes—A Survey in 6 Cities of Japan. Epidemiology, 2011, 22, S40-S41.	1.2	2
124	Spouse caregivers and behavioral and psychological symptoms of dementia. Aging and Mental Health, 2013, 17, 966-972.	1.5	2
125	The Association between Prenatal Yoga and the Administration of Ritodrine Hydrochloride during Pregnancy: An Adjunct Study of the Japan Environment and Children's Study. PLoS ONE, 2016, 11, e0158155.	1.1	2
126	Late Gadolinium Enhancement on Cardiac MRI Correlates with QT Dynamicity Represented by QT/RR Relationship in Patients with Ventricular Arrhythmias. Annals of Noninvasive Electrocardiology, 2016, 21, 126-135.	0.5	2

#	Article	IF	CITATIONS
127	Functional characteristics of left ventricular synchronization via right ventricular outflowâ€tract pacing detected by twoâ€dimensional strain echocardiography. Journal of Arrhythmia, 2017, 33, 28-34.	0.5	2
128	Effort–reward imbalance at work and tooth loss: a cross-sectional study from the J-SHINE project. Industrial Health, 2020, 58, 26-34.	0.4	2
129	Trajectories of the Psychological Status of Mothers of Infants With Nonsyndromic Orofacial Clefts: A Prospective Cohort Study From the Japan Environment and Children's Study. Cleft Palate-Craniofacial Journal, 2021, 58, 369-377.	0.5	2
130	Parental educational level and childhood wheezing and asthma: A prospective cohort study from the Japan Environment and Children's Study. PLoS ONE, 2021, 16, e0250255.	1.1	2
131	Association between Early Life Child Development and Family Dog Ownership: A Prospective Birth Cohort Study of the Japan Environment and Children's Study. International Journal of Environmental Research and Public Health, 2021, 18, 7082.	1.2	2
132	Factors correlating with serum birch pollen IgE status in pregnant women in Hokkaido, Japan: The Japan Environment and Children's Study (JECS). World Allergy Organization Journal, 2020, 13, 100128.	1.6	2
133	Silent cerebral ischemia detected by magnetic resonance imaging can predict postoperative delirium after total arch replacement for aneurysm. JTCVS Open, 2022, 10, 87-96.	0.2	2
134	Dampness, Food Habits, and Sick Building Syndrome Symptoms Among Elementary School Pupils. Epidemiology, 2011, 22, S39.	1.2	1
135	The relationship between repolarization parameters and serum electrolyte levels in patients with J wave syndromes. Magnesium Research, 2015, 28, 1-13.	0.4	1
136	Acute myocardial infarction and stoke after the enactment of smoke-free legislation in public places in Bibai city: data analysis of hospital admissions and ambulance transports. Hypertension Research, 2019, 42, 1801-1807.	1.5	1
137	Gender disparities in academic dermatology in Japan: Results from the first national survey. Journal of Dermatological Science, 2021, 102, 2-6.	1.0	1
138	Association between the Concentrations of Metallic Elements in Maternal Blood during Pregnancy and Prevalence of Abdominal Congenital Malformations: The Japan Environment and Children's Study. International Journal of Environmental Research and Public Health, 2021, 18, 10103.	1.2	1
139	Relation of Dampness to Sick Building Syndrome in Japanese Public Apartment Houses. Epidemiology, 2009, 20, S150.	1.2	1
140	Low HDL-cholesterol, hypertension and impaired glucose tolerance as predictors of acute myocardial infarction in northern area of Japan. Hokkaido Journal of Medical Science, 2006, 81, 25-30.	0.1	1
141	Recombinant Activated Factor VII in Aortic Surgery for Patients Under Hypothermic Circulatory Arrest. Therapeutics and Clinical Risk Management, 2022, Volume 18, 337-348.	0.9	1
142	Cut-off values of fasting plasma glucose for detecting impaired glucose tolerance in Japanese population – Relation between fasting and 2-h plasma glucose. Nutrition, Metabolism and Cardiovascular Diseases, 2007, 17, e5-e6.	1.1	0
143	Relationships of gamma-glutamyltransferase and beta 2-microglobulin on high sensitivity C-reactive protein among Japanese elementary school children. Journal of Pediatric Endocrinology and Metabolism, 2014, 27, 643-50.	0.4	0
144	PS 14-43 CHRONOTHERAPY WITH AN ARB AND CCB COMBINATION. Journal of Hypertension, 2016, 34, e446.	0.3	0

#	Article	IF	CITATIONS
145	Comparing Conscious Sedation With Regional Anesthesia Versus General Anesthesia in Minimally Invasive Mitral Valve Surgery With Right-Sided Minithoracotomy: A Retrospective Study. Journal of Cardiothoracic and Vascular Anesthesia, 2021, , .	0.6	0
146	Maternal psychological distress, education, household income, and congenital heart defects: a prospective cohort study from the Japan environment and children's study. BMC Pregnancy and Childbirth, 2021, 21, 544.	0.9	0
147	Diffuse Abdominal Uptake of Ga-67 Citrate in Two Cases. Clinical Nuclear Medicine, 2000, 25, 74.	0.7	0
148	Sick Building/House Syndrome. Current Topics in Environmental Health and Preventive Medicine, 2020, , 21-38.	0.1	0
149	Five-minute test to prevent postcardiotomy reexploration. JTCVS Techniques, 2022, 12, 121-129.	0.2	0
150	Post-cardiotomy pericardial effusion and postoperative atrial fibrillation risk. International Journal of Cardiovascular Imaging, 2022, 38, 1873-1879.	0.2	0
151	Association between maternal caffeine intake during pregnancy and child development at 6 and 12Âmonths: The Japan Environment and Children's Study. Early Human Development, 2022, 171, 105607.	0.8	0