Pier Luigi Temporelli

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/5372731/pier-luigi-temporelli-publications-by-year.pdf

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

127
papers5,248
citations33
h-index71
g-index161
ext. papers6,117
ext. citations4.6
avg, IF5.04
L-index

#	Paper	IF	Citations
127	Left atrial strain predicts exercise capacity in heart failure independently of left ventricular ejection fraction ESC Heart Failure, 2022,	3.7	3
126	Atrial fibrillation ablation in heart failure: Findings from the ESC-EHRA EORP Atrial Fibrillation Ablation long-term (AFA LT) registry. <i>International Journal of Cardiology</i> , 2022 , 346, 19-26	3.2	O
125	Clinical Impact and Prognostic Role of Triglyceride to High-Density Lipoprotein Cholesterol Ratio in Patients With Chronic Coronary Syndromes at Very High Risk: Insights From the START Study <i>Frontiers in Cardiovascular Medicine</i> , 2022 , 9, 874087	5.4	O
124	Is physical activity always good for you? The physical activity paradox. <i>European Heart Journal Supplements</i> , 2021 , 23, E168-E171	1.5	2
123	Heart valve calcification and cardiac hemodynamics. <i>Echocardiography</i> , 2021 , 38, 525-530	1.5	1
122	Impact of history of depression on 1-year outcomes in patients with chronic coronary syndromes: An analysis of a contemporary, prospective, nationwide registry. <i>International Journal of Cardiology</i> , 2021 , 331, 273-280	3.2	2
121	Prognostic relevance of Doppler echocardiographic re-assessment in HFrEF patients. <i>International Journal of Cardiology</i> , 2021 , 327, 111-116	3.2	2
120	Comparing the Prognostic Impact of Prediabetes with Diabetes in a Nationwide Cohort of Patients with Chronic Coronary Syndromes: An Analysis of the START Registry. <i>Cardiology</i> , 2021 , 146, 547-555	1.6	
119	Nonresponse to Acute Vasodilator Challenge and Prognosis in Heart Failure With Pulmonary Hypertension. <i>Journal of Cardiac Failure</i> , 2021 , 27, 869-876	3.3	O
118	Pre-existing type 2 diabetes is associated with increased all-cause death independently of echocardiographic predictors of poor prognosis only in ischemic heart disease. <i>Nutrition, Metabolism and Cardiovascular Diseases,</i> 2020 , 30, 2036-2040	4.5	1
117	Current lipid lowering treatment and attainment of LDL targets recommended by ESC/EAS guidelines in very high-risk patients with established atherosclerotic cardiovascular disease: Insights from the START registry. <i>International Journal of Cardiology</i> , 2020 , 316, 229-235	3.2	7
116	Relation between plasma ceramides and cardiovascular death in chronic heart failure: A subset analysis of the GISSI-HF trial. <i>ESC Heart Failure</i> , 2020 , 7, 3288	3.7	6
115	Lipid Lowering Treatment and Eligibility for PCSK9 Inhibition in Post-Myocardial Infarction Patients in Italy: Insights from Two Contemporary Nationwide Registries. <i>Cardiovascular Therapeutics</i> , 2020 , 2020, 3856242	3.3	5
114	External applicability of the ISCHEMIA trial: an analysis of a prospective, nationwide registry of patients with stable coronary artery disease. <i>EuroIntervention</i> , 2020 , 16, e966-e973	3.1	12
113	Fifteen-Year Trends of Cardiogenic Shock and Mortality in Patients with Diabetes and Acute Coronary Syndromes. <i>American Journal of Medicine</i> , 2020 , 133, 331-339.e2	2.4	4
112	Which patients with atrial fibrillation undergo an ablation procedure today in Europe? A report from the ESC-EHRA-EORP Atrial Fibrillation Ablation Long-Term and Atrial Fibrillation General Pilot Registries. <i>Europace</i> , 2020 , 22, 250-258	3.9	2
111	Association of Hypertriglyceridemia with All-Cause Mortality and Atherosclerotic Cardiovascular Events in a Low-Risk Italian Population: The TG-REAL Retrospective Cohort Analysis. <i>Journal of the American Heart Association</i> , 2020 , 9, e015801	6	17

110	Prevention after acute coronary syndrome the Qess is moreQohilosophy. <i>European Heart Journal Supplements</i> , 2020 , 22, E153-E156	1.5	
109	Additive Value of Biomarkers and Echocardiography to Stratify the Risk of Death in Heart Failure Patients with Reduced Ejection Fraction. <i>Cardiology Research and Practice</i> , 2019 , 2019, 1824816	1.9	3
108	Awareness and appropriateness of the management of preclinical heart failure in outpatient clinics in Italy: Insights from the VASTISSIMO study - EValuation of the AppropriateneSs of The preclinical phase (Stage A and Stage B) of Heart Failure Management in Outpatient Clinics in Italy. <i>Monaldi</i>	2.7	2
107	Clinical characteristics of heart failure patients undergoing atrial fibrillation ablation today in Europe. Data from the atrial fibrillation registries of the European Society of Cardiology and the European Heart Rhythm Association. <i>European Journal of Heart Failure</i> , 2019 , 21, 690-693	12.3	2
106	Four golden rules to halve the risk of cardiovascular events. <i>European Heart Journal Supplements</i> , 2019 , 21, B103-B104	1.5	
105	Similar predictive value of six-minute walking distance and B-type natriuretic peptide in heart failure with reduced to mid-range ejection fraction. <i>Monaldi Archives for Chest Disease</i> , 2019 , 89,	2.7	1
104	Referral from vascular surgery to cardiovascular rehabilitation and related outcomes in patients with peripheral arterial disease: the THINKPAD-RELOADED survey. <i>Monaldi Archives for Chest Disease</i> , 2019 , 89,	2.7	1
103	Brachial pulse pressure in acute heart failure. Results of the Heart Failure Registry. <i>ESC Heart Failure</i> , 2019 , 6, 1167-1177	3.7	2
102	Clinical outcomes, pharmacological treatment, and quality of life of patients with stable coronary artery diseases managed by cardiologists: 1-year results of the START study. <i>European Heart Journal Quality of Care & Dutcomes</i> , 2019, 5, 334-342	4.6	8
101	Left atrial dilatation in systolic heart failure: a marker of poor prognosis, not just a buffer between the left ventricle and pulmonary circulation. <i>Journal of Echocardiography</i> , 2018 , 16, 155-161	1.6	9
100	Prognostic impact of elevated serum uric acid levels on long-term outcomes in patients with chronic heart failure: A post-hoc analysis of the GISSI-HF (Gruppo Italiano per lo Studio della Sopravvivenza nella Insufficienza Cardiaca-Heart Failure) trial. <i>Metabolism: Clinical and Experimental</i>	12.7	20
99	, 2018, 83, 205-215 Hypertriglyceridemia and omega-3 fatty acids: Their often overlooked role in cardiovascular disease prevention. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018, 28, 197-205	4.5	24
98	Prevalence and management of familial hypercholesterolemia in patients with coronary artery disease: The heredity survey. <i>International Journal of Cardiology</i> , 2018 , 252, 193-198	3.2	25
97	Characteristics, treatments and 1-year prognosis of hospitalized and ambulatory heart failure patients with chronic obstructive pulmonary disease in the European Society of Cardiology Heart Failure Long-Term Registry. <i>European Journal of Heart Failure</i> , 2018 , 20, 100-110	12.3	49
96	Characteristics, treatment and quality of life of stable coronary artery disease patients with or without angina: Insights from the START study. <i>PLoS ONE</i> , 2018 , 13, e0199770	3.7	6
95	Questioning the Associations of EB Fatty Acid Supplement Use With Cardiovascular Disease Risks. JAMA Cardiology, 2018 , 3, 781	16.2	1
94	Haemodynamic effects of an acute vasodilator challenge in heart failure patients with reduced ejection fraction and different forms of post-capillary pulmonary hypertension. <i>European Journal of Heart Failure</i> , 2018 , 20, 725-734	12.3	17
93	Current management and treatment of patients with stable coronary artery diseases presenting to cardiologists in different clinical contexts: A prospective, observational, nationwide study. European Journal of Preventive Cardiology, 2018, 25, 43-53	3.9	19

92	Cardiac Prevention and Rehabilitation "3.0": From acute to chronic phase. Position Paper of the Italian Association for Cardiovascular Prevention and Rehabilitation (GICR-IACPR). <i>Monaldi Archives for Chest Disease</i> , 2018 , 88, 1004	2.7	7
91	Exercise gas exchange in continuous-flow left ventricular assist device recipients. <i>PLoS ONE</i> , 2018 , 13, e0187112	3.7	10
90	Prevalence and pharmacologic management of familial hypercholesterolemia in an unselected contemporary cohort of patients with stable coronary artery disease. <i>Clinical Cardiology</i> , 2018 , 41, 1075	;- 1 ∙∂83	5
89	Association between decreasing estimated glomerular filtration rate and risk of cardiac conduction defects in patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2018 , 44, 473-481	5.4	1
88	Refined 4-group classification of left ventricular hypertrophy based on ventricular concentricity and volume dilatation outlines distinct noninvasive hemodynamic profiles in a large contemporary echocardiographic population. <i>Echocardiography</i> , 2018 , 35, 1258-1265	1.5	3
87	Frequent coexistence of chronic heart failure and chronic obstructive pulmonary disease in respiratory and cardiac outpatients: Evidence from SUSPIRIUM, a multicentre Italian survey. <i>European Journal of Preventive Cardiology</i> , 2017 , 24, 567-576	3.9	23
86	Association Between Diabetes and 1-Year Adverse Clinical Outcomes in a Multinational Cohort of Ambulatory Patients With Chronic Heart Failure: Results From the ESC-HFA Heart Failure Long-Term Registry. <i>Diabetes Care</i> , 2017 , 40, 671-678	14.6	65
85	Prognostic effects of rosuvastatin in patients with co-existing chronic obstructive pulmonary disease and chronic heart failure: A sub-analysis of GISSI-HF trial. <i>Pulmonary Pharmacology and Therapeutics</i> , 2017 , 44, 16-23	3.5	12
84	Persistent abnormalities in pulmonary arterial compliance after heart transplantation in patients with combined post-capillary and pre-capillary pulmonary hypertension. <i>PLoS ONE</i> , 2017 , 12, e0188383	3.7	8
83	Jogging and mortality: is there a U-shaped curve?. <i>Journal of Cardiovascular Medicine</i> , 2017 , 18 Suppl 1, e177-e179	1.9	1
82	Prognostic Impact of Diabetes and Prediabetes on Survival Outcomes in Patients With Chronic Heart Failure: A Post-Hoc Analysis of the GISSI-HF (Gruppo Italiano per lo Studio della Sopravvivenza nella Insufficienza Cardiaca-Heart Failure) Trial. <i>Journal of the American Heart</i>	6	39
81	Association, 2017 , 6, Pulmonary hemodynamics in heart failure patients with reduced or preserved ejection fraction and pulmonary hypertension: Similarities and disparities. <i>American Heart Journal</i> , 2017 , 192, 120-127	4.9	23
80	In-hospital and 1-year mortality associated with diabetes in patients with acute heart failure: results from the ESC-HFA Heart Failure Long-Term Registry. <i>European Journal of Heart Failure</i> , 2017 , 19, 54-65	12.3	92
79	Different correlates but similar prognostic implications for right ventricular dysfunction in heart failure patients with reduced or preserved ejection fraction. <i>European Journal of Heart Failure</i> , 2017 , 19, 873-879	12.3	122
78	Prevalence and Prognostic Impact of Chronic Obstructive Pulmonary Disease in Patients with Chronic Heart Failure: Data from the GISSI-HF Trial. <i>Cardiology</i> , 2017 , 136, 128-137	1.6	27
77	ANMCO/GICR-IACPR/SICI-GISE Consensus Document: the clinical management of chronic ischaemic cardiomyopathy. <i>European Heart Journal Supplements</i> , 2017 , 19, D163-D189	1.5	
76	Functional mitral regurgitation: are we treating the real target?. <i>Journal of Cardiovascular Medicine</i> , 2016 , 17, 767-73	1.9	3
75	Effects of Polyunsaturated Fatty Acid Treatment on Postdischarge Outcomes After Acute Myocardial Infarction. <i>American Journal of Cardiology</i> , 2016 , 117, 340-6	3	12

(2013-2016)

74	Prognostic impact of in-hospital hyperglycemia in hospitalized patients with acute heart failure: Results of the IN-HF (Italian Network on Heart Failure) Outcome registry. <i>International Journal of Cardiology</i> , 2016 , 203, 587-93	3.2	27
73	Cardiopulmonary rehabilitation in patiens with heart failure and chronic pulmonary disease. <i>Monaldi Archives for Chest Disease</i> , 2016 , 84, 739	2.7	
72	Echo and BNP serial assessment in ambulatory heart failure care: Data on loop diuretic use and renal function. <i>Data in Brief</i> , 2016 , 9, 1074-1076	1.2	1
71	Echo and natriuretic peptide guided therapy improves outcome and reduces worsening renal function in systolic heart failure: An observational study of 1137 outpatients. <i>International Journal of Cardiology</i> , 2016 , 224, 416-423	3.2	20
70	Right ventricular recovery during follow-up is associated with improved survival in patients with chronic heart failure with reduced ejection fraction. <i>European Journal of Heart Failure</i> , 2016 , 18, 1462-14	4 7 7.3	24
69	Beta-blockers can improve survival in medically-treated patients with severe symptomatic aortic stenosis. <i>International Journal of Cardiology</i> , 2015 , 190, 15-7	3.2	13
68	Echocardiography of Right Ventriculoarterial Coupling Combined With Cardiopulmonary Exercise Testing to Predict Outcome in Heart Failure. <i>Chest</i> , 2015 , 148, 226-234	5.3	85
67	A retrospective multicenter study on long-term prevalence of chronic pain after cardiac surgery. Journal of Cardiovascular Medicine, 2015 , 16, 768-74	1.9	22
66	A retrospective multicenter study on long-term prevalence of chronic pain after cardiac surgery. Journal of Cardiovascular Medicine, 2015 , 16, 857	1.9	78
65	The prognostic impact of diastolic dysfunction in patients with chronic heart failure and post-acute myocardial infarction: Can age-stratified E/A ratio alone predict survival?. <i>International Journal of Cardiology</i> , 2015 , 181, 362-8	3.2	11
64	Worksite health and wellness in the European union. <i>Progress in Cardiovascular Diseases</i> , 2014 , 56, 508-	18 .5	25
63	Mitral and aortic valve sclerosis/calcification and carotid atherosclerosis: results from 1065 patients. <i>Heart and Vessels</i> , 2014 , 29, 776-83	2.1	20
62	Obesity paradox in patients with aortic valve stenosis. Protective effect of body mass index independently of age, disease severity, treatment modality and non-cardiac comorbidities. <i>International Journal of Cardiology</i> , 2014 , 176, 1441-3	3.2	9
61	Prognostic relevance of pulmonary arterial compliance in patients with chronic heart failure. <i>Chest</i> , 2014 , 145, 1064-1070	5.3	101
60	Prevalence and 1-year prognosis of transient heart failure following coronary revascularization. <i>Internal and Emergency Medicine</i> , 2014 , 9, 641-7	3.7	2
59	Italian Survey on Prevalence and Disease Management of Chronic Heart Failure and Chronic Obstructive Pulmonary Disease comorbidity in ambulatory patients. SUSPIRIUM study rationale and design. <i>Monaldi Archives for Chest Disease</i> , 2014 , 82, 29-34	2.7	2
58	Lower extremities peripheral arterial disease among patients admitted to cardiac rehabilitation: the THINKPAD registry. <i>International Journal of Cardiology</i> , 2014 , 171, 192-8	3.2	13
57	Effective secondary prevention through cardiac rehabilitation after coronary revascularization and predictors of poor adherence to lifestyle modification and medication. Results of the ICAROS Survey. <i>International Journal of Cardiology</i> , 2013 , 167, 1390-5	3.2	57

56	Tricuspid annular plane systolic excursion and pulmonary arterial systolic pressure relationship in heart failure: an index of right ventricular contractile function and prognosis. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2013 , 305, H1373-81	5.2	282
55	Flow-mediated dilation normalization predicts outcome in chronic heart failure patients. <i>Journal of Cardiac Failure</i> , 2013 , 19, 260-7	3.3	27
54	Effects on survival of loop diuretic dosing in ambulatory patients with chronic heart failure using a propensity score analysis. <i>International Journal of Clinical Practice</i> , 2013 , 67, 656-64	2.9	33
53	Severe aortic valve stenosis with normal left ventricular function and low vs. high pressure gradient: different hemodynamic profiles but similar clinical presentation, comorbidity and outcome. <i>International Journal of Cardiology</i> , 2013 , 167, 2326-8	3.2	6
52	How to face emergencies in heart failure patients with ventricular assist device. <i>International Journal of Cardiology</i> , 2013 , 168, 5143-8	3.2	7
51	Cardiovascular risk profile and lifestyle habits in a cohort of Italian cardiologists (from the SOCRATES Survey). <i>American Journal of Cardiology</i> , 2013 , 112, 226-30	3	11
50	Relationship of pulmonary hypertension and right ventricular dysfunction with survival of elderly patients with chronic systolic heart failure. <i>European Heart Journal</i> , 2013 , 34, 3630-3630	9.5	
49	Prognostic relevance of a non-invasive evaluation of right ventricular function and pulmonary artery pressure in patients with chronic heart failure. <i>European Journal of Heart Failure</i> , 2013 , 15, 408-1	4 ^{12.3}	89
48	Prognostic implications of left ventricular dilation in patients with nonischemic heart failure: interactions with restrictive filling pattern and mitral regurgitation. <i>Congestive Heart Failure</i> , 2012 , 18, 198-204		1
47	Gender differences in cardiac rehabilitation programs from the Italian survey on cardiac rehabilitation (ISYDE-2008). <i>International Journal of Cardiology</i> , 2012 , 160, 133-9	3.2	15
46	Aortic valve sclerosis is a marker of atherosclerosis independently of traditional clinical risk factors. Analysis in 712 patients without ischemic heart disease. <i>International Journal of Cardiology</i> , 2012 , 158, 163-4	3.2	3
45	The inability to perform a 6 minute walking test after cardio-thoracic surgery is a marker of clinical severity and poor outcome. Data from the ISYDE-2008 Italian survey. <i>International Journal of Cardiology</i> , 2011 , 151, 115-6	3.2	7
44	Late postoperative atrial fibrillation after cardiac surgery: a national survey within the cardiac rehabilitation setting. <i>Journal of Cardiovascular Medicine</i> , 2011 , 12, 390-5	1.9	16
43	Independent prognostic value of functional mitral regurgitation in patients with heart failure. A quantitative analysis of 1256 patients with ischaemic and non-ischaemic dilated cardiomyopathy. Heart, 2011 , 97, 1675-80	5.1	360
42	Doppler echocardiography in advanced systolic heart failure: a noninvasive alternative to Swan-Ganz catheter. <i>Circulation: Heart Failure</i> , 2010 , 3, 387-94	7.6	67
41	Validation of an echo-Doppler decision model to predict left ventricular filling pressure in patients with heart failure independently of ejection fraction. <i>European Journal of Echocardiography</i> , 2010 , 11, 703-10		37
40	Long-term secondary prevention programs after cardiac rehabilitation for the reduction of future cardiovascular events: focus on regular physical activity. <i>Future Cardiology</i> , 2009 , 5, 297-314	1.3	7
39	Independent relationship of left atrial size and mortality in patients with heart failure: an individual patient meta-analysis of longitudinal data (MeRGE Heart Failure). <i>European Journal of Heart Failure</i> , 2009 , 11, 929-36	12.3	109

(2001-2009)

38	Early right coronary vasospasm presenting with malignant arrhythmias in a heart transplantation recipient without allograft vasculopathy. <i>International Journal of Cardiology</i> , 2009 , 131, e120-3	3.2	6
37	Global secondary prevention strategies to limit event recurrence after myocardial infarction: results of the GOSPEL study, a multicenter, randomized controlled trial from the Italian Cardiac Rehabilitation Network. <i>Archives of Internal Medicine</i> , 2008 , 168, 2194-204		258
36	Perioperative and postoperative predictors of outcome in patients with low ejection fraction early after coronary artery bypass grafting: the additional value of left ventricular remodeling. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2008 , 15, 441-7		4
35	Italian survey on cardiac rehabilitation and secondary prevention after cardiac revascularization: ICAROS study. A survey from the Italian cardiac rehabilitation network: rationale and design. <i>Monaldi Archives for Chest Disease</i> , 2008 , 70, 99-104, 106	2.7	4
34	Low plasma levels of brain natriuretic peptide in severe acute heart failure: merely a case?. <i>International Journal of Cardiology</i> , 2007 , 122, e18-20	3.2	1
33	Statement on cardiopulmonary exercise testing in chronic heart failure due to left ventricular dysfunction: recommendations for performance and interpretation Part II: How to perform cardiopulmonary exercise testing in chronic heart failure. European Journal of Cardiovascular		3
32	Right ventricular functional recovery after acute myocardial infarction: relation with left ventricular function and interventricular septum motion. GISSI-3 echo substudy. <i>Heart</i> , 2005 , 91, 484-8	5.1	42
31	The impact of growth hormone/insulin-like growth factor-I axis and nocturnal breathing disorders on cardiovascular features of adult patients with Prader-Willi syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 5639-46	5.6	36
30	GlObal Secondary Prevention strategiEs to Limit event recurrence after myocardial infarction: the GOSPEL study. A trial from the Italian Cardiac Rehabilitation Network: rationale and design. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2005 , 12, 555-561		11
29	Left atrium remodeling after acute myocardial infarction (results of the GISSI-3 Echo Substudy). <i>American Journal of Cardiology</i> , 2004 , 93, 1156-9	3	23
28	Doppler-derived mitral deceleration time as a strong prognostic marker of left ventricular remodeling and survival after acute myocardial infarction: results of the GISSI-3 echo substudy. <i>Journal of the American College of Cardiology</i> , 2004 , 43, 1646-53	15.1	117
27	Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). <i>BMJ</i> , <i>The</i> , 2004 , 328, 189	5.9	388
26	Limited predictive value of cardiopulmonary exercise indices in patients with moderate chronic heart failure treated with carvedilol. <i>American Heart Journal</i> , 2004 , 147, 553-60	4.9	24
25	Antiremodeling effect of long-term exercise training in patients with stable chronic heart failure: results of the Exercise in Left Ventricular Dysfunction and Chronic Heart Failure (ELVD-CHF) Trial. <i>Circulation</i> , 2003 , 108, 554-9	16.7	251
24	Relation between early mitral regurgitation and left ventricular thrombus formation after acute myocardial infarction: results of the GISSI-3 echo substudy. <i>British Heart Journal</i> , 2002 , 88, 131-6		14
23	Prediction of 6 months left ventricular dilatation after myocardial infarction in relation to cardiac morbidity and mortality. Application of a new dilatation model to GISSI-3 data. <i>European Heart Journal</i> , 2002 , 23, 536-42	9.5	20
22	Accurate noninvasive estimation of pulmonary vascular resistance by Doppler echocardiography in patients with chronic failure heart failure. <i>Journal of the American College of Cardiology</i> , 2001 , 37, 1813	-9 ^{15.1}	84
21	Heterogeneity of left ventricular remodeling after acute myocardial infarction: results of the Gruppo Italiano per lo Studio della Sopravvivenza nell@nfarto Miocardico-3 Echo Substudy. <i>American Heart Journal</i> , 2001 , 141, 131-8	4.9	83

20	Chronic mitral regurgitation and Doppler estimation of left ventricular filling pressures in patients with heart failure. <i>Journal of the American Society of Echocardiography</i> , 2001 , 14, 1094-9	5.8	15
19	Electrocardiographic evolutionary changes and left ventricular remodeling after acute myocardial infarction: results of the GISSI-3 Echo substudy. <i>Journal of the American College of Cardiology</i> , 2000 , 35, 127-35	15.1	39
18	Estimation of pulmonary wedge pressure by transmitral Doppler in patients with chronic heart failure and atrial fibrillation. <i>American Journal of Cardiology</i> , 1999 , 83, 724-7	3	82
17	Doppler-derived acceleration rate of right ventricular early filling as a measurement of right atrial pressure in chronic heart failure secondary to ischemic or idiopathic dilated cardiomyopathy. <i>American Journal of Cardiology</i> , 1998 , 81, 513-5	3	10
16	Heterogeneous fate of perfusion and contraction after anterior wall acute myocardial infarction and effects on left ventricular remodeling. <i>American Journal of Cardiology</i> , 1998 , 82, 1457-62	3	О
15	Reversible restrictive left ventricular diastolic filling with optimized oral therapy predicts a more favorable prognosis in patients with chronic heart failure. <i>Journal of the American College of Cardiology</i> , 1998 , 31, 1591-7	15.1	112
14	Amlodipine Improves the Anti-Ischaemic Effect of Atenolol in Postinfarction Patients with Effort-Induced Ischaemia. <i>Clinical Drug Investigation</i> , 1997 , 13, 22-28	3.2	
13	Attenuation of unfavorable remodeling by exercise training in postinfarction patients with left ventricular dysfunction: results of the Exercise in Left Ventricular Dysfunction (ELVD) trial. <i>Circulation</i> , 1997 , 96, 1790-7	16.7	123
12	Independent and incremental prognostic value of Doppler-derived mitral deceleration time of early filling in both symptomatic and asymptomatic patients with left ventricular dysfunction. <i>Journal of the American College of Cardiology</i> , 1996 , 28, 383-90	15.1	207
11	Independent and Incremental Prognostic Value of Doppler-Derived Mitral Deceleration Time of Early Filling in Both Symptomatic and Asymptomatic Patients With Left Ventricular Dysfunction. <i>Journal of the American College of Cardiology</i> , 1996 , 28, 383-390	15.1	198
10	Soho: A modular spacecraft concept to allow flexible payload integration and efficient development. <i>Acta Astronautica</i> , 1995 , 37, 277-291	2.9	2
9	Technetium-99m sestamibi tomographic evaluation of residual ischemia after anterior myocardial infarction. <i>Journal of the American College of Cardiology</i> , 1995 , 25, 590-6	15.1	13
8	Residual exertional ischemia and unfavorable left ventricular remodeling in patients with systolic dysfunction after anterior myocardial infarction. <i>Journal of the American College of Cardiology</i> , 1995 , 25, 1539-46	15.1	16
7	Images in clinical medicine. Negative U waves as an indicator of stress-induced myocardial ischemia. <i>New England Journal of Medicine</i> , 1994 , 330, 1791	59.2	1
6	Doppler-derived mitral deceleration time of early filling as a strong predictor of pulmonary capillary wedge pressure in postinfarction patients with left ventricular systolic dysfunction. <i>Journal of the American College of Cardiology</i> , 1994 , 23, 1630-7	15.1	312
5	Spontaneous delayed recovery of perfusion and contraction after the first 5 weeks after anterior infarction. Evidence for the presence of hibernating myocardium in the infarcted area. <i>Circulation</i> , 1994 , 90, 1386-97	16.7	112
4	Long-term physical training and left ventricular remodeling after anterior myocardial infarction: results of the Exercise in Anterior Myocardial Infarction (EAMI) trial. EAMI Study Group. <i>Journal of the American College of Cardiology</i> , 1993 , 22, 1821-9	15.1	124
3	Increased fibrin turnover and high PAI-1 activity as predictors of ischemic events in atherosclerotic patients. A case-control study. The PLAT Group. <i>Arteriosclerosis and Thrombosis: A Journal of Vascular Biology.</i> 1993 13, 1412-7		177

LIST OF PUBLICATIONS

2	EAMI E xercise Training in Anterior Myocardial Infarction: An Ongoing Multicenter Randomized Study*. <i>Chest</i> , 1992 , 101, 315S-321S	5.3	6	
1	Inaccuracy of various proposed electrocardiographic criteria in the diagnosis of apical myocardial infarctiona critical review. <i>European Heart Journal</i> , 1989 , 10, 880-6	9.5	7	