## Elena Anufriieva

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5381129/publications.pdf

Version: 2024-02-01

516215 642321 65 827 16 23 citations g-index h-index papers 69 69 69 358 docs citations times ranked citing authors all docs

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Human-Induced Sharp Salinity Changes in the World's Largest Hypersaline Lagoon Bay Sivash (Crimea) and Their Effects on the Ecosystem. Water (Switzerland), 2022, 14, 403.   | 1.2 | 11        |
| 2  | Spatio-temporal variability of zooplankton and zoobenthos as the elements of integrated zoocenosis in a marine lake (Crimea, Black Sea): What is a general pattern?. Journal of Sea Research, 2022, 185, 102231.   | 0.6 | 4         |
| 3  | Feeding behavior of <i>Gammarus aequicauda</i> in the presence of two prey species of <i>Artemia</i> sp. and <i>Baeotendipes noctivagus</i> Ecological and Integrative Physiology, 2022, 337, 768-775.   | 0.9 | 2         |
| 4  | <i>Cladophora</i> spp. (Chlorophyta) modulate environment and create a habitat for microalgae in hypersaline waters. European Journal of Phycology, 2021, 56, 231-243.   | 0.9 | 16        |
| 5  | Can Gammarus aequicauda (Amphipoda) suppress a population of Baeotendipes noctivagus<br>(Chironomidae) in a hypersaline lake? A case of Lake Moynaki (Crimea). Aquaculture Research, 2021, 52,<br>1705-1714.   | 0.9 | 8         |
| 6  | Mercury in the world's largest hypersaline lagoon Bay Sivash, the Sea of Azov. Environmental Science and Pollution Research, 2021, 28, 28704-28712.  | 2.7 | 5         |
| 7  | Microphytobenthos in the Hypersaline Water Bodies, the Case of Bay Sivash (Crimea): Is Salinity the Main Determinant of Species Composition?. Water (Switzerland), 2021, 13, 1542.   | 1.2 | 8         |
| 8  | The behavior of <i>Gammarus aequicauda</i> (crustacea, amphipoda) during predation on chironomid larvae: Sex differences and changes in precopulatory mateâ€guarding state. Journal of Experimental Zoology Part A: Ecological and Integrative Physiology, 2021, 335, 572-582. | 0.9 | 3         |
| 9  | Salt to conserve: a review on the ecology and preservation of hypersaline ecosystems. Biological Reviews, 2021, 96, 2828-2850.   | 4.7 | 47        |
| 10 | Spatial and temporal variability of microphytobenthos in a marine hypersaline lake (Crimea): Are there some general patterns?. Journal of Sea Research, 2021, 177, 102121.   | 0.6 | 3         |
| 11 | Appearance of a New Species of Cladocera (Anomopoda, Chydoridae, Bosminidae) in the Hypersaline<br>Moynaki Lake, Crimea. Biology Bulletin, 2021, 48, 934-937.  | 0.1 | 4         |
| 12 | The longâ€term changes in plankton composition: Is Bay Sivash transforming back into one of the world's largest habitats of <i>Artemia </i> sp. (Crustacea, Anostraca)?. Aquaculture Research, 2020, 51, 341-350.  | 0.9 | 21        |
| 13 | Ecosystems of artificial saline lakes. A case of Lake Magic in Wadi El-Rayan depression (Egypt).<br>Knowledge and Management of Aquatic Ecosystems, 2020, , 31.  | 0.5 | 9         |
| 14 | Trace Elements in the Bottom Sediments of the Crimean Saline Lakes. Is It Possible to Explain Their Concentration Variability?. Water (Switzerland), 2020, 12, 2364.   | 1.2 | 7         |
| 15 | Behavior of <i>Gammarus aequicauda</i> (Crustacea, Amphipoda) during predation on <i>Artemia</i> (Crustacea, Anostraca): New experimental results. International Review of Hydrobiology, 2020, 105, 143-150.   | 0.5 | 6         |
| 16 | Is biomass of filamentous green algae <i>Cladophora</i> spp. (Chlorophyta, Ulvophyceae) an unlimited cheap and valuable resource for medicine and pharmacology? A review. Reviews in Aquaculture, 2020, 12, 2493-2510.   | 4.6 | 10        |
| 17 | Structure and Trophic Relations in Hypersaline Environments. Biology Bulletin Reviews, 2020, 10, 48-56.  | 0.3 | 12        |
| 18 | Natural radionuclides in bottom sediments of the saline lakes. What factors determine their concentration?. Environmental Earth Sciences, 2020, 79, 1.   | 1.3 | 9         |

| #  | Article  | IF        | CITATIONS     |
|----|--|-----------|---------------|
| 19 | Does Salinity Affect the Distribution of the Artificial Radionuclides 90Sr and 137Cs in Water of the Saline Lakes? A Case of the Crimean Peninsula. Water (Switzerland), 2020, 12, 349.                          | 1.2       | 14            |
| 20 | <i>Gammarus aequicauda</i> and <i>Moina salina</i> in the Crimean saline waters: New experimental and field data on their trophic relation. Aquaculture Research, 2020, 51, 3091-3099.                           | 0.9       | 12            |
| 21 | Macrostructure of benthos along a salinity gradient: The case of Sivash Bay (the Sea of Azov), the largest hypersaline lagoon worldwide. Journal of Sea Research, 2019, 154, 101811.                             | 0.6       | 15            |
| 22 | Do separated taxa react differently to a long-term salinity increase? The meiobenthos changes in Bay Sivash, largest hypersaline lagoon worldwide. Knowledge and Management of Aquatic Ecosystems, 2019, , 36.   | 0.5       | 12            |
| 23 | Suppression of <i>Artemia</i> spp. (Crustacea, Anostraca) populations by predators in the Crimean hypersaline lakes: A review of the evidence. International Review of Hydrobiology, 2019, 104, 5-13.            | 0.5       | 25            |
| 24 | Distribution and Population Dynamics of the Highly Halotolerant Species Eucypris mareotica (Fischer,) Tj ETQq0   | 0 OrgBT / | Overlock 10 1 |
| 25 | The role of salinity as an environmental filtering factor in the determination of the Diptera taxonomic composition in the Crimean waters. Knowledge and Management of Aquatic Ecosystems, 2019, , 3.            | 0.5       | 6             |
| 26 | The Effect of Gamma Radiation on Parthenogenetic Artemia (Branchiopoda, Anostraca) Cysts: Nauplius Hatching and Postnauplial Survival under Varying Salinity. Biology Bulletin, 2019, 46, 1390-1396.             | 0.1       | 2             |
| 27 | Long-term changes (1979-2015) in the nematode fauna in Sivash Bay (Sea of Azov), Russia, worldwide the largest hypersaline lagoon, during salinity transformations. Nematology, 2019, 21, 337-347.               | 0.2       | 15            |
| 28 | Does salinity affect body proportions and "size/mass―ratios of highly halotolerant <i>Baeotendipes noctivagus</i> larvae (Diptera, Chironomidae)?. Oceanological and Hydrobiological Studies, 2019, 48, 305-315. | 0.3       | 4             |
| 29 | The political decision caused the drastic ecosystem shift of the Sivash Bay (the Sea of Azov).<br>Quaternary International, 2018, 475, 4-10.   | 0.7       | 31            |
| 30 | Tintinnina (Ciliophora) and Foraminifera in plankton of hypersaline Lagoon Bardawil (Egypt): spatial and temporal variability. Turkish Journal of Zoology, 2018, 42, 218-229.                                    | 0.4       | 15            |
| 31 | Cladophora mats in a Crimean hypersaline lake: structure, dynamics, and inhabiting animals. Journal of Oceanology and Limnology, 2018, 36, 1930-1940.  | 0.6       | 19            |
| 32 | Effect of Salinity on Chironomid Larvae (Diptera, Chironomidae) in Hypersaline Lakes of Crimea.<br>Biology Bulletin, 2018, 45, 1211-1218.  | 0.1       | 6             |
| 33 | How can saline and hypersaline lakes contribute to aquaculture development? A review. Journal of Oceanology and Limnology, 2018, 36, 2002-2009.  | 0.6       | 30            |
| 34 | Anthropogenic Transformation of Kyzyl-Yar Lake in Crimea: Multiyear Research Findings. Arid Ecosystems, 2018, 8, 299-306.  | 0.2       | 11            |
| 35 | Integral Indicators of Variability of Arctodiaptomus salinus (Daday, 1885) (Copepoda, Diaptomidae) and Their Possible Use in Assessing the Population State. Inland Water Biology, 2018, 11, 456-464.            | 0.2       | 2             |
| 36 | Copepoda in the shallow hypersaline Bardawil coastal lake (Egypt): Are there long-term changes in composition and abundance?. Oceanological and Hydrobiological Studies, 2018, 47, 219-229.                      | 0.3       | 10            |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | First data on predation of Eucypris mareotica (Crustacea, Ostracoda) in hypersaline waters. Food Webs, 2018, 16, e00090.  | 0.5 | 6         |
| 38 | Extreme hydrological events destabilize aquatic ecosystems and open doors for alien species. Quaternary International, 2018, 475, 11-15.  | 0.7 | 24        |
| 39 | Gamma Radiation Effect of Partnogenetic Artemia (Branchiopoda, Anostraca) Cysts on Nauplius<br>Hatching and Postnauplius Survival under Different Salinity. Povolzhskii Ekologicheskii Zhurnal,<br>2018, 17, 418-432.                               | 0.0 | 2         |
| 40 | Chironomidae larvae in hypersaline waters of the Crimea: diversity, distribution, abundance and production., 2017, 84, 61-72.   |     | 41        |
| 41 | History of research on biodiversity in Crimean hypersaline waters. Arid Ecosystems, 2017, 7, 52-58.   | 0.2 | 14        |
| 42 | Mesochra rostrata Gurney, 1927 (Copepoda, Harpacticoida) in Sivash Bay (Sea of Azov): Is it a new alien species or a relict of Tethys?. Russian Journal of Biological Invasions, 2017, 8, 244-250.  | 0.2 | 6         |
| 43 | Size polymorphism and fluctuating asymmetry of Artemia (Branchiopoda: Anostraca) populations from the Crimea. Journal of Siberian Federal University - Biology, 2017, 10, 114-126.  | 0.2 | 6         |
| 44 | Brief review of phototrophs in the Crimean hypersaline lakesand lagoons: diversity, ecological role, the possibility of using. Marine Biological Journal, 2017, 2, 80-85.   | 0.3 | 2         |
| 45 | Editorial: the Scientific Сonference "Biodiversity and Productivity of Aquatic Ecosystemsâ€; Dedicated to the 100th Anniversary of Vladimir Nikolaevich Greze (1915–1988). Part 2. Journal of Siberian Federal University - Biology, 2017, 10, 5-8. | 0.2 | 0         |
| 46 | Current invasions of East Asian cyclopoids (Copepoda, Cyclopoida)in Europe: new records from eastern Ukraine. Turkish Journal of Zoology, 2016, 40, 282-285.  | 0.4 | 8         |
| 47 | Long-term changes of physicochemical parameters and benthos in Lake Qarun (Egypt): Can we make a correct forecast of ecosystem future?. Knowledge and Management of Aquatic Ecosystems, 2016, , 18.   | 0.5 | 19        |
| 48 | First Record of Ranatra linearis (Hemiptera, Nepidae) in Hypersaline Water Bodies of the Crimea.<br>Hydrobiological Journal, 2016, 52, 49-53.   | 0.2 | 3         |
| 49 | Cyclopoida in Hypersaline Waters of the Crimea and the World: Diversity, the Impact of Environmental Factors, Ecological Role. Journal of Siberian Federal University - Biology, 2016, 10, 398-408.   | 0.2 | 3         |
| 50 | Transformation of Gulf Sivash (the Sea of Azov) in Conditions of Growing Salinity: Changes of Meiobenthos and Other Ecosystem Components (2013-2015). Journal of Siberian Federal University - Biology, 2016, 10, 452-466.                          | 0.2 | 10        |
| 51 | Second records of the endangered species Artemia urmiana (Anostraca) in the Crimea habitat. Marine<br>Biological Journal, 2016, 1, 75.  | 0.3 | 1         |
| 52 | Intentional introduction of Artemia sinica (Anostraca) in the high-altitude Tibetan lake Dangxiong Co: the new population and consequences for the environment and for humans. Chinese Journal of Oceanology and Limnology, 2015, 33, 1451-1460.    | 0.7 | 27        |
| 53 | Dormant stages of crustaceans as a mechanism of propagation in the extreme and unpredictable environment in the Crimean hypersaline lakes. Chinese Journal of Oceanology and Limnology, 2015, 33, 1362-1367.  | 0.7 | 17        |
| 54 | Does salinity change determine zooplankton variability in the saline Qarun Lake (Egypt)?. Chinese Journal of Oceanology and Limnology, 2015, 33, 1368-1377.   | 0.7 | 26        |

| #  | Article   | IF      | CITATIONS   |
|----|---|---------|-------------|
| 55 | Do copepods inhabit hypersaline waters worldwide? A short review and discussion. Chinese Journal of Oceanology and Limnology, 2015, 33, 1354-1361.  | 0.7     | 31          |
| 56 | Morphometric variability of Arctodiaptomus salinus (Copepoda) in the Mediterranean-Black Sea region. Zoological Research, 2015, 36, 328-36.   | 0.6     | 2           |
| 57 | Current Invasions of Asian Cyclopid Species (Copepoda: Cyclopidae) in Crimea, with Taxonomical and Zoogeographical Remarks on the Hypersaline and Freshwater Fauna. Annales Zoologici, 2014, 64, 109-130. | 0.1     | 42          |
| 58 | The swimming behavior of Artemia (Anostraca): new experimental and observational data. Zoology, 2014, 117, 415-421.   | 0.6     | 17          |
| 59 | Factors determining the average body size of geographically separated Arctodiaptomus salinus (Daday,) Tj ETQq1  | 10.7843 | 14 rgBT /Ov |
| 60 | Title is missing!. Turkish Journal of Fisheries and Aquatic Sciences, 2013, 13, .   | 0.4     | 36          |
| 61 | The intramolecular mobility of poly-n-alkylmethacrylates. Polymer Science USSR, 1975, 17, 675-682.  | 0.2     | 3           |
| 62 | Intramolecular mobility and flexibility of polyamidoacid molecules in solution. Polymer Science USSR, 1975, 17, 3220-3229.  | 0.2     | 2           |
| 63 | Potentiometric titration of polyacrylic acid, polymethacrylic acid and poly-l-glutamic acid. Polymer Science USSR, 1965, 7, 1008-1018.  | 0.2     | 23          |
| 64 | OSTRACODS IN THE PLANKTON OF THE SIVASH BAY (THE SEA OF AZOV) DURING ITS TRANSFORMATION FROM BRACKISH TO HYPERSALINE STATE. Ecologica Montenegrina, 0, 14, 102-108.                                       | 0.5     | 6           |
| 65 | MICROALGAE IN THE DIET OF EUCYPRIS MAREOTICA (CRUSTACEA, OSTRACODA) IN THE HYPERSALINE LAKE CHERSONESSKOYE (CRIMEA). Ecologica Montenegrina, 0, 17, 100-104.  | 0.5     | 3           |