

# Jussi Pihlajamki

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/5382997/jussi-pihlajamaki-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

204  
papers

9,843  
citations

51  
h-index

93  
g-index

219  
ext. papers

11,559  
ext. citations

6.4  
avg, IF

5.62  
L-index

#	Paper	IF	Citations
204	PSD3 downregulation confers protection against fatty liver disease.. <i>Nature Metabolism</i> , <b>2022</b> , 4, 60-75	14.6	1
203	Immigrants Perspectives on healthy life and healthy lifestyle counseling: a focus group study.. <i>Scandinavian Journal of Public Health</i> , <b>2022</b> , 14034948221075021	3	0
202	Digitally Supported Lifestyle Intervention to Prevent Type 2 Diabetes Through Healthy Habits: Secondary Analysis of Long-Term User Engagement Trajectories in a Randomized Controlled Trial.. <i>Journal of Medical Internet Research</i> , <b>2022</b> , 24, e31530	7.6	1
201	Robotic versus hybrid assisted ventral hernia repair: a prospective one-year comparative study of clinical outcomes.. <i>Acta Chirurgica Belgica</i> , <b>2022</b> , 1-20	0.9	
200	Digitalization as an Engine for Change? Building a Vision Pathway towards a Sustainable Health Care System by Using the MLP and Health Economic Decision Modelling. <i>Sustainability</i> , <b>2021</b> , 13, 13007	3.6	1
199	Distinct contributions of metabolic dysfunction and genetic risk factors in the pathogenesis of non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , <b>2021</b> ,	13.4	9
198	Indole-3-Propionic Acid, a Gut-Derived Tryptophan Metabolite, Associates with Hepatic Fibrosis. <i>Nutrients</i> , <b>2021</b> , 13,	6.7	6
197	Endothelial function and concentrations of high-sensitivity C-reactive protein, interleukin-6, and tumor necrosis factor-alpha during a long agonist IVF protocol. <i>Journal of Reproductive Immunology</i> , <b>2021</b> , 148, 103434	4.2	0
196	FADS1 rs174550 genotype and high linoleic acid diet modify plasma PUFA phospholipids in a dietary intervention study. <i>European Journal of Nutrition</i> , <b>2021</b> , 1	5.2	0
195	Integrative analysis of liver-specific non-coding regulatory SNPs associated with the risk of coronary artery disease. <i>American Journal of Human Genetics</i> , <b>2021</b> , 108, 411-430	11	4
194	The FADS1 Genotype Modifies Metabolic Responses to the Linoleic Acid and Alpha-linolenic Acid Containing Plant Oils-Genotype Based Randomized Trial FADSDIET2. <i>Molecular Nutrition and Food Research</i> , <b>2021</b> , 65, e2001004	5.9	5
193	The Importance of Intestinal Length in Triglyceride Metabolism and in Predicting the Outcomes of Comorbidities in Laparoscopic Roux-en-Y Gastric Bypass-a Narrative Review. <i>Obesity Surgery</i> , <b>2021</b> , 31, 3291-3295	3.7	
192	Exome-Wide Association Study on Alanine Aminotransferase Identifies Sequence Variants in the GPAM and APOE Associated With Fatty Liver Disease. <i>Gastroenterology</i> , <b>2021</b> , 160, 1634-1646.e7	13.3	23
191	Sleep-time physiological recovery is associated with eating habits in distressed working-age Finns with overweight: secondary analysis of a randomised controlled trial. <i>Journal of Occupational Medicine and Toxicology</i> , <b>2021</b> , 16, 23	2.7	0
190	Fatal complications after an interrupted gastric bypass operation in a patient with non-alcoholic fatty liver disease and massive obesity: a case report. <i>Journal of Surgical Case Reports</i> , <b>2021</b> , 2021, rjab247	0.6	
189	LPIAT1/MBOAT7 depletion increases triglyceride synthesis fueled by high phosphatidylinositol turnover. <i>Gut</i> , <b>2021</b> , 70, 180-193	19.2	39
188	Dietary Fiber from Oat and Rye Brans Ameliorate Western Diet-Induced Body Weight Gain and Hepatic Inflammation by the Modulation of Short-Chain Fatty Acids, Bile Acids, and Tryptophan Metabolism. <i>Molecular Nutrition and Food Research</i> , <b>2021</b> , 65, e1900580	5.9	14

187	Serum aromatic and branched-chain amino acids associated with NASH demonstrate divergent associations with serum lipids. <i>Liver International</i> , <b>2021</b> , 41, 754-763	7.9	8
186	Protein Phosphatase 1 Regulatory Subunit 3B Genotype at rs4240624 Has a Major Effect on Gallbladder Bile Composition. <i>Hepatology Communications</i> , <b>2021</b> , 5, 244-257	6	1
185	PCSK9 rs11591147 R46L loss-of-function variant protects against liver damage in individuals with NAFLD. <i>Liver International</i> , <b>2021</b> , 41, 321-332	7.9	10
184	Formation and Validation of the Healthy Diet Index (HDI) for Evaluation of Diet Quality in Healthcare. <i>International Journal of Environmental Research and Public Health</i> , <b>2021</b> , 18,	4.6	2
183	Differential Mitochondrial Gene Expression in Adipose Tissue Following Weight Loss Induced by Diet or Bariatric Surgery. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2021</b> , 106, 1312-1324	5.6	1
182	NR1H4 rs35724 G>C variant modulates liver damage in nonalcoholic fatty liver disease. <i>Liver International</i> , <b>2021</b> , 41, 2712-2719	7.9	3
181	Identification of TBX15 as an adipose master trans regulator of abdominal obesity genes. <i>Genome Medicine</i> , <b>2021</b> , 13, 123	14.4	2
180	Comparison of Communication Channels for Large-Scale Type 2 Diabetes Risk Screening and Intervention Recruitment: Empirical Study. <i>JMIR Diabetes</i> , <b>2021</b> , 6, e21356	2.7	2
179	The Effects of Acceptance and Commitment Therapy (ACT) Intervention on Inflammation and Stress Biomarkers: a Randomized Controlled Trial. <i>International Journal of Behavioral Medicine</i> , <b>2020</b> , 27, 539-555	2.6	6
178	The effect of different estradiol levels on carotid artery distensibility during a long agonist IVF protocol. <i>Reproductive Biology and Endocrinology</i> , <b>2020</b> , 18, 44	5	2
177	Genetic variants in the MTHFR are not associated with fatty liver disease. <i>Liver International</i> , <b>2020</b> , 40, 1934-1940	7.9	1
176	Effect of metabolic state on implicit and explicit responses to food in young healthy females. <i>Appetite</i> , <b>2020</b> , 148, 104593	4.5	0
175	Parental metabolic syndrome epigenetically reprograms offspring hepatic lipid metabolism in mice. <i>Journal of Clinical Investigation</i> , <b>2020</b> , 130, 2391-2407	15.9	14
174	Internet-Based Lifestyle Intervention to Prevent Type 2 Diabetes Through Healthy Habits: Design and 6-Month Usage Results of Randomized Controlled Trial. <i>JMIR Diabetes</i> , <b>2020</b> , 5, e15219	2.7	7
173	Diabetic phenotype in mouse and humans reduces the number of microglia around amyloid plaques. <i>Molecular Neurodegeneration</i> , <b>2020</b> , 15, 66	19	6
172	Mucosal-associated invariant T cell alterations during the development of human type 1 diabetes. <i>Diabetologia</i> , <b>2020</b> , 63, 2396-2409	10.3	5
171	Change in abdominal, but not femoral subcutaneous fat CT-radiodensity is associated with improved metabolic profile after bariatric surgery. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , <b>2020</b> , 30, 2363-2371	4.5	4
170	Epigenetic markers associated with metformin response and intolerance in drug-naïve patients with type 2 diabetes. <i>Science Translational Medicine</i> , <b>2020</b> , 12,	17.5	14

169	Oxygen-18 and Carbon-13 isotopes in eCOand erythrocytes carbonic anhydrase activity of Finnish prediabetic population. <i>Journal of Breath Research</i> , <b>2020</b> ,	3.1	1
168	Liver DNA methylation of FADS2 associates with FADS2 genotype. <i>Clinical Epigenetics</i> , <b>2019</b> , 11, 10	7.7	12
167	Total liver phosphatidylcholine content associates with non-alcoholic steatohepatitis and glycine N-methyltransferase expression. <i>Liver International</i> , <b>2019</b> , 39, 1895-1905	7.9	3
166	Decreased plasma serotonin and other metabolite changes in healthy adults after consumption of wholegrain rye: an untargeted metabolomics study. <i>American Journal of Clinical Nutrition</i> , <b>2019</b> , 109, 1630-1639	7	11
165	Digitally supported program for type 2 diabetes risk identification and risk reduction in real-world setting: protocol for the StopDia model and randomized controlled trial. <i>BMC Public Health</i> , <b>2019</b> , 19, 255	4.1	12
164	gene variation bridges atherogenic dyslipidemia with hepatic inflammation in NAFLD patients. <i>Journal of Lipid Research</i> , <b>2019</b> , 60, 1144-1153	6.3	27
163	Healthy Nordic Diet Modulates the Expression of Genes Related to Mitochondrial Function and Immune Response in Peripheral Blood Mononuclear Cells from Subjects with Metabolic Syndrome-A SYSDIET Sub-Study. <i>Molecular Nutrition and Food Research</i> , <b>2019</b> , 63, e1801405	5.9	8
162	Laparoscopic Roux-en-Y gastric bypass surgery influenced pharmacokinetics of several drugs given as a cocktail with the highest impact observed for CYP1A2, CYP2C8 and CYP2E1 substrates. <i>Basic and Clinical Pharmacology and Toxicology</i> , <b>2019</b> , 125, 123-132	3.1	5
161	Prevalence and Risk Factors of Significant Fibrosis in Patients With Nonalcoholic Fatty Liver Without Steatohepatitis. <i>Clinical Gastroenterology and Hepatology</i> , <b>2019</b> , 17, 2310-2319.e6	6.9	42
160	Novel Lipid Long Intervening Noncoding RNA, Oligodendrocyte Maturation-Associated Long Intergenic Noncoding RNA, Regulates the Liver Steatosis Gene Stearoyl-Coenzyme A Desaturase As an Enhancer RNA. <i>Hepatology Communications</i> , <b>2019</b> , 3, 1356-1372	6	17
159	Eating Competence Is Associated with Lower Prevalence of Obesity and Better Insulin Sensitivity in Finnish Adults with Increased Risk for Type 2 Diabetes: The StopDia Study. <i>Nutrients</i> , <b>2019</b> , 12,	6.7	4
158	Association of fatty liver disease with mortality outcomes in an Eastern Finland male cohort. <i>BMJ Open Gastroenterology</i> , <b>2019</b> , 6, e000219	3.9	3
157	Fatty liver index as a predictor of increased risk of cardiometabolic disease: finding from the Kuopio Ischaemic Heart Disease Risk Factor Study Cohort. <i>BMJ Open</i> , <b>2019</b> , 9, e031420	3	2
156	Epigenome-Wide Association Study of Incident Type 2 Diabetes in a British Population: EPIC-Norfolk Study. <i>Diabetes</i> , <b>2019</b> , 68, 2315-2326	0.9	40
155	An Isocaloric Nordic Diet Modulates and Gene Expression in Peripheral Blood Mononuclear Cells in Individuals with Metabolic Syndrome-A SYSDIET Sub-Study. <i>Nutrients</i> , <b>2019</b> , 11,	6.7	9
154	Inflammatory response to dietary linoleic acid depends on FADS1 genotype. <i>American Journal of Clinical Nutrition</i> , <b>2019</b> , 109, 165-175	7	33
153	Incidence, Comorbidities, and Mortality in Idiopathic Normal PressureHydrocephalus. <i>World Neurosurgery</i> , <b>2018</b> , 112, e624-e631	2.1	22
152	Brown adipose tissue lipid metabolism in morbid obesity: Effect of bariatric surgery-induced weight loss. <i>Diabetes, Obesity and Metabolism</i> , <b>2018</b> , 20, 1280-1288	6.7	23

151	Causal relationship of hepatic fat with liver damage and insulin resistance in nonalcoholic fatty liver. <i>Journal of Internal Medicine</i> , <b>2018</b> , 283, 356-370	10.8	140
150	The effects of acceptance and commitment therapy on eating behavior and diet delivered through face-to-face contact and a mobile app: a randomized controlled trial. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , <b>2018</b> , 15, 22	8.4	28
149	Serum, liver and bile sitosterol and sitostanol in obese patients with and without NAFLD. <i>Bioscience Reports</i> , <b>2018</b> , 38,	4.1	4
148	Effects of Genetic Variants on Carboxylesterase 1 Gene Expression, and Clopidogrel Pharmacokinetics and Antiplatelet Effects. <i>Basic and Clinical Pharmacology and Toxicology</i> , <b>2018</b> , 122, 341-345	3.1	8
147	Comprehensive Pharmacogenomic Study Reveals an Important Role of UGT1A3 in Montelukast Pharmacokinetics. <i>Clinical Pharmacology and Therapeutics</i> , <b>2018</b> , 104, 158-168	6.1	12
146	Effector T Cell Resistance to Suppression and STAT3 Signaling during the Development of Human Type 1 Diabetes. <i>Journal of Immunology</i> , <b>2018</b> , 201, 1144-1153	5.3	14
145	Small Intestinal Length Associates with Serum Triglycerides Before and After LRYGB. <i>Obesity Surgery</i> , <b>2018</b> , 28, 3969-3975	3.7	3
144	Association of fatty liver index with the risk of incident cardiovascular disease and acute myocardial infarction. <i>European Journal of Gastroenterology and Hepatology</i> , <b>2018</b> , 30, 1047-1054	2.2	20
143	Increased Liver Fatty Acid Uptake Is Partly Reversed and Liver Fat Content Normalized After Bariatric Surgery. <i>Diabetes Care</i> , <b>2018</b> , 41, 368-371	14.6	18
142	Evaluation of the Effect of Bariatric Surgery-Induced Weight Loss on Knee Gait and Cartilage Degeneration. <i>Journal of Biomechanical Engineering</i> , <b>2018</b> , 140,	2.1	15
141	Sex Differences in the Methylome and Transcriptome of the Human Liver and Circulating HDL-Cholesterol Levels. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2018</b> , 103, 4395-4408	5.6	26
140	Measured energy content of frequently purchased restaurant meals: multi-country cross sectional study. <i>BMJ, The</i> , <b>2018</b> , 363, k4864	5.9	20
139	Associations of serum indolepropionic acid, a gut microbiota metabolite, with type 2 diabetes and low-grade inflammation in high-risk individuals. <i>Nutrition and Diabetes</i> , <b>2018</b> , 8, 35	4.7	75
138	Protein phosphatase 1 regulatory subunit 3B gene variation protects against hepatic fat accumulation and fibrosis in individuals at high risk of nonalcoholic fatty liver disease. <i>Hepatology Communications</i> , <b>2018</b> , 2, 666-675	6	30
137	Molecular evaluation of vitamin D responsiveness of healthy young adults. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , <b>2017</b> , 174, 314-321	5.1	24
136	Sleep of professional athletes: Underexploited potential to improve health and performance. <i>Journal of Sports Sciences</i> , <b>2017</b> , 35, 704-710	3.6	58
135	Obstructive sleep apnea: the effect of bariatric surgery after 12 months. A prospective multicenter trial. <i>Sleep Medicine</i> , <b>2017</b> , 35, 85-90	4.6	45
134	Interorgan cross talk between fatty acid metabolism, tissue inflammation, and FADS2 genotype in humans with obesity. <i>Obesity</i> , <b>2017</b> , 25, 545-552	8	20

133	Human liver epigenetic alterations in non-alcoholic steatohepatitis are related to insulin action. <i>Epigenetics</i> , <b>2017</b> , 12, 287-295	5.7	39
132	Indolepropionic acid and novel lipid metabolites are associated with a lower risk of type 2 diabetes in the Finnish Diabetes Prevention Study. <i>Scientific Reports</i> , <b>2017</b> , 7, 46337	4.9	137
131	Fatty acid uptake and blood flow in adipose tissue compartments of morbidly obese subjects with or without type 2 diabetes: effects of bariatric surgery. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , <b>2017</b> , 313, E175-E182	6	20
130	Fasting serum hippuric acid is elevated after bilberry ( <i>Vaccinium myrtillus</i> ) consumption and associates with improvement of fasting glucose levels and insulin secretion in persons at high risk of developing type 2 diabetes. <i>Molecular Nutrition and Food Research</i> , <b>2017</b> , 61, 1700019	5.9	36
129	Epigenetic alterations in blood mirror age-associated DNA methylation and gene expression changes in human liver. <i>Epigenomics</i> , <b>2017</b> , 9, 105-122	4.4	33
128	Identification and characterization of a FOXA2-regulated transcriptional enhancer at a type 2 diabetes intronic locus that controls GCKR expression in liver cells. <i>Genome Medicine</i> , <b>2017</b> , 9, 63	14.4	16
127	Diabetes medication associates with DNA methylation of metformin transporter genes in the human liver. <i>Clinical Epigenetics</i> , <b>2017</b> , 9, 102	7.7	28
126	Alterations in fatty acid metabolism in response to obesity surgery combined with dietary counseling. <i>Nutrition and Diabetes</i> , <b>2017</b> , 7, e285	4.7	17
125	Hepatic DPP4 DNA Methylation Associates With Fatty Liver. <i>Diabetes</i> , <b>2017</b> , 66, 25-35	0.9	41
124	Serum Plant Sterols Associate with Gallstone Disease Independent of Weight Loss and Non-Alcoholic Fatty Liver Disease. <i>Obesity Surgery</i> , <b>2017</b> , 27, 1284-1291	3.7	4
123	A liquid chromatography-tandem mass spectrometry analysis of nine cytochrome P450 probe drugs and their corresponding metabolites in human serum and urine. <i>Analytical and Bioanalytical Chemistry</i> , <b>2017</b> , 409, 251-268	4.4	12
122	Circulating CXCR5+PD-1+ICOS+ Follicular T Helper Cells Are Increased Close to the Diagnosis of Type 1 Diabetes in Children With Multiple Autoantibodies. <i>Diabetes</i> , <b>2017</b> , 66, 437-447	0.9	68
121	Regulation of alternative splicing in human obesity loci. <i>Obesity</i> , <b>2016</b> , 24, 2033-7	8	9
120	Association of MBOAT7 gene variant with plasma ALT levels in children: the PANIC study. <i>Pediatric Research</i> , <b>2016</b> , 80, 651-655	3.2	34
119	Reply to "Statin treatment for non-alcoholic steatohepatitis". <i>Journal of Hepatology</i> , <b>2016</b> , 64, 242-3	13.4	
118	FADS2 genotype regulates delta-6 desaturase activity and inflammation in human adipose tissue. <i>Journal of Lipid Research</i> , <b>2016</b> , 57, 56-65	6.3	35
117	Fatty acid metabolism is altered in non-alcoholic steatohepatitis independent of obesity. <i>Metabolism: Clinical and Experimental</i> , <b>2016</b> , 65, 655-666	12.7	57
116	Associations of TM6SF2 167K allele with liver enzymes and lipid profile in children: the PANIC Study. <i>Pediatric Research</i> , <b>2016</b> , 79, 684-8	3.2	11

115	Probiotics modulated gut microbiota suppresses hepatocellular carcinoma growth in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2016</b> , 113, E1306-15	11.5	288
114	The MBOAT7-TMC4 Variant rs641738 Increases Risk of Nonalcoholic Fatty Liver Disease in Individuals of European Descent. <i>Gastroenterology</i> , <b>2016</b> , 150, 1219-1230.e6	13.3	347
113	Effect of Bariatric Surgery on Adipose Tissue Glucose Metabolism in Different Depots in Patients With or Without Type 2 Diabetes. <i>Diabetes Care</i> , <b>2016</b> , 39, 292-9	14.6	39
112	Association of plasma fatty acid composition with plasma irisin levels in normal weight and overweight/obese children. <i>Pediatric Obesity</i> , <b>2016</b> , 11, 299-305	4.6	17
111	Gene-diet interaction of a common FADS1 variant with marine polyunsaturated fatty acids for fatty acid composition in plasma and erythrocytes among men. <i>Molecular Nutrition and Food Research</i> , <b>2016</b> , 60, 381-9	5.9	17
110	Plasma cathepsin D correlates with histological classifications of fatty liver disease in adults and responds to intervention. <i>Scientific Reports</i> , <b>2016</b> , 6, 38278	4.9	22
109	Prevalence of Obstructive Sleep Apnoea Among Patients Admitted for Bariatric Surgery. A Prospective Multicentre Trial. <i>Obesity Surgery</i> , <b>2016</b> , 26, 1384-90	3.7	38
108	High perceived stress is associated with unfavorable eating behavior in overweight and obese Finns of working age. <i>Appetite</i> , <b>2016</b> , 103, 249-258	4.5	48
107	DNA methylation of loci within ABCG1 and PHOSPHO1 in blood DNA is associated with future type 2 diabetes risk. <i>Epigenetics</i> , <b>2016</b> , 11, 482-8	5.7	99
106	Transmembrane 6 superfamily member 2 gene variant disentangles nonalcoholic steatohepatitis from cardiovascular disease. <i>Hepatology</i> , <b>2015</b> , 61, 506-14	11.2	311
105	Markers of cholesterol metabolism as biomarkers in predicting diabetes in the Finnish Diabetes Prevention Study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , <b>2015</b> , 25, 635-42	4.5	11
104	The 148 M allele of the PNPLA3 is associated with plasma irisin levels in a population sample of Caucasian children: The PANIC Study. <i>Metabolism: Clinical and Experimental</i> , <b>2015</b> , 64, 793-6	12.7	14
103	Ketone body production is differentially altered in steatosis and non-alcoholic steatohepatitis in obese humans. <i>Liver International</i> , <b>2015</b> , 35, 1853-61	7.9	48
102	Associations of I148M variant in PNPLA3 gene with plasma ALT levels during 2-year follow-up in normal weight and overweight children: the PANIC Study. <i>Pediatric Obesity</i> , <b>2015</b> , 10, 84-90	4.6	15
101	Dietary polyunsaturated fatty acids and the Pro12Ala polymorphisms of PPARG regulate serum lipids through divergent pathways: a randomized crossover clinical trial. <i>Genes and Nutrition</i> , <b>2015</b> , 10, 43	4.3	11
100	Plasma fatty acids as predictors of glycaemia and type 2 diabetes. <i>Diabetologia</i> , <b>2015</b> , 58, 2533-44	10.3	70
99	Cross-linking of sodium caseinate-structured emulsion with transglutaminase alters postprandial metabolic and appetite responses in healthy young individuals. <i>British Journal of Nutrition</i> , <b>2015</b> , 114, 418-29	3.6	6
98	Persistent organic pollutants and non-alcoholic fatty liver disease in morbidly obese patients: a cohort study. <i>Environmental Health</i> , <b>2015</b> , 14, 79	6	39

97	MFAP5 is related to obesity-associated adipose tissue and extracellular matrix remodeling and inflammation. <i>Obesity</i> , <b>2015</b> , 23, 1371-8	8	20
96	Statin use and non-alcoholic steatohepatitis in at risk individuals. <i>Journal of Hepatology</i> , <b>2015</b> , 63, 705-12	3.4	227
95	Epigenetic Alterations in Human Liver From Subjects With Type 2 Diabetes in Parallel With Reduced Folate Levels. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2015</b> , 100, E1491-501	5.6	109
94	Plasma IL-1 receptor antagonist levels correlate with the development of non-alcoholic steatohepatitis. <i>Biomarkers in Medicine</i> , <b>2015</b> , 9, 1301-9	2.3	3
93	Healthy Nordic diet downregulates the expression of genes involved in inflammation in subcutaneous adipose tissue in individuals with features of the metabolic syndrome. <i>American Journal of Clinical Nutrition</i> , <b>2015</b> , 101, 228-39	7	38
92	Effect of bariatric surgery on liver glucose metabolism in morbidly obese diabetic and non-diabetic patients. <i>Journal of Hepatology</i> , <b>2014</b> , 60, 377-83	13.4	70
91	Specific collagen XVIII isoforms promote adipose tissue accrual via mechanisms determining adipocyte number and affect fat deposition. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2014</b> , 111, E3043-52	11.5	31
90	DNA methylation in obesity and type 2 diabetes. <i>Annals of Medicine</i> , <b>2014</b> , 46, 103-13	1.5	56
89	The effectiveness and applicability of different lifestyle interventions for enhancing wellbeing: the study design for a randomized controlled trial for persons with metabolic syndrome risk factors and psychological distress. <i>BMC Public Health</i> , <b>2014</b> , 14, 310	4.1	31
88	High-fat diet increases tau expression in the brain of T2DM and AD mice independently of peripheral metabolic status. <i>Journal of Nutritional Biochemistry</i> , <b>2014</b> , 25, 634-41	6.3	40
87	A population-based study on the prevalence of NASH using scores validated against liver histology. <i>Journal of Hepatology</i> , <b>2014</b> , 60, 839-46	13.4	89
86	Lipoprotein subclass metabolism in nonalcoholic steatohepatitis. <i>Journal of Lipid Research</i> , <b>2014</b> , 55, 2676-84	6.3	51
85	Postprandial glucose metabolism and SCFA after consuming wholegrain rye bread and wheat bread enriched with bioprocessed rye bran in individuals with mild gastrointestinal symptoms. <i>Nutrition Journal</i> , <b>2014</b> , 13, 104	4.3	26
84	Adipose tissue INSR splicing in humans associates with fasting insulin level and is regulated by weight loss. <i>Diabetologia</i> , <b>2014</b> , 57, 347-51	10.3	30
83	Genetic risk score does not predict the outcome of obesity surgery. <i>Obesity Surgery</i> , <b>2014</b> , 24, 128-33	3.7	16
82	Desmosterol in human nonalcoholic steatohepatitis. <i>Hepatology</i> , <b>2013</b> , 58, 976-82	11.2	34
81	Downregulation of CPPED1 expression improves glucose metabolism in vitro in adipocytes. <i>Diabetes</i> , <b>2013</b> , 62, 3747-50	0.9	18
80	Comparative nontargeted profiling of metabolic changes in tissues and biofluids in high-fat diet-fed Ossabaw pig. <i>Journal of Proteome Research</i> , <b>2013</b> , 12, 3980-92	5.6	27



79	Association of ketone body levels with hyperglycemia and type 2 diabetes in 9,398 Finnish men. <i>Diabetes</i> , <b>2013</b> , 62, 3618-26	0.9	77
78	Link between GIP and osteopontin in adipose tissue and insulin resistance. <i>Diabetes</i> , <b>2013</b> , 62, 2088-94	0.9	64
77	Regulation of alternative splicing in obesity and weight loss. <i>Adipocyte</i> , <b>2013</b> , 2, 143-7	3.2	12
76	Non-cholesterol sterol levels predict hyperglycemia and conversion to type 2 diabetes in Finnish men. <i>PLoS ONE</i> , <b>2013</b> , 8, e67406	3.7	13
75	Glomerular filtration rate and parathyroid hormone are associated with 1,25-dihydroxyvitamin D in men without chronic kidney disease. <i>Journal of Internal Medicine</i> , <b>2012</b> , 271, 573-80	10.8	5
74	Adipose tissue TCF7L2 splicing is regulated by weight loss and associates with glucose and fatty acid metabolism. <i>Diabetes</i> , <b>2012</b> , 61, 2807-13	0.9	57
73	Serum interleukin 1 receptor antagonist as an independent marker of non-alcoholic steatohepatitis in humans. <i>Journal of Hepatology</i> , <b>2012</b> , 56, 663-70	13.4	69
72	Response to Brosch et al. <i>Cell Metabolism</i> , <b>2012</b> , 15, 267-269	24.6	4
71	Conjugated bile acids associate with altered rates of glucose and lipid oxidation after Roux-en-Y gastric bypass. <i>Obesity Surgery</i> , <b>2012</b> , 22, 1473-80	3.7	127
70	Impact of obesity and associated diseases on outcome after laparoscopic cholecystectomy. <i>Surgical Laparoscopy, Endoscopy and Percutaneous Techniques</i> , <b>2012</b> , 22, 509-13	1.3	25
69	Hyperglycemia and a common variant of GCKR are associated with the levels of eight amino acids in 9,369 Finnish men. <i>Diabetes</i> , <b>2012</b> , 61, 1895-902	0.9	195
68	Expression of the splicing factor gene SFRS10 is reduced in human obesity and contributes to enhanced lipogenesis. <i>Cell Metabolism</i> , <b>2011</b> , 14, 208-18	24.6	98
67	Association of indices of liver and adipocyte insulin resistance with 19 confirmed susceptibility loci for type 2 diabetes in 6,733 non-diabetic Finnish men. <i>Diabetologia</i> , <b>2011</b> , 54, 563-71	10.3	34
66	PKC $\beta$ regulates hepatic insulin sensitivity and hepatosteatosis in mice and humans. <i>Journal of Clinical Investigation</i> , <b>2011</b> , 121, 2504-17	15.9	98
65	Diverse associations of 25-hydroxyvitamin D and 1,25-dihydroxy-vitamin D with dyslipidaemias. <i>Journal of Internal Medicine</i> , <b>2010</b> , 268, 604-10	10.8	80
64	SIRT1 mRNA expression may be associated with energy expenditure and insulin sensitivity. <i>Diabetes</i> , <b>2010</b> , 59, 829-35	0.9	83
63	Insulin sensitivity regulates cholesterol metabolism to a greater extent than obesity: lessons from the METSIM Study. <i>Journal of Lipid Research</i> , <b>2010</b> , 51, 2422-7	6.3	48
62	Changes in cytokine levels during acute hyperinsulinemia in offspring of type 2 diabetic subjects. <i>Atherosclerosis</i> , <b>2010</b> , 210, 536-41	3.1	15

61	Cholesterol absorption decreases after Roux-en-Y gastric bypass but not after gastric banding. <i>Metabolism: Clinical and Experimental</i> , <b>2010</b> , 59, 866-72	12.7	64
60	Thyroid hormone-related regulation of gene expression in human fatty liver. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2009</b> , 94, 3521-9	5.6	108
59	The metabolism of plant sterols is disturbed in postmenopausal women with coronary artery disease. <i>Metabolism: Clinical and Experimental</i> , <b>2009</b> , 58, 401-7	12.7	25
58	The Pro12Ala polymorphism of the PPARgamma2 gene is associated with hepatic glucose uptake during hyperinsulinemia in subjects with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , <b>2009</b> , 58, 541-6	12.7	8
57	Serum bile acids are higher in humans with prior gastric bypass: potential contribution to improved glucose and lipid metabolism. <i>Obesity</i> , <b>2009</b> , 17, 1671-7	8	431
56	Markers of endothelial dysfunction and low-grade inflammation are associated in the offspring of type 2 diabetic subjects. <i>Atherosclerosis</i> , <b>2008</b> , 197, 271-7	3.1	27
55	Single-nucleotide polymorphism rs7754840 of CDKAL1 is associated with impaired insulin secretion in nondiabetic offspring of type 2 diabetic subjects and in a large sample of men with normal glucose tolerance. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2008</b> , 93, 1924-30	5.6	71
54	Cholesterol synthesis prevails over absorption in metabolic syndrome. <i>Translational Research</i> , <b>2007</b> , 149, 310-6	11	44
53	Single nucleotide polymorphisms of the MCHR1 gene do not affect metabolism in humans. <i>Obesity</i> , <b>2007</b> , 15, 2902-7	8	8
52	Single nucleotide polymorphisms of the melanocortin-3 receptor gene are associated with substrate oxidation and first-phase insulin secretion in offspring of type 2 diabetic subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2007</b> , 92, 1112-7	5.6	16
51	Serum adipokines are associated with cholesterol metabolism in the metabolic syndrome. <i>Clinica Chimica Acta</i> , <b>2007</b> , 383, 126-32	6.2	20
50	Cholesterol metabolism and non-cholesterol sterol distribution in lipoproteins of type 1 diabetes: the effect of improved glycemic control. <i>Atherosclerosis</i> , <b>2007</b> , 194, 465-72	3.1	9
49	Common polymorphisms of calpain-10 are associated with abdominal obesity in subjects at high risk of type 2 diabetes. <i>Diabetologia</i> , <b>2006</b> , 49, 1560-6	10.3	16
48	Association analysis of peroxisome proliferator-activated receptor gamma polymorphisms and late onset Alzheimer's disease in the Finnish population. <i>Dementia and Geriatric Cognitive Disorders</i> , <b>2006</b> , 22, 449-53	2.6	26
47	Changes in inflammatory cytokines are related to impaired glucose tolerance in offspring of type 2 diabetic subjects. <i>Diabetes Care</i> , <b>2006</b> , 29, 2714-20	14.6	63
46	Gene dose effect of the DQB1*0201 allele contributes to severity of coeliac disease. <i>Scandinavian Journal of Gastroenterology</i> , <b>2006</b> , 41, 191-9	2.4	69
45	HLA genotyping is useful in the evaluation of the risk for coeliac disease in the 1st-degree relatives of patients with coeliac disease. <i>Scandinavian Journal of Gastroenterology</i> , <b>2006</b> , 41, 1299-304	2.4	27
44	Chapter 12 Regulation of PGC-1 in humans with insulin resistance and type 2 diabetes: Functional implications. <i>Advances in Molecular and Cellular Endocrinology</i> , <b>2006</b> , 5, 233-253		1

43	Endothelial function in hypercholesterolemic subjects: Effects of plant stanol and sterol esters. <i>Atherosclerosis</i> , <b>2006</b> , 188, 425-32	3.1	55
42	Visceral obesity is associated with high levels of serum squalene. <i>Obesity</i> , <b>2006</b> , 14, 1155-63	8	24
41	Interleukin-6 promoter polymorphism and late-onset Alzheimer's disease in the Finnish population. <i>Journal of Neurogenetics</i> , <b>2005</b> , 19, 155-61	1.6	22
40	The effect of the Ala12 allele of the peroxisome proliferator-activated receptor-gamma2 gene on skeletal muscle glucose uptake depends on obesity: a positron emission tomography study. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2005</b> , 90, 4249-54	5.6	26
39	High amount of visceral fat mass is associated with multiple metabolic changes in offspring of type 2 diabetic patients. <i>International Journal of Obesity</i> , <b>2005</b> , 29, 1464-70	5.5	15
38	Haplotypes of PPARGC1A are associated with glucose tolerance, body mass index and insulin sensitivity in offspring of patients with type 2 diabetes. <i>Diabetologia</i> , <b>2005</b> , 48, 1331-4	10.3	21
37	Single nucleotide polymorphisms in the peroxisome proliferator-activated receptor delta gene are associated with skeletal muscle glucose uptake. <i>Diabetes</i> , <b>2005</b> , 54, 3587-91	0.9	51
36	The G-250A promoter polymorphism of the hepatic lipase gene predicts the conversion from impaired glucose tolerance to type 2 diabetes mellitus: the Finnish Diabetes Prevention Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2004</b> , 89, 2019-23	5.6	58
35	Insulin resistance is associated with increased cholesterol synthesis and decreased cholesterol absorption in normoglycemic men. <i>Journal of Lipid Research</i> , <b>2004</b> , 45, 507-12	6.3	134
34	Multiple abnormalities in glucose and energy metabolism and coordinated changes in levels of adiponectin, cytokines, and adhesion molecules in subjects with metabolic syndrome. <i>Circulation</i> , <b>2004</b> , 110, 3842-8	16.7	207
33	Polymorphisms in the ABCG5 and ABCG8 genes associate with cholesterol absorption and insulin sensitivity. <i>Journal of Lipid Research</i> , <b>2004</b> , 45, 1660-5	6.3	125
32	Aldose reductase gene polymorphisms and peripheral nerve function in patients with type 2 diabetes. <i>Diabetes Care</i> , <b>2004</b> , 27, 2021-6	14.6	27
31	Polymorphisms of the SUR1 (ABCC8) and Kir6.2 (KCNJ11) genes predict the conversion from impaired glucose tolerance to type 2 diabetes. The Finnish Diabetes Prevention Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2004</b> , 89, 6286-90	5.6	76
30	The Val103Ile polymorphism of melanocortin-4 receptor regulates energy expenditure and weight gain. <i>Obesity</i> , <b>2004</b> , 12, 1060-6		22
29	The Pro12Ala polymorphism of the PPAR gamma 2 gene regulates weight from birth to adulthood. <i>Obesity</i> , <b>2004</b> , 12, 187-90		45
28	Common polymorphisms in the genes regulating the early insulin signalling pathway: effects on weight change and the conversion from impaired glucose tolerance to Type 2 diabetes. The Finnish Diabetes Prevention Study. <i>Diabetologia</i> , <b>2004</b> , 47, 871-7	10.3	50
27	Genomewide search and association studies in a Finnish celiac disease population: Identification of a novel locus and replication of the HLA and CTLA4 loci <b>2004</b> , 130A, 345-50		27
26	The C-174G promoter polymorphism of the IL-6 gene affects energy expenditure and insulin sensitivity. <i>Diabetes</i> , <b>2003</b> , 52, 558-61	0.9	122

25	Insulin resistance is related to left ventricular hypertrophy in patients with polycystic kidney disease type 1. <i>American Journal of Kidney Diseases</i> , <b>2003</b> , 41, 1219-24	7.4	26
24	The effect of the -308A allele of the TNF-alpha gene on insulin action is dependent on obesity. <i>Obesity</i> , <b>2003</b> , 11, 912-7		29
23	The Leu7Pro polymorphism of the neuropeptide Y gene regulates free fatty acid metabolism. <i>Metabolism: Clinical and Experimental</i> , <b>2003</b> , 52, 643-6	12.7	17
22	Promoter polymorphisms of the TNF-alpha (G-308A) and IL-6 (C-174G) genes predict the conversion from impaired glucose tolerance to type 2 diabetes: the Finnish Diabetes Prevention Study. <i>Diabetes</i> , <b>2003</b> , 52, 1872-6	0.9	209
21	The K121Q polymorphism of the PC-1 gene is associated with insulin resistance but not with dyslipidemia. <i>Diabetes Care</i> , <b>2003</b> , 26, 464-7	14.6	63
20	Mutations in the cardiac myosin-binding protein C gene are the predominant cause of familial hypertrophic cardiomyopathy in eastern Finland. <i>Journal of Molecular Medicine</i> , <b>2002</b> , 80, 412-22	5.5	57
19	Apolipoprotein E gene promoter (-219G/T) polymorphism is associated with premature coronary heart disease. <i>Journal of Molecular Medicine</i> , <b>2001</b> , 79, 732-7	5.5	39
18	The hormone sensitive lipase gene in familial combined hyperlipidemia and insulin resistance. <i>European Journal of Clinical Investigation</i> , <b>2001</b> , 31, 302-8	4.6	28
17	A major gene effect on fasting insulin and insulin sensitivity in familial combined hyperlipidemia. <i>Diabetes</i> , <b>2001</b> , 50, 2396-401	0.9	5
16	Association of angiotensin converting enzyme and plasminogen activator inhibitor-1 promoter gene polymorphisms with features of the insulin resistance syndrome in patients with premature coronary heart disease. <i>Atherosclerosis</i> , <b>2001</b> , 157, 57-64	3.1	42
15	Sulfonylurea receptor 1 gene variants are associated with gestational diabetes and type 2 diabetes but not with altered secretion of insulin. <i>Diabetes Care</i> , <b>2000</b> , 23, 70-3	14.6	74
14	Impaired free fatty acid suppression during hyperinsulinemia is a characteristic finding in familial combined hyperlipidemia, but insulin resistance is observed only in hypertriglyceridemic patients. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>2000</b> , 20, 164-70	9.4	32
13	G-250A substitution in promoter of hepatic lipase gene is associated with dyslipidemia and insulin resistance in healthy control subjects and in members of families with familial combined hyperlipidemia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>2000</b> , 20, 1789-95	9.4	64
12	The Pro12A1a substitution in the peroxisome proliferator activated receptor gamma 2 is associated with an insulin-sensitive phenotype in families with familial combined hyperlipidemia and in nondiabetic elderly subjects with dyslipidemia. <i>Atherosclerosis</i> , <b>2000</b> , 151, 567-74	3.1	41
11	Two polymorphisms in the peroxisome proliferator-activated receptor-gamma gene are associated with severe overweight among obese women. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>1999</b> , 84, 3708-12	5.6	186
10	Genomewide scan for familial combined hyperlipidemia genes in Finnish families, suggesting multiple susceptibility loci influencing triglyceride, cholesterol, and apolipoprotein B levels. <i>American Journal of Human Genetics</i> , <b>1999</b> , 64, 1453-63	11	126
9	A Pro12Ala substitution in PPARgamma2 associated with decreased receptor activity, lower body mass index and improved insulin sensitivity. <i>Nature Genetics</i> , <b>1998</b> , 20, 284-7	36.3	1117
8	Linkage of familial combined hyperlipidaemia to chromosome 1q21-q23. <i>Nature Genetics</i> , <b>1998</b> , 18, 369-373	36.3	226

7	Different regulation of free fatty acid levels and glucose oxidation by the Trp64Arg polymorphism of the beta3-adrenergic receptor gene and the promoter variant (A-3826G) of the uncoupling protein 1 gene in familial combined hyperlipidemia. <i>Metabolism: Clinical and Experimental</i> , <b>1998</b> , 47, 1397-402	12.7	18
6	The cardiac beta-myosin heavy chain gene is not the predominant gene for hypertrophic cardiomyopathy in the Finnish population. <i>Journal of the American College of Cardiology</i> , <b>1998</b> , 32, 1709-15	15.1	54
5	Impaired insulin-stimulated glucose oxidation and free fatty acid suppression in patients with familial combined hyperlipidemia: a precursor defect for dyslipidemia?. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>1998</b> , 18, 1548-53	9.4	28
4	The Ala54Thr polymorphism of the fatty acid binding protein 2 gene does not influence insulin sensitivity in Finnish nondiabetic and NIDDM subjects. <i>Diabetes</i> , <b>1997</b> , 46, 711-2	0.9	36
3	The Trp64Arg polymorphism of the beta 3-Adrenergic receptor gene. Lack of association with NIDDM and features of insulin resistance syndrome. <i>Diabetes Care</i> , <b>1997</b> , 20, 1319-23	14.6	56
2	Codon 54 polymorphism of the human intestinal fatty acid binding protein 2 gene is associated with dyslipidemias but not with insulin resistance in patients with familial combined hyperlipidemia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , <b>1997</b> , 17, 1039-44	9.4	45
1	Robotic versus hybrid assisted ventral hernia repair: a prospective one-year comparative study of clinical outcomes. <i>Acta Chirurgica Belgica</i> , 1-7	0.9	