

Michael W Krause

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/539517/publications.pdf>

Version: 2024-02-01

65
papers

5,553
citations

126858
33
h-index

106281
65
g-index

70
all docs

70
docs citations

70
times ranked

7020
citing authors

#	ARTICLE	IF	CITATIONS
1	A trans-spliced leader sequence on actin mRNA in <i>C. elegans</i> . <i>Cell</i> , 1987, 49, 753-761.	13.5	683
2	O-Linked GlcNAc Transferase Is a Conserved Nucleocytoplasmic Protein Containing Tetratricopeptide Repeats. <i>Journal of Biological Chemistry</i> , 1997, 272, 9316-9324.	1.6	462
3	linking metabolism to epigenetics through O-GlcNAcylation. <i>Nature Reviews Molecular Cell Biology</i> , 2012, 13, 312-321.	16.1	364
4	Scalable and Versatile Genome Editing Using Linear DNAs with Microhomology to Cas9 Sites in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , 2014, 198, 1347-1356.	1.2	292
5	The hexosamine signaling pathway: O-GlcNAc cycling in feast or famine. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2010, 1800, 80-95.	1.1	284
6	Haem homeostasis is regulated by the conserved and concerted functions of HRG-1 proteins. <i>Nature</i> , 2008, 453, 1127-1131.	13.7	275
7	The <i>pie-1</i> and <i>mex-1</i> genes and maternal control of blastomere identity in early <i>C. elegans</i> embryos. <i>Cell</i> , 1992, 70, 163-176.	13.5	231
8	CeMyoD accumulation defines the body wall muscle cell fate during <i>C. elegans</i> embryogenesis. <i>Cell</i> , 1990, 63, 907-919.	13.5	211
9	A <i>Caenorhabditis elegans</i> model of insulin resistance: Altered macronutrient storage and dauer formation in an OGT-1 knockout. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 11266-11271.	3.3	208
10	RNA editing by ADARs is important for normal behavior in <i>Caenorhabditis elegans</i> . <i>EMBO Journal</i> , 2002, 21, 6025-6035.	3.5	196
11	<i>Caenorhabditis elegans</i> ortholog of a diabetes susceptibility locus: <i>oga-1</i> (O-GlcNAcase) knockout impacts O-GlcNAc cycling, metabolism, and dauer. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 11952-11957.	3.3	151
12	Dynamic O-GlcNAc cycling at promoters of <i>Caenorhabditis elegans</i> genes regulating longevity, stress, and immunity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 7413-7418.	3.3	136
13	ELT-2 is the predominant transcription factor controlling differentiation and function of the <i>C. elegans</i> intestine, from embryo to adult. <i>Developmental Biology</i> , 2009, 327, 551-565.	0.9	129
14	Sequence analysis of the complete <i>Caenorhabditis elegans</i> myosin heavy chain gene family. <i>Journal of Molecular Biology</i> , 1989, 205, 603-613.	2.0	113
15	The myogenic potency of HLH-1 reveals wide-spread developmental plasticity in early <i>C. elegans</i> embryos. <i>Development (Cambridge)</i> , 2005, 132, 1795-1805.	1.2	103
16	O-GlcNAc cycling: Emerging roles in development and epigenetics. <i>Seminars in Cell and Developmental Biology</i> , 2010, 21, 646-654.	2.3	101
17	Elements Regulating Cell- and Stage-Specific Expression of the <i>C. elegans</i> MyoD Family Homolog <i>hlh-1</i> . <i>Developmental Biology</i> , 1994, 166, 133-148.	0.9	99
18	Defining the transcriptional redundancy of early bodywall muscle development in <i>C. elegans</i> : evidence for a unified theory of animal muscle development. <i>Genes and Development</i> , 2006, 20, 3395-3406.	2.7	98

#	ARTICLE	IF	CITATIONS
19	Dopamine and Glutamate Induce Distinct Striatal Splice Forms of Ania-6, an RNA Polymerase II-Associated Cyclin. <i>Neuron</i> , 2001, 32, 277-287.	3.8	91
20	Wild-type and mutant actin genes in <i>Caenorhabditis elegans</i> . <i>Journal of Molecular Biology</i> , 1989, 208, 381-392.	2.0	90
21	O-GlcNAc cycling mutants modulate proteotoxicity in <i>Caenorhabditis elegans</i> models of human neurodegenerative diseases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 17669-17674.	3.3	86
22	Selenocysteine-Containing Thioredoxin Reductase in <i>C. elegans</i> . <i>Biochemical and Biophysical Research Communications</i> , 1999, 259, 244-249.	1.0	82
23	The embryonic muscle transcriptome of <i>Caenorhabditis elegans</i> . <i>Genome Biology</i> , 2007, 8, R188.	3.8	75
24	O-Linked-N-Acetylglucosamine Cycling and Insulin Signaling Are Required for the Glucose Stress Response in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , 2011, 188, 369-382.	1.2	66
25	Insulinoma-Associated Protein IA-2, a Vesicle Transmembrane Protein, Genetically Interacts with UNC-31/CAPS and Affects Neurosecretion in <i>Caenorhabditis elegans</i> . <i>Journal of Neuroscience</i> , 2004, 24, 3115-3124.	1.7	63
26	Cyclin E expression during development in <i>caenorhabditis elegans</i> . <i>Developmental Biology</i> , 2003, 254, 102-115.	0.9	53
27	NHR-23 dependent collagen and hedgehog-related genes required for molting. <i>Biochemical and Biophysical Research Communications</i> , 2011, 413, 515-520.	1.0	49
28	DNA rearrangements of the actin gene cluster in <i>Caenorhabditis-elegans</i> accompany reversion of three muscle mutants. <i>Journal of Molecular Biology</i> , 1984, 180, 497-513.	2.0	48
29	Evolutionary Conservation of MyoD Function and Differential Utilization of E Proteins. <i>Developmental Biology</i> , 1999, 208, 465-472.	0.9	42
30	Use of an Activated Beta-Catenin to Identify Wnt Pathway Target Genes in <i>Caenorhabditis elegans</i> , Including a Subset of Collagen Genes Expressed in Late Larval Development. <i>G3: Genes, Genomes, Genetics</i> , 2014, 4, 733-747.	0.8	39
31	Caudal-like PAL-1 directly activates the bodywall muscle module regulator <i>hlh-1</i> in <i>C. elegans</i> to initiate the embryonic muscle gene regulatory network. <i>Development (Cambridge)</i> , 2009, 136, 1241-1249.	1.2	38
32	Chapter 20 Transcription and Translation. <i>Methods in Cell Biology</i> , 1995, , 483-512.	0.5	37
33	Heme Utilization in the <i>Caenorhabditis elegans</i> Hypodermal Cells Is Facilitated by Heme-responsive Gene-2. <i>Journal of Biological Chemistry</i> , 2012, 287, 9601-9612.	1.6	37
34	The MADS-Box Factor CeMEF2 Is Not Essential for <i>Caenorhabditis elegans</i> Myogenesis and Development. <i>Developmental Biology</i> , 2000, 223, 431-440.	0.9	34
35	Transcriptional regulation of gene expression in <i>C. elegans</i> . <i>WormBook</i> , 2013, , 1-31.	5.3	34
36	Genome-Wide Analysis Reveals Novel Genes Essential for Heme Homeostasis in <i>Caenorhabditis elegans</i> . <i>PLoS Genetics</i> , 2010, 6, e1001044.	1.5	32

#	ARTICLE	IF	CITATIONS
37	Chapter 21 Techniques for Analyzing Transcription and Translation. <i>Methods in Cell Biology</i> , 1995, 48, 513-529.	0.5	27
38	A Novel H ⁺ -coupled Oligopeptide Transporter (OPT3) from <i>Caenorhabditis elegans</i> with a Predominant Function as a H ⁺ Channel and an Exclusive Expression in Neurons. <i>Journal of Biological Chemistry</i> , 2000, 275, 9563-9571.	1.6	27
39	MyoD and myogenesis in <i>C. elegans</i> . <i>BioEssays</i> , 1995, 17, 219-228.	1.2	26
40	Promotion of Bone Morphogenetic Protein Signaling by Tetraspanins and Glycosphingolipids. <i>PLoS Genetics</i> , 2015, 11, e1005221.	1.5	26
41	Perilipin-related protein regulates lipid metabolism in <i>C. elegans</i> . <i>PeerJ</i> , 2015, 3, e1213.	0.9	25
42	A Widespread Distribution of Genomic CeMyoD Binding Sites Revealed and Cross Validated by ChIP-Chip and ChIP-Seq Techniques. <i>PLoS ONE</i> , 2010, 5, e15898.	1.1	24
43	A Genetic Analysis of the <i>Caenorhabditis elegans</i> Detoxification Response. <i>Genetics</i> , 2017, 206, 939-952.	1.2	21
44	Inter-organ signalling by HRG-7 promotes systemic haem homeostasis. <i>Nature Cell Biology</i> , 2017, 19, 799-807.	4.6	21
45	Hrg1 promotes heme-iron recycling during hemolysis in the zebrafish kidney. <i>PLoS Genetics</i> , 2018, 14, e1007665.	1.5	21
46	Characterization of a dominant negative <i>C. elegans</i> Twist mutant protein with implications for human Saethre-Chotzen syndrome. <i>Development (Cambridge)</i> , 2002, 129, 2761-2772.	1.2	20
47	Identification of Wnt Pathway Target Genes Regulating the Division and Differentiation of Larval Seam Cells and Vulval Precursor Cells in <i>Caenorhabditis elegans</i> . <i>G3: Genes, Genomes, Genetics</i> , 2015, 5, 1551-1566.	0.8	18
48	Functional conservation of nematode and vertebrate myogenic regulatory factors. <i>Journal of Cell Science</i> , 1992, 1992, 111-115.	1.2	17
49	Transgenic <i>C. elegans</i> Dauer Larvae Expressing Hookworm Phospho Null DAF-16/FoxO Exit Dauer. <i>PLoS ONE</i> , 2011, 6, e25996.	1.1	16
50	Disruption of O-GlcNAc Cycling in <i>C. elegans</i> Perturbs Nucleotide Sugar Pools and Complex Glycans. <i>Frontiers in Endocrinology</i> , 2014, 5, 197.	1.5	15
51	X Chromosome Crossover Formation and Genome Stability in <i>Caenorhabditis elegans</i> Are Independently Regulated by <i>xnd-1</i> . <i>G3: Genes, Genomes, Genetics</i> , 2016, 6, 3913-3925.	0.8	15
52	Nutrient-Driven O-GlcNAcylation at Promoters Impacts Genome-Wide RNA Pol II Distribution. <i>Frontiers in Endocrinology</i> , 2018, 9, 521.	1.5	13
53	Somatic muscle specification during embryonic and post-embryonic development in the nematode <i>C. elegans</i> . <i>Wiley Interdisciplinary Reviews: Developmental Biology</i> , 2012, 1, 203-214.	5.9	11
54	The Paired-box protein PAX-3 regulates the choice between lateral and ventral epidermal cell fates in <i>C. elegans</i> . <i>Developmental Biology</i> , 2016, 412, 191-207.	0.9	11

#	ARTICLE	IF	CITATIONS
55	<i>Caenorhabditis elegans</i> DAF-2 as a Model for Human Insulin Receptoropathies. G3: Genes, Genomes, Genetics, 2017, 7, 257-268.	0.8	10
56	Loss of the Transcriptional Repressor PAG-3/Gfi-1 Results in Enhanced Neurosecretion that is Dependent on the Dense-Core Vesicle Membrane Protein IDA-1/IA-2. PLoS Genetics, 2009, 5, e1000447.	1.5	9
57	Multiple Ribonuclease Hâ€œEncoding Genes in the <i>Caenorhabditis elegans</i> Genome Contrasts with the Two Typical Ribonuclease Hâ€œEncoding Genes in the Human Genome. Molecular Biology and Evolution, 2002, 19, 1910-1919.	3.5	7
58	New Roles for the Heterochronic Transcription Factor LIN-29 in Cuticle Maintenance and Lipid Metabolism at the Larval-to-Adult Transition in <i>Caenorhabditis elegans</i> . Genetics, 2020, 214, 669-690.	1.2	7
59	GEI-8, a Homologue of Vertebrate Nuclear Receptor Corepressor NCoR/SMRT, Regulates Gonad Development and Neuronal Functions in <i>Caenorhabditis elegans</i> . PLoS ONE, 2013, 8, e58462.	1.1	7
60	Myogenic conversion and transcriptional profiling of embryonic blastomeres in <i>Caenorhabditis elegans</i> . Methods, 2012, 56, 50-54.	1.9	6
61	Regulation of UNC-130/FOXD-mediated mesodermal patterning in <i>C. elegans</i> . Developmental Biology, 2016, 416, 300-311.	0.9	6
62	Proteomic analysis uncovers a metabolic phenotype in <i>C. elegans</i> after <i>nhr-40</i> reduction of function. Biochemical and Biophysical Research Communications, 2008, 374, 49-54.	1.0	5
63	Diversification of fasting regulated transcription in a cluster of duplicated nuclear hormone receptors in <i>C. elegans</i> . Gene Expression Patterns, 2010, 10, 227-236.	0.3	5
64	<i>C. elegans</i> Genetic Networks Predict Roles for O-GlcNAc Cycling in Key Signaling Pathways. Current Signal Transduction Therapy, 2010, 5, 60-73.	0.3	2
65	Cell Fate Determination in <i>Caenorhabditis elegans</i> . , 1999, , 251-267.		1