

Jing Kan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5395677/publications.pdf>

Version: 2024-02-01

31
papers

1,828
citations

687363

13
h-index

434195

31
g-index

32
all docs

32
docs citations

32
times ranked

1686
citing authors

#	ARTICLE	IF	CITATIONS
1	Intravascular Ultrasound Versus Angiography-Guided Drug-Eluting Stent Implantation. <i>Journal of the American College of Cardiology</i> , 2018, 72, 3126-3137.	2.8	392
2	Double Kissing Crush Versus Provisional Stenting for Left Main Distal Bifurcation Lesions. <i>Journal of the American College of Cardiology</i> , 2017, 70, 2605-2617.	2.8	256
3	A Randomized Clinical Study Comparing Double Kissing Crush With Provisional Stenting for Treatment of Coronary Bifurcation Lesions. <i>Journal of the American College of Cardiology</i> , 2011, 57, 914-920.	2.8	247
4	3-Year Outcomes of the ULTIMATE Trial Comparing Intravascular Ultrasound Versus Angiography-Guided Drug-Eluting Stent Implantation. <i>JACC: Cardiovascular Interventions</i> , 2021, 14, 247-257.	2.9	149
5	3-Year Outcomes of the DKCRUSH-V Trial Comparing DK Crush With Provisional Stenting for Left Main Bifurcation Lesions. <i>JACC: Cardiovascular Interventions</i> , 2019, 12, 1927-1937.	2.9	130
6	Multicentre, randomized comparison of two-stent and provisional stenting techniques in patients with complex coronary bifurcation lesions: the DEFINITION II trial. <i>European Heart Journal</i> , 2020, 41, 2523-2536.	2.2	124
7	Cutoff Value and Long-Term Prediction of Clinical Events by FFR Measured Immediately After Implantation of a Drug-Eluting Stent in Patients With Coronary Artery Disease. <i>JACC: Cardiovascular Interventions</i> , 2017, 10, 986-995.	2.9	111
8	Clinical Outcome of Double Kissing Crush Versus Provisional Stenting of Coronary Artery Bifurcation Lesions. <i>Circulation: Cardiovascular Interventions</i> , 2017, 10, .	3.9	81
9	Incidence and Clinical Outcomes of Stent Fractures on the Basis of 6,555 Patients and 16,482 Drug-Eluting Stents From 4 Centers. <i>JACC: Cardiovascular Interventions</i> , 2016, 9, 1115-1123.	2.9	66
10	Pulmonary Artery Denervation Significantly Increases 6-Min Walk Distance for Patients With Combined Pre- and Post-Capillary Pulmonary Hypertension Associated With Left Heart Failure. <i>JACC: Cardiovascular Interventions</i> , 2019, 12, 274-284.	2.9	65
11	Comparison of one-year clinical outcomes between intravascular ultrasound-guided versus angiography-guided implantation of drug-eluting stents for left main lesions: a single-center analysis of a 1,016-patient cohort. <i>Patient Preference and Adherence</i> , 2014, 8, 1299.	1.8	43
12	Improved 3-Year Cardiac Survival After IVUS-Guided Long DES Implantation. <i>JACC: Cardiovascular Interventions</i> , 2022, 15, 208-216.	2.9	38
13	Acute renal injury after thoracic endovascular aortic repair of Stanford type B aortic dissection: Incidence, risk factors, and prognosis. <i>Journal of the Formosan Medical Association</i> , 2014, 113, 612-619.	1.7	23
14	Comparison of intravascular ultrasound-guided with angiography-guided double kissing crush stenting for patients with complex coronary bifurcation lesions: Rationale and design of a prospective, randomized, and multicenter DKCRUSH VIII trial. <i>American Heart Journal</i> , 2021, 234, 101-110.	2.7	12
15	Impact of Intravascular Ultrasound-Guided Optimal Stent Expansion on 3-Year Hard Clinical Outcomes. <i>Circulation: Cardiovascular Interventions</i> , 2021, 14, e011124.	3.9	11
16	Is Routine Postdilation During Angiography-Guided Stent Implantation as Good as Intravascular Ultrasound Guidance?: An Analysis Using Data From IVUS-XPL and ULTIMATE. <i>Circulation: Cardiovascular Interventions</i> , 2022, 15, e011366.	3.9	10
17	3-Year Outcomes After 2-Stent With Provisional Stenting for Complex Bifurcation Lesions Defined by DEFINITION Criteria. <i>JACC: Cardiovascular Interventions</i> , 2022, 15, 1310-1320.	2.9	9
18	Obstructive Sleep Apnea Affecting Platelet Reactivity in Patients Undergoing Percutaneous Coronary Intervention. <i>Chinese Medical Journal</i> , 2018, 131, 1023-1029.	2.3	7

#	ARTICLE	IF	CITATIONS
19	Comparison in Prevalence, Predictors, and Clinical Outcome of VSR Versus FWR after Acute Myocardial Infarction: The Prospective, Multicenter Registry MOODY Trial-Heart Rupture Analysis. <i>Cardiovascular Revascularization Medicine</i> , 2019, 20, 1158-1164.	0.8	6
20	Analysis of Serum MicroRNAs as Potential Biomarker in Coronary Bifurcation Lesion. <i>Disease Markers</i> , 2015, 2015, 1-5.	1.3	5
21	Stent fracture is associated with a higher mortality in patients with type-2 diabetes treated by implantation of a second-generation drug-eluting stent. <i>International Journal of Cardiovascular Imaging</i> , 2017, 33, 1873-1881.	1.5	5
22	Optical coherence tomography predictors of target vessel myocardial infarction after provisional stenting in patients with coronary bifurcation disease. <i>Catheterization and Cardiovascular Interventions</i> , 2021, 97, 1331-1340.	1.7	5
23	Long-term mortality after pulmonary artery denervation stratified by baseline functional class in patients with pulmonary arterial hypertension. <i>Asian Intervention</i> , 2022, 8, 58-68.	0.4	5
24	A review in enormity of OCT and its enduring understanding of vulnerable plaque in coronary bifurcation lesion. <i>International Journal of Cardiovascular Imaging</i> , 2018, 34, 1679-1684.	1.5	4
25	Improvement of clinical outcome in patients with ST-elevation myocardial infarction between 1999 and 2016 in China: The Prospective, Multicentre Registry MOODY study. <i>European Journal of Clinical Investigation</i> , 2020, 50, e13197.	3.4	4
26	Association of periprocedural myocardial infarction with mortality after stenting true coronary bifurcation lesions: A pooled individual participant data analysis from four randomized controlled trials. <i>Catheterization and Cardiovascular Interventions</i> , 2021, , .	1.7	4
27	Antiviral Abidol is Associated with the Reduction of In-Hospital Mortality in COVID-19 Patients. <i>Cardiology Discovery</i> , 2021, 1, 37-43.	0.5	4
28	Comparison of two and three dimensional quantitative coronary angiography to intravascular ultrasound in the assessment of left main coronary artery bifurcation lesions. <i>Chinese Medical Journal</i> , 2014, 127, 1012-21.	2.3	4
29	Rationale and design for comparison of non-compliant balloon with drug-coating balloon angioplasty for side branch after provisional stenting for patients with true coronary bifurcation lesions: a prospective, multicentre and randomised DCB-BIF trial. <i>BMJ Open</i> , 2022, 12, e052788.	1.9	4
30	Overlapping Drug-Eluting Stent Is Associated with Increased Definite Stent Thrombosis and Revascularization: Results from 15,561 Patients in the AUTHENTIC Study. <i>Cardiovascular Drugs and Therapy</i> , 2021, 35, 331-341.	2.6	3
31	Quantitative assessment of late lumen loss after biodegradable polymer and permanent polymer sirolimus-eluting stents implantation. <i>Chinese Medical Journal</i> , 2013, 126, 1081-5.	2.3	0