

Albert Escuer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5398771/publications.pdf>

Version: 2024-02-01

305
papers

11,574
citations

28736

57
h-index

49824

91
g-index

322
all docs

322
docs citations

322
times ranked

4659
citing authors

#	ARTICLE	IF	CITATIONS
1	Quasi-isotropic SMMs: slow relaxation of the magnetization in polynuclear CuII/MnII complexes. Dalton Transactions, 2022, 51, 1779-1783.	1.6	9
2	Metalloâ€Ligand Based 3d/4f Coordination Polymers: Synthesis, Structure and Magnetic Properties. European Journal of Inorganic Chemistry, 2022, 2022, .	1.0	3
3	An Unprecedented Dodecanuclear Copper(II) Complex Derived from an Unsymmetrical Schiffâ€Base Ligand. ChemistrySelect, 2022, 7, .	0.7	2
4	Slow magnetic relaxation for cobalt(ⁱⁱ) complexes in axial bipyramidal environment: an $S = 1/2$ spin case. Dalton Transactions, 2022, .	1.6	2
5	Further synthetic investigation of the general lanthanoid(ⁱⁱⁱ) [Ln(ⁱⁱⁱ)]/copper(ⁱⁱ)/pyridine-2,6-dimethanol/carboxylate reaction system: {CuII5LnIII4} coordination clusters (Ln = Dy, Tb, Ho) and their yttrium(ⁱⁱⁱ) analogue. Dalton Transactions, 2021, 50, 240-251.	1.6	4
6	Oxalamide based coordination polymers. Journal of Coordination Chemistry, 2021, 74, 252-265.	0.8	3
7	Synthesis and characterization of new coordination compounds by the use of 2-pyridinemethanol and di- or tricarboxylic acids. CrystEngComm, 2021, 23, 5489-5497.	1.3	3
8	Correlating the axial Zero Field Splitting with the slow magnetic relaxation in Gd ^{III} SIMs. Chemical Communications, 2021, 57, 721-724.	2.2	20
9	Structural and magnetic studies of mononuclear lanthanide complexes derived from N-rich chiral Schiff bases. Dalton Transactions, 2021, 50, 1746-1753.	1.6	6
10	Single-Ion Anisotropy and Intramolecular Interactions in Ce ^{III} and Nd ^{III} Dimers. Inorganic Chemistry, 2021, 60, 8692-8703.	1.9	7
11	New classes of organic Chelate-Free coordination Polymers: An End-On Azido-Bridged Cu(II) 1-D chain composed of {Cu6(N3)12} repeating units. Polyhedron, 2021, 206, 115315.	1.0	4
12	New Highly Charged Iron(III) Metalâ€Organic Cube Stabilized by a Bulky Amine. ACS Omega, 2020, 5, 22238-22247.	1.6	2
13	A biocompatible ZnNa2-based metalâ€organic framework with high ibuprofen, nitric oxide and metal uptake capacity. Materials Advances, 2020, 1, 2248-2260.	2.6	8
14	Chiral Oxazolidine Complexes Derived from Phenolic Schiff Bases. Crystal Growth and Design, 2020, 20, 4176-4184.	1.4	5
15	Chiral Versus Nonâ€Chiral [Mn ^{III} 6Mn ^{II} Na ^I], [Mn ^{III} 6Mn ^{II} Na ^I 2] and [Mn ^{III} 3Mn ^{II} Na ^I 2] Clusters Derived from Schiff Bases or the Fight for Symmetry. Chemistry - A European Journal, 2020, 26, 13053-13062.	1.7	2
16	Rare Nuclearities in Ni(II) Cluster Chemistry: An Unprecedented {Ni12} Nanosized Cage from the Use of N-Naphthalidene-2-Amino-5-Chlorobenzoic Acid. Inorganics, 2020, 8, 32.	1.2	0
17	Na2MnIII6Ln ^{III} clusters with a non-equivalent core: chiral <i>vs</i> <i>i</i> . <i>meso</i> isomerism. Dalton Transactions, 2020, 49, 4216-4219.	1.6	2
18	â€Metal Complexes as Ligandsâ€™ for the Synthesis of Coordination Polymers: A MnIII Monomer as a Building Block for the Preparation of an Unprecedented 1-D {MnIIIMnIII}n Linear Chain. Materials, 2020, 13, 1352.	1.3	2

#	ARTICLE	IF	CITATIONS
19	From Bowls to Capsules: Assembly of Hexanuclear Ni II Rings Tailored by Alkali Cations. <i>Chemistry - A European Journal</i> , 2020, 26, 11158-11169.	1.7	0
20	Multifunctionality in Two Families of Dinuclear Lanthanide(III) Complexes with a Tridentate Schiff-Base Ligand. <i>Inorganic Chemistry</i> , 2019, 58, 9581-9585.	1.9	12
21	Decanuclear FeIII clusters with hemiacetal ligands: a new {M10(μ ₃ -O) ₈ } cluster core. <i>Dalton Transactions</i> , 2019, 48, 13139-13142.	1.6	0
22	Click chemistry as a route to the synthesis of structurally new and magnetically interesting coordination clusters: a {NiII ₈ } complex with a trapezoidal prismatic topology. <i>Dalton Transactions</i> , 2019, 48, 11632-11636.	1.6	4
23	Chiral mononuclear lanthanide complexes derived from chiral Schiff bases: Structural and magnetic studies. <i>Polyhedron</i> , 2019, 170, 264-270.	1.0	12
24	{Ni ₄ } Cubanes from enantiomerically pure 2-(1-hydroxyethyl)pyridine ligands: supramolecular chirality. <i>Dalton Transactions</i> , 2019, 48, 10427-10434.	1.6	5
25	Editorial: Molecular Magnets. <i>Frontiers in Chemistry</i> , 2019, 7, 229.	1.8	9
26	Lithium cations in a self-assembled electrostatic nanocapsule. <i>Dalton Transactions</i> , 2019, 48, 16158-16161.	1.6	6
27	Enhancement of magnetic relaxation properties with 3d diamagnetic cations in [Zn ^{II} Ln ^{III}] and [Ni ^{II} Ln ^{III}], Ln ^{III} = Kramers lanthanides. <i>Dalton Transactions</i> , 2019, 48, 641-652.	1.6	35
28	From Mesocates to Helicates: Structural, Magnetic and Chiro-Optical Studies on Nickel(II) Supramolecular Assemblies Derived from Tetradentate Schiff Bases. <i>Chemistry - A European Journal</i> , 2018, 24, 7653-7663.	1.7	13
29	Cu ^{II} Frameworks from Di-2-pyridyl Ketone and Benzene-1,3,5-triphosphonic Acid. <i>European Journal of Inorganic Chemistry</i> , 2018, 2018, 91-98.	1.0	8
30	Polynuclear ampyrone based 3d coordination clusters. <i>CrystEngComm</i> , 2018, 20, 1411-1421.	1.3	4
31	Triple Halide Bridges in Chiral Mn ^{II} ₂ Mn ^{III} ₆ Na ^I ₂ Cages: Structural and Magnetic Characterization. <i>Inorganic Chemistry</i> , 2018, 57, 926-929.	1.9	9
32	Mononuclear Lanthanide(III)-Salicylideneaniline Complexes: Synthetic, Structural, Spectroscopic, and Magnetic Studies. <i>Magnetochemistry</i> , 2018, 4, 45.	1.0	12
33	Oximate-Based Ligands in 3d ⁴ -Metal Cluster Chemistry: A Family of {Cu ₃ Ln} Complexes with a "Propeller"-like Topology and Single-Molecule Magnetic Behavior. <i>Inorganic Chemistry</i> , 2018, 57, 13944-13952.	1.9	22
34	Chiral [Mn ^{II} Mn ^{III}] ₃ (M ² = Na ^I , Ca ^{II} ,) Tj ETQq0 0 0 rgBT /Overloc Clusters Built from an Enantiomerically Pure Schiff Base: Synthetic, Chiroptical, and Magnetic Properties. <i>Chemistry - A European Journal</i> , 2018, 24, 18705-18717.	1.7	10
35	Chiroptical and magnetic properties of star-shaped FeIII ₄ complexes from chiral Schiff bases. Structural and magnetic correlations based on continuous shape measures. <i>Dalton Transactions</i> , 2018, 47, 8392-8401.	1.6	17
36	Slow magnetic relaxation and luminescence properties in lanthanide(III)/anil complexes. <i>Dalton Transactions</i> , 2018, 47, 11859-11872.	1.6	15

#	ARTICLE	IF	CITATIONS
37	Chiral tetranuclear Ni II clusters derived from Schiff bases and azido co-ligands. <i>Polyhedron</i> , 2018, 150, 10-14.	1.0	8
38	Trinuclear Complexes Derived from R/S Schiff Bases - Chiral Single-Molecule Magnets. <i>European Journal of Inorganic Chemistry</i> , 2017, 2017, 990-990.	1.0	0
39	Four New Families of Polynuclear Zn-Ln Coordination Clusters. Synthetic, Topological, Magnetic, and Luminescent Aspects. <i>Crystal Growth and Design</i> , 2017, 17, 1524-1538.	1.4	30
40	Binding of ligands containing carbonyl and phenol groups to iron(III): new Fe_6 , Fe_{10} and Fe_{12} coordination clusters. <i>Dalton Transactions</i> , 2017, 46, 3240-3251.	1.6	17
41	Syntheses, structures, and chiroptical and magnetic properties of chiral clusters built from Schiff bases: a novel $[\text{Mn}^{\text{II}}\text{Mn}^{\text{III}}_6\text{Na}_2]$ core. <i>Dalton Transactions</i> , 2017, 46, 6514-6517.	1.6	17
42	Trinuclear Complexes Derived from R/S Schiff Bases - Chiral Single-Molecule Magnets. <i>European Journal of Inorganic Chemistry</i> , 2017, 2017, 991-998.	1.0	11
43	Transition Metal Single-Molecule Magnets: A Mn_{31} Nanosized Cluster with a Large Energy Barrier of ~ 460 K and Magnetic Hysteresis at ~ 45 K. <i>Journal of the American Chemical Society</i> , 2017, 139, 15644-15647.	6.6	66
44	Heptanuclear Disk-Like $\text{M}^{\text{II}}_3\text{Ln}^{\text{III}}_4$ ($\text{M} = \text{Ni}, \text{Co}$) Coordination Clusters: Synthesis, Structures and Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2017, 2017, 3938-3945.	1.0	8
45	Structural Diversities in Heterometallic $\text{Mn}^{\text{II}}\text{Ca}$ Cluster Chemistry from the Use of Salicylhydroxamic Acid: $\{\text{Mn}^{\text{III}}_4\text{Ca}_2\}$, $\{\text{Mn}^{\text{II/III}}_6\text{Ca}_2\}$, $\{\text{Mn}^{\text{III/IV}}_8\text{Ca}\}$, and $\{\text{Mn}^{\text{III}}_8\text{Ca}_2\}$ Complexes with Relevance to Both High- and Low-Valent States of the Oxygen-Evolving Complex. <i>Inorganic Chemistry</i> , 2017, 56, 10766-10774.	1.9	15
46	2-hydroxybenzophenone-controlled self-assembly of enneanuclear lanthanide(III) hydroxo coordination clusters with an "hourglass"-like topology. <i>Inorganic Chemistry Communication</i> , 2017, 83, 118-122.	1.8	8
47	Carbonato- and methanediolato(-2)-bridged nickel(II) coordination clusters from the use of N-salicylidene-4-methyl-o-aminophenol. <i>Inorganic Chemistry Communication</i> , 2017, 83, 113-117.	1.8	4
48	Cu(II) frameworks from a "mixed-ligand" approach. <i>CrystEngComm</i> , 2017, 19, 4355-4367.	1.3	7
49	Using the Singly Deprotonated Triethanolamine to Prepare Dinuclear Lanthanide(III) Complexes: Synthesis, Structural Characterization and Magnetic Studies. <i>Magnetochemistry</i> , 2017, 3, 5.	1.0	16
50	A Ni(II) Coordination Cluster from the Use of the Di-2-Pyridyl Ketone/Acetate Ligand Combination: Synthetic, Structural and Magnetic Studies. <i>Magnetochemistry</i> , 2016, 2, 30.	1.0	6
51	Manganese "Salicyloximate Clusters Starting from $[\text{Mn}^{\text{II}}(\text{hfacac})_2]$: From Mn_4 to Mn_{12} . <i>European Journal of Inorganic Chemistry</i> , 2016, 2016, 1232-1241.	1.0	7
52	Structural diversity in Ni^{II} cluster chemistry: Ni_5 , Ni_6 , and $\{\text{NiNa}_2\}_n$ complexes bearing the Schiff-base ligand N-naphthalidene-2-amino-5-chlorobenzoic acid. <i>Dalton Transactions</i> , 2016, 45, 10256-10270.	1.6	15
53	Linked Nickel Metallacrowns from a Phosphonate/2-Pyridyloximate Blend of Ligands: Structure and Magnetic Properties. <i>Inorganic Chemistry</i> , 2016, 55, 3161-3168.	1.9	10
54	Dinuclear cobalt(III) and mixed valence trinuclear $\text{Mn}^{\text{III}}\text{Mn}^{\text{II}}\text{Mn}^{\text{III}}$ complexes with a tripodal bridging pyridylaminophenol ligand. <i>New Journal of Chemistry</i> , 2016, 40, 6164-6170.	1.4	10

#	ARTICLE	IF	CITATIONS
55	Bis(di-2-pyridyl ketoximate- O , N , N $\hat{=}$)bis(di-2-pyridyl ketoxime- N , N $\hat{=}$)dicopper(II) diperchlorate: A plausible, weakly ferromagnetically-coupled intermediate in the formation of the neutral, strongly antiferromagnetically-coupled neutral dimer bearing only deprotonated ligands. <i>Inorganic Chemistry Communication</i> , 2016, 70, 95-98.	1.8	4
56	Topological insights in polynuclear Ni/Na coordination clusters derived from a schiff base ligand. <i>Structural Chemistry</i> , 2016, 27, 1703-1714.	1.0	6
57	A discrete Cu_6 cluster and a 3D $Mn^{II}Cu^{II}$ framework based on assembly of Mn_2Cu_4 clusters: synthesis, structure and magnetic properties. <i>Dalton Transactions</i> , 2016, 45, 15523-15531.	1.6	9
58	New structural motifs in Mn cluster chemistry from the ketone/gem-diol and bis(gem-diol) forms of 2,6-di-(2-pyridylcarbonyl)pyridine: $\{Mn^{II}_4Mn^{III}_2\}$ and $\{Mn^{II}_4Mn^{III}_6\}$ complexes. <i>RSC Advances</i> , 2016, 6, 105969-105979.	1.7	6
59	A Ferromagnetic Salicylaldoximate/Azide $Mn^{II}_2Mn^{III}_6$ Cluster with an $S = 17$ Ground State and a Single-Molecule-Magnet Response. <i>Inorganic Chemistry</i> , 2016, 55, 5735-5737.	1.9	11
60	α -Ligands-with-Benefits: Naphthalene-Substituted Schiff Bases Yielding New Ni^{II} Metal Clusters with Ferromagnetic and Emissive Properties and Undergoing Exciting Transformations. <i>Inorganic Chemistry</i> , 2016, 55, 1270-1277.	1.9	20
61	Interesting copper(II)-assisted transformations of 2-acetylpyridine and 2-benzoylpyridine. <i>Dalton Transactions</i> , 2016, 45, 1063-1077.	1.6	23
62	Copper(II) cubanes with a $\{Cu_4O\}$ core and well defined $S = 1$ ground state. <i>Dalton Transactions</i> , 2016, 45, 1604-1613.	1.6	5
63	Synthesis, Crystal Structures, and Magnetic and Catalytic Studies on a Linear Trinuclear Mn^{II}_3 Complex. <i>ChemPlusChem</i> , 2015, 80, 1440-1447.	1.3	3
64	Ni^{II}_{20} α -Bowls from the Use of Tridentate Schiff Bases. <i>Inorganic Chemistry</i> , 2015, 54, 5615-5617.	1.9	25
65	A new family of high nuclearity Co/Dy^{III} coordination clusters possessing robust and unseen topologies. <i>Dalton Transactions</i> , 2015, 44, 12788-12795.	1.6	12
66	New structural topologies in 4f-metal cluster chemistry from vertex-sharing butterfly units: $\{Ln^{III}_7\}$ complexes exhibiting slow magnetization relaxation and ligand-centred emissions. <i>RSC Advances</i> , 2015, 5, 92534-92538.	1.7	24
67	Synthesis and first use of pyridine-2,6-diylbis(pyrazine-2-ylmethanone) in metal cluster chemistry: a $\{Mn^{III}_3Na_2\}$ complex with an ideal trigonal bipyramidal geometry. <i>Dalton Transactions</i> , 2015, 44, 4318-4327.	1.6	2
68	A Mn_6 Cluster inside a Mn_{10} Wheel: Characterization of a Mn_{16} Oximate Complex Resulting from a Tetrazole-2-pyridylketoneoximate Ligand. <i>European Journal of Inorganic Chemistry</i> , 2015, 2015, 1326-1329.	1.0	6
69	Cobalt(II), nickel(II) and copper(II) coordination clusters employing a monoanionic Schiff base ligand: synthetic, topological and computational mechanistic aspects. <i>CrystEngComm</i> , 2015, 17, 6753-6764.	1.3	20
70	Mn^{II} - and Co^{II} -Catalyzed Transformation of 2-Cyanopyrimidine to Methylimidate by Sodium Azide: Isolation, Structural Characterization, and Magnetic Studies on 2D Mn^{II} - and Cu^{II} -Complexes. <i>Inorganic Chemistry</i> , 2015, 54, 7030-7037.	1.9	9
71	Nonemployed Simple Carboxylate Ions in Well-Investigated Areas of Heterometallic Carboxylate Cluster Chemistry: A New Family of $\{Cu^{II}_4Ln^{III}_8\}$ Complexes Bearing <i>tert</i> -Butylacetate Bridging Ligands. <i>Inorganic Chemistry</i> , 2015, 54, 7555-7561.	1.9	24
72	A family of dinuclear lanthanide(III) complexes from the use of a tridentate Schiff base. <i>Dalton Transactions</i> , 2015, 44, 10200-10209.	1.6	60

#	ARTICLE	IF	CITATIONS
73	A Disk-Like Co ^{II} ₃ Dy ^{III} ₄ Coordination Cluster Exhibiting Single Molecule Magnet Behavior. <i>European Journal of Inorganic Chemistry</i> , 2015, 2015, 2646-2649.	1.0	17
74	Spin Frustration in Triangular Cu ₃ Complexes with 6-Methyl-2-pyridyloxime as Ligand – Synthesis, Structural, and Magnetic Characterization. <i>European Journal of Inorganic Chemistry</i> , 2014, 2014, 83-92.	1.0	21
75	Structural aesthetics in molecular nanoscience: a unique Ni ₂₆ cluster with a “rabbit-face” topology and a discrete Ni ₁₈ – molecular chain™. <i>Chemical Communications</i> , 2014, 50, 14942-14945.	2.2	36
76	Template arrangement of {Ni ₃ (¹ / ₄ -OH)(SO ₄)(R-NO) ₂ } ⁺ fragments around Na ⁺ cations: An unprecedented {Ni ₁₂ Na} unit linked by oximate and sulfate bridges. <i>Inorganic Chemistry Communication</i> , 2014, 43, 169-172.	1.8	2
77	The bridging azido ligand as a central “player” in high-nuclearity 3d-metal cluster chemistry. <i>Coordination Chemistry Reviews</i> , 2014, 275, 87-129.	9.5	158
78	2-Pyridylcyanoxime-NiII Clusters with Unusual Topologies: Lone-Pair- π Interactions and Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2014, 2014, 5443-5450.	1.0	6
79	Rare nuclearities in Ni(λ -N-salicylidene-2-amino-5-chlorobenzoic acid) cluster chemistry: a Ni ₁₁ cage from the first use of N-salicylidene-2-amino-5-chlorobenzoic acid in metal cluster chemistry. <i>RSC Advances</i> , 2014, 4, 12680-12684.	1.7	10
80	Unexpected metal ion-assisted transformations leading to unexplored bridging ligands in Ni ^{II} coordination chemistry: the case of PO ₃ F ₂ ³⁻ group. <i>Dalton Transactions</i> , 2014, 43, 14520-14524.	1.6	11
81	Anionic Guests in Prismatic Cavities Generated by Enneanuclear Nickel Metallacycles. <i>Inorganic Chemistry</i> , 2014, 53, 1113-1121.	1.9	16
82	New Classes of Ferromagnetic Materials with Exclusively End-on Azido Bridges: From Single-Molecule Magnets to 2D Molecular-Based Magnets. <i>Chemistry - A European Journal</i> , 2014, 20, 13860-13864.	1.7	25
83	Structural and magnetic variations in tetranuclear Ni ^{II} clusters: the effect of the reaction solvent and ligand substitution on product identity. <i>Dalton Transactions</i> , 2014, 43, 16605-16609.	1.6	32
84	New Topologies in Pentanuclear Nickel/Oximate Clusters: Structural and Magnetic Characterization. <i>Inorganic Chemistry</i> , 2014, 53, 3194-3203.	1.9	21
85	The templating effect of halides in the tetrameric copper(II) [Cu ₂ (LH) ₂ (¹ / ₄ -X)Cu ₂ (LH) ₂] ₃₊ complexes (LH ₂ =N-(2-pyridylmethyl)-N,N-bis-[2-hydroxy-5-methyl-benzyl]-amine; X=Br, Cl). <i>Synthesis and magneto-structural characterization</i> . <i>Polyhedron</i> , 2014, 76, 117-121.	1.0	3
86	Manganese clusters derived from 2-pyridylcyanoxime: new topologies and a large spin ground state in pyridyloximate chemistry. <i>Dalton Transactions</i> , 2013, 42, 12334.	1.6	15
87	Ni ^{II} -Pyridyloximate Triangles with a Central ¹ / ₄ -OH Ligand: Magnetostructural Correlations. <i>European Journal of Inorganic Chemistry</i> , 2013, 2013, 5274-5280.	1.0	7
88	Molecular Nanoscale Magnetic Refrigerants: A Ferrimagnetic {Cu ^{II} ₁₅ Gd ^{III} ₇ } Cagelike Cluster from the Use of Pyridine-2,6-dimethanol. <i>Inorganic Chemistry</i> , 2013, 52, 10235-10237.	1.9	58
89	Ligand-to-metal ratio dependent assembly of two distinct 1D and 3D copper(II)-dicyanamide magnetic coordination polymers with a tripyridyltriazole co-tecton. <i>Inorganica Chimica Acta</i> , 2013, 403, 142-146.	1.2	23
90	Slow Magnetization Relaxation in Unprecedented Mn ^{IV} ₄ Dy ^{III} ₃ and Mn ^{IV} ₄ Dy ^{III} ₅ Clusters from the Use of <i>N</i> -Salicylidene- <i>o</i> -aminophenol. <i>Inorganic Chemistry</i> , 2013, 52, 1179-1181.	1.9	41

#	ARTICLE	IF	CITATIONS
91	Polynuclear pyridyldioximato-nickel(II) clusters: Synthesis, structure and magnetic study. <i>Polyhedron</i> , 2013, 52, 339-345.	1.0	5
92	Copper and manganese derivatives of pyrazinyloximato ligands: Synthesis, structure and magnetic study. <i>Polyhedron</i> , 2013, 64, 84-90.	1.0	4
93	Copper(II)/di-2-pyridyl ketone chemistry: A triangular cluster displaying antisymmetric exchange versus an 1D coordination polymer. <i>Polyhedron</i> , 2013, 64, 30-37.	1.0	13
94	In search of 3d/4f-metal single-molecule magnets: Nickel(II)/lanthanide(III) coordination clusters. <i>Pure and Applied Chemistry</i> , 2013, 85, 315-327.	0.9	37
95	Dinuclear Lanthanide(III) Complexes by Metal-Ion-Assisted Hydration of Di-2-pyridyl Ketone Azine. <i>Inorganic Chemistry</i> , 2013, 52, 4145-4147.	1.9	21
96	Azido-bridging cobalt(II) systems: crystal structures and magnetic properties. <i>Acta Crystallographica Section A: Foundations and Advances</i> , 2013, 69, s624-s624.	0.3	2
97	The "periodic table" of di-2-pyridyl ketone: vanadium complexes. <i>Dalton Transactions</i> , 2012, 41, 11984.	1.6	13
98	Anion coordination by metallamacrocycles: a cryptand-like cavity. <i>Chemical Communications</i> , 2012, 48, 9777.	2.2	11
99	Employment of methyl 2-pyridyl ketone oxime in 3d/4f-metal chemistry: dinuclear nickel(ii)/lanthanide(iii) species and complexes containing the metals in separate ions. <i>Dalton Transactions</i> , 2012, 41, 13755.	1.6	34
100	[Mn ₂ (N ₃) ₅] ²⁻ : Four different azide bridging modes and dicubane subunits observed in a new Mn(II)-azide only 2D system. <i>Inorganic Chemistry Communication</i> , 2012, 21, 4-7.	1.8	21
101	High Nuclearity in Azido/Oximate Chemistry: Ni ₁₄ and Ni ₁₃ Clusters with $\langle i \rangle S \langle /i \rangle = 6$ and 9 Ground States. <i>Inorganic Chemistry</i> , 2012, 51, 5503-5505.	1.9	34
102	Single-Strand Molecular Wheels and Coordination Polymers in Copper(II) Benzoate Chemistry by the Employment of \pm -Benzoin Oxime and Azides: Synthesis, Structures, and Magnetic Characterization. <i>European Journal of Inorganic Chemistry</i> , 2012, 2012, 3121-3131.	1.0	27
103	Triangular Nickel Complexes Derived from 2-Pyridylcyanoxime: An Approach to the Magnetic Properties of the [Ni ₃ (μ_3 -OH){pyC(R)NO} ₃] ²⁺ Core. <i>Chemistry - A European Journal</i> , 2012, 18, 3637-3648.	1.7	35
104	Synthesis and characterization of CoIII inverse metallacrowns via use of 6-methyl-2-pyridylalldoxime. <i>Inorganic Chemistry Communication</i> , 2012, 16, 78-80.	1.8	9
105	A novel ferromagnetically-coupled trinuclear nickel(II) complex constructed from the new 1,2-di(pyridin-2-yl)ethanone ligand in its enolate form. <i>Inorganic Chemistry Communication</i> , 2012, 20, 184-187.	1.8	3
106	Dinuclear, tetranuclear and polymeric complexes in copper(ii) perchlorate/pyridine-2,6-diamidoxime chemistry: synthetic, structural and magnetic studies. <i>Dalton Transactions</i> , 2011, 40, 225-233.	1.6	23
107	Use of 6-methylpyridine-2-carbaldehydeoxime in nickel(ii) carboxylate chemistry: synthetic, structural and magnetic properties of penta and hexanuclear complexes. <i>Dalton Transactions</i> , 2011, 40, 10109.	1.6	21
108	Building of a novel Mn ₁₂ Single Molecule Magnet by assembly of anisotropic {Mn ₃ (μ_3 -O)(salox) ₃ } ⁺ triangles. <i>Dalton Transactions</i> , 2011, 40, 7127.	1.6	20

#	ARTICLE	IF	CITATIONS
109	Ni ₅ , Ni ₈ , and Ni ₁₀ Clusters with 2,6-Diacetylpyridine-dioxime as a Ligand. <i>Inorganic Chemistry</i> , 2011, 50, 8893-8901.	1.9	25
110	Trinuclear, Tetranuclear, and Polymeric Cu ^{II} Complexes from the First Use of 2-Pyridylcyanoxime in Transition Metal Chemistry: Synthetic, Structural, and Magnetic Studies. <i>Inorganic Chemistry</i> , 2011, 50, 2468-2478.	1.9	57
111	Assembly of [MnII ₂ MnIII ₂]S=9 Clusters via Azido Bridges: a New Single-Chain Magnet. <i>Inorganic Chemistry</i> , 2011, 50, 2717-2719.	1.9	44
112	Triangular NiII ₂ LnIII and NiII ₂ YIII complexes derived from di-2-pyridyl ketone: Synthesis, structures and magnetic properties. <i>Polyhedron</i> , 2011, 30, 2978-2986.	1.0	25
113	Cyanato bridged binuclear nickel(II) and copper(II) complexes with pyridylpyrazole ligand: Synthesis, structure and magnetic properties. <i>Inorganica Chimica Acta</i> , 2011, 375, 333-337.	1.2	29
114	Two new azido bridging Mn(II) 1D systems: Synthesis and characterization of trans-[Mn(N ₃) ₂ (2-aminopyridine) ₂] _n and trans-[Mn(N ₃) ₂ (4-azidopyridine) ₂] _n . <i>Journal of Molecular Structure</i> , 2010, 969, 192-196.	1.8	19
115	Halo and azido copper(II) coordination polymers featuring the gem-diolate forms of di-2-pyridyl ketone. <i>Polyhedron</i> , 2010, 29, 100-109.	1.0	15
116	Strong antiferromagnetic coupling in doubly N,O oximato-bridged dinuclear copper(II) complexes. <i>Polyhedron</i> , 2010, 29, 204-211.	1.0	31
117	Triangular Ni(II) complexes from the use of 2-pyridyl oximes. <i>Polyhedron</i> , 2010, 29, 627-633.	1.0	16
118	Manganese clusters derived from a 2,6-diacetylpyridine dioximato ligand: structure and magnetic study. <i>Dalton Transactions</i> , 2010, 39, 4817.	1.6	14
119	Anionic Tuning of the Dimensionality in Copper Oximato Chemistry. <i>Inorganic Chemistry</i> , 2010, 49, 9752-9754.	1.9	38
120	Depolymerization Approach in Mn Cluster Chemistry: Controlled Cleavage of a 1D Coordination Polymer Consisting of Mn ₈ Units in Its Constituent, Discrete Mn ₈ Complex. <i>Inorganic Chemistry</i> , 2010, 49, 359-361.	1.9	20
121	First Structural and Magnetic Studies of Ni Clusters Containing 2,6-Diacetylpyridine-dioxime as a Ligand. <i>Inorganic Chemistry</i> , 2010, 49, 2259-2266.	1.9	22
122	Combining Azide, Carboxylate, and 2-Pyridyloximate Ligands in Transition-Metal Chemistry: Ferromagnetic NiII ₅ Clusters with a Bowtie Skeleton. <i>Inorganic Chemistry</i> , 2010, 49, 10486-10496.	1.9	76
123	Topologic ferrimagnetism in cobalt(ii)-azido chains. <i>Dalton Transactions</i> , 2010, 39, 4482.	1.6	18
124	Initial employment of di-2-pyridyl ketone as a route to nickel(ii)/lanthanide(iii) clusters: triangular Ni ₂ Ln complexes. <i>Dalton Transactions</i> , 2010, 39, 8603.	1.6	42
125	The sulfate ligand as a promising player in 3d-metal cluster chemistry. <i>Inorganica Chimica Acta</i> , 2009, 362, 634-650.	1.2	42
126	A three-dimensional copper(II) coordination polymer featuring the 2,3-dioxyquinoxalinate(-2) ligand: Preparation, structural characterization and magnetic study. <i>Polyhedron</i> , 2009, 28, 1646-1651.	1.0	8

#	ARTICLE	IF	CITATIONS
127	Employment of the sulfate ligand in 3d-metal cluster chemistry: A novel hexanuclear nickel(II) complex with a chair metal topology. <i>Polyhedron</i> , 2009, 28, 3177-3184.	1.0	25
128	A tetranuclear complex from the employment of pyridine-2,6-dimethanol in copper(II) nitrate chemistry: Synthetic, structural and magnetic studies. <i>Polyhedron</i> , 2009, 28, 3235-3242.	1.0	22
129	Interpretation of the Magnetic Properties of a Compound Consisting of Cocrystallized Cu_{13} and Cu_{14} Clusters through the Targeted Synthesis and Study of Its Discrete Cu_{14} Component. <i>Inorganic Chemistry</i> , 2009, 48, 4610-4612.	1.9	32
130	Octanuclear manganese(ii,iii) clusters stabilized with diamino-alcoxo ligands. <i>Dalton Transactions</i> , 2009, , 9924.	1.6	3
131	Manganese(II) Trimeric Systems Derived from Pyridyldioximate Ligands: Synthesis, Crystal Structure, and Magnetic Characterization. <i>European Journal of Inorganic Chemistry</i> , 2008, 2008, 5082-5087.	1.0	39
132	First use of 1,4-dihydro-2,3-quinoxalinedione in the chemistry of coordination polymers: A 3D copper(II) complex containing the 2,3-dioxyquinoxalinate(η^2) ligand in a novel coordination mode. <i>Inorganic Chemistry Communication</i> , 2008, 11, 186-191.	1.8	12
133	Two new tetranuclear copper(II) complexes: Structure and magnetic studies. <i>Inorganica Chimica Acta</i> , 2008, 361, 4065-4069.	1.2	8
134	Solvatomorphs of dimeric transition metal complexes based on the V_4O_{12} cyclic anion as building block: Crystalline packing and magnetic properties. <i>Inorganica Chimica Acta</i> , 2008, 361, 3681-3689.	1.2	24
135	Unusual Structural Types in Nickel Cluster Chemistry from the Use of Pyridyl Oximes: Ni_5 , $Ni_{12}Na_2$, and Ni_{14} Clusters. <i>Inorganic Chemistry</i> , 2008, 47, 11825-11838.	1.9	76
136	An unusual 3D-topology and dominant ferromagnetic couplings in two Cu(ii)-azide coordination polymers. <i>Dalton Transactions</i> , 2008, , 3553.	1.6	68
137	Copper(II) Complexes with New Polypodal Ligands Presenting Axial~Equatorial Phenoxo Bridges {2-[(Bis(2-pyridylmethyl)amino)methyl]-4-methylphenol, 2-[(Bis(2-pyridylmethyl)amino)methyl]-4-methyl-6-(methylthio)phenol]: Examples of Ferromagnetically Coupled Bi- and Trinuclear Copper(II) Complexes. <i>Inorganic Chemistry</i> , 2007, 46, 6924-6932.	1.9	44
138	Ferromagnetic Coupling in a 1D Coordination Polymer Containing a Symmetric $[Cu(\frac{1}{4}1,1-N_3)_2Cu(\frac{1}{4}1,1-N_3)_2Cu]^{2+}$ Core and Based on an Organic Ligand Obtained from the Solid State. <i>Inorganic Chemistry</i> , 2007, 46, 8843-8850.	1.9	71
139	Acetate/Di-2-pyridyl Ketone Oximate Blend as a Source of High-Nuclearity Nickel(II) Clusters: Dependence of the Nuclearity on the Nature of the Inorganic Anion Present. <i>Inorganic Chemistry</i> , 2007, 46, 2350-2352.	1.9	65
140	New Ternary Complexes of Copper(II) with 2,2'-Bipyridine (Bpy) and Phosphocholine (PCh) or the Quaternary 1-(2-Phosphonomethoxy)ethyl Derivative of 2,4-Diaminopyrimidine (PMEDAPy). <i>European Journal of Inorganic Chemistry</i> , 2007, 2007, 1867-1873.	1.0	7
141	Use of the Sulfato Ligand in 3d-Metal Cluster Chemistry: A Family of Hexanuclear Nickel(II) Complexes with 2-Pyridyl-Substituted Oxime Ligands. <i>European Journal of Inorganic Chemistry</i> , 2007, 2007, 2761-2774.	1.0	54
142	The benzoate/nitrate/ $2,2' : 6,6' : 2,2'$ -terpyridine blend™ in lanthanide(III) chemistry: Relevance to the separation of lanthanides and actinides by solvent extraction. <i>Inorganic Chemistry Communication</i> , 2007, 10, 800-804.	1.8	13
143	A rare all-Mn ²⁺ decametallc cage from distorted face-sharing cubes. <i>Inorganica Chimica Acta</i> , 2007, 360, 61-68.	1.2	25
144	Initial use of di-2-pyridyl ketone oxime in chromium carboxylate chemistry: Triangular compounds and unexpected formation of a carboxylate-free dichromium(II,II) complex. <i>Inorganica Chimica Acta</i> , 2007, 360, 69-83.	1.2	26

#	ARTICLE	IF	CITATIONS
145	Initial use of the di-2-pyridyl ketone/sulfate blend in 3d-metal cluster chemistry: Preparation, X-ray structures and physical studies of zinc(II) and nickel(II) cubanes. <i>Journal of Molecular Structure</i> , 2007, 829, 176-188.	1.8	41
146	A general synthetic route for the preparation of high-spin molecules: Replacement of bridging hydroxo ligands in molecular clusters by end-on azido ligands. <i>Polyhedron</i> , 2007, 26, 2089-2094.	1.0	25
147	Synthesis, structure and magnetic properties of new 1D and 2D manganese azido complexes. <i>Acta Crystallographica Section A: Foundations and Advances</i> , 2007, 63, s153-s153.	0.3	0
148	Complementarity and countercomplementarity in polynuclear copper(ii) complexes with R ₂ NCH ₂ CH(OH)CH ₂ NR ₂ (R = H, CH ₃): crystal structures and magnetic study. <i>Dalton Transactions</i> , 2006, , 2934.	1.6	36
149	A new 2D network built from potassium sandwiches {K[CuII ₃ (bdap) ₃] ₂ } and { $\frac{1}{4}$ [1,3-SCN) ₃ CuI(NCS)]} anions: structure and magnetic behaviour. <i>Chemical Communications</i> , 2006, , 3113-3115.	2.2	19
150	Five New Manganese ^{1/4} -Azido 2D Compounds Synthesized in Aqueous Hydrazoic Acid from Pyrazine Derivatives. <i>Inorganic Chemistry</i> , 2006, 45, 868-876.	1.9	46
151	A rare mixed-valence state manganese(II/IV) tetranuclear cage formed using phenyl 2-pyridyl ketone oxime and azide as ligands. <i>Inorganic Chemistry Communication</i> , 2006, 9, 638-641.	1.8	39
152	An unusual dichromium(II,II) compound bearing di-2-pyridyl ketone oximate ligands and prepared by the ligand-assisted reduction of a trichromium(III,II,III) complex in air. <i>Inorganic Chemistry Communication</i> , 2006, 9, 1178-1182.	1.8	13
153	Enneanuclear Ni(II) complexes from the use of the flexible ligand 2-pyridinealdoxime: The nature of the inorganic anion does not affect the chemical and structural identity of the cationic cluster. <i>Inorganica Chimica Acta</i> , 2006, 359, 4149-4157.	1.2	36
154	Synthesis, Structure, and Magnetic Behavior of Two New 1D Polymeric Manganese Azido Complexes. <i>European Journal of Inorganic Chemistry</i> , 2006, 2006, 3177-3184.	1.0	29
155	Azide as a Bridging Ligand and Magnetic Coupler in Transition Metal Clusters. <i>European Journal of Inorganic Chemistry</i> , 2006, 2006, 4721-4736.	1.0	330
156	New homogeneous and alternating Mn(II)-azido 1D systems. <i>Polyhedron</i> , 2005, 24, 557-562.	1.0	23
157	Use of the Di-2-pyridyl Ketone/Acetate/Dicyanamide Blend in Manganese(II), Cobalt(II) and Nickel(II) Chemistry: Neutral Cubane Complexes. <i>European Journal of Inorganic Chemistry</i> , 2005, 2005, 879-893.	1.0	82
158	New Polymeric Manganese Azide Derivatives with Quinazoline. <i>European Journal of Inorganic Chemistry</i> , 2005, 2005, 4659-4664.	1.0	23
159	Ferromagnetic Interaction in 1,3-Cyanamido-Derived Copper(II) Cryptates. <i>Chemistry - A European Journal</i> , 2005, 11, 398-405.	1.7	17
160	Di-2-pyridyl ketone oxime [(py) ₂ CNOH] in manganese carboxylate chemistry: mononuclear, dinuclear and tetranuclear complexes, and partial transformation of (py) ₂ CNOH to the gem-diolate(2 ⁻) derivative of di-2-pyridyl ketone leading to the formation of NO ₃ ⁻ . <i>Dalton Transactions</i> , 2005, , 501-511.	1.6	71
161	Topological ferrimagnetic behavior of one new chain with the new AF/F ² /F ² /F alternating sequence. <i>New Journal of Chemistry</i> , 2005, 29, 306-314.	1.4	40
162	A two-dimensional azido-based topologic ferrimagnet. <i>Chemical Communications</i> , 2005, , 605.	2.2	100

#	ARTICLE	IF	CITATIONS
163	The First Cobalt Metallacrowns: Preparation and Characterization of Mixed-Valence Cobalt(II/III), Inverse 12-Metallacrown-4 Complexes. <i>Inorganic Chemistry</i> , 2005, 44, 3374-3376.	1.9	77
164	Hexanuclear Manganese(III) Single-Molecule Magnets. <i>Angewandte Chemie - International Edition</i> , 2004, 43, 210-212.	7.2	232
165	Phenylcyanamido Ligand Building Molecular, 1-D, and 2-D Systems Syntheses, Crystal Structures and Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2004, 2004, 309-316.	1.0	10
166	Phenyl 2-Pyridyl Ketone and Its Oxime in Manganese Carboxylate Chemistry: Synthesis, Characterisation, X-ray Studies and Magnetic Properties of Mononuclear, Trinuclear and Octanuclear Complexes. <i>European Journal of Inorganic Chemistry</i> , 2004, 2004, 2885-2901.	1.0	102
167	Three New Dinuclear Copper(II) Complexes with $[Cu(\frac{1}{4}1,3-N3)2Cu]^{2+}$ and $[Cu(\frac{1}{4}1,1-N3)2Cu]^{2+}$ Asymmetrical Cores: Syntheses, Structures and Magnetic Behavior.. <i>ChemInform</i> , 2004, 35, no.	0.1	0
168	2-Pyridinealdoxime [(py)CHNOH] in manganese(II) carboxylate chemistry: mononuclear, dinuclear, tetranuclear and polymeric complexes, and partial transformation of (py)CHNOH to picolinate(λ^1). <i>Polyhedron</i> , 2004, 23, 83-95.	1.0	92
169	New examples of the unusual X-phenylcyanamido bridging ligands. Crystal structure and magnetic properties of MnII derivatives. <i>Polyhedron</i> , 2004, 23, 1409-1417.	1.0	20
170	Polynuclear and 1-D derivatives of 1,3-bis(dimethylamino)-2-propanolato ligand. Structure and magnetic characterization. <i>Dalton Transactions</i> , 2004, , 1867.	1.6	20
171	Three new dinuclear copper(ii) complexes with $[Cu(\frac{1}{4}1,3-N3)2Cu]^{2+}$ and $[Cu(\frac{1}{4}1,1-N3)2Cu]^{2+}$ asymmetrical cores: syntheses, structures and magnetic behaviour. <i>New Journal of Chemistry</i> , 2004, 28, 681-686.	1.4	106
172	Crystal Structure and Magnetic Interactions in Nickel(II) Dibrigded Complexes Formed by Two Azide Groups or by Both Phenolate Oxygen Azide, Thiocyanate, Carboxylate, or Cyanate Groups. <i>Inorganic Chemistry</i> , 2004, 43, 2427-2434.	1.9	181
173	Syntheses and Magneto Structural Study of Several Polynuclear Copper(II) Complexes Derived from 1,3-Bis(dimethylamino)-2-propanolato. <i>Inorganic Chemistry</i> , 2004, 43, 7218-7226.	1.9	29
174	Syntheses, Structure and Magnetic Properties of the First $\mu_1,5$ -Dicyanamido-Bridged Dinuclear Compounds $[Ni(\mu_1,5-dca)(dca)(dpt)]_2$ and $[Ni(\mu_1,5-dca)(dca)(medpt)]_2$. <i>European Journal of Inorganic Chemistry</i> , 2003, 2003, 3929-3933.	1.0	39
175	Synthesis, structural characterisation and magnetic behaviour of two new 1-D trans- $(\frac{1}{4}1,3-N3)$ nickel(II) compounds. Observation of magnetic transition at 288 and 137 K and determination of the Haldane gap for the trans- $\{(\frac{1}{4}1,3-N3)[Ni(meso-cth)]\}_n(ClO_4)_n$ compound. <i>Inorganica Chimica Acta</i> , 2003, 344, 133-142.	1.2	13
176	Di-2-pyridyl ketone oxime in 3d-metal carboxylate cluster chemistry: a new family of mixed-valence MnII/MnIII complexes. <i>Inorganic Chemistry Communication</i> , 2003, 6, 1056-1060.	1.8	38
177	Molecular, 1D, and 2D Systems Built from Phenylcyanamido Ligands. Syntheses, Crystal Structures, and Characterization of Their Magnetic Properties. <i>Inorganic Chemistry</i> , 2003, 42, 541-551.	1.9	21
178	Supramolecular systems built from phenylcyanamido ligands. Syntheses, crystal structures and magnetic properties. <i>Dalton Transactions</i> , 2003, , 2121.	1.6	7
179	Octanuclearity and tetradecanuclearity in manganese chemistry: an octanuclear manganese(ii)/(iii) complex featuring the novel $[Mn_8(\mu_4-O)_2(\mu_3-OH)_2]^{14+}$ core and $[Mn_{10}IIIMn_4IIIIO_4(O_2CMe)_{20}\{(2-py)_2C(OH)O\}_4]$ (2-py = 2-pyridyl). <i>Chemical Communications</i> , 2003, , 819-821.	2.2	97
180	Synthesis, crystal structure and magnetic properties of the triangulo-tricopper(ii) complex $[Cu_3(cpse)_3(H_2O)_3]\cdot 8.5H_2O$. <i>Dalton Transactions RSC</i> , 2002, , 2648-2653.	2.3	43

#	ARTICLE	IF	CITATIONS
181	Letter: Comments on a cyclic tetranuclear copper(II) complex. Dalton Transactions RSC, 2002, , 3778-3778.	2.3	6
182	[M(N3)2(L)]n: building 3-D Millâ€“azido networks with new topologies. Chemical Communications, 2002, , 64-65.	2.2	67
183	Three Novel End-to-End Single Azido-Bridged Ferromagnetic Copper(II) Chains: Synthesis, Crystal Structure, and Magnetic Behavior. European Journal of Inorganic Chemistry, 2002, 2002, 943-949.	1.0	78
184	A One-Dimensional Manganese(II) Coordination Polymer Derived from Zerovalent Manganese and 1-Hydroxybenzotriazole â” Synthesis, Characterization, Crystal Structure and Magnetic Properties. European Journal of Inorganic Chemistry, 2002, 2002, 2488-2493.	1.0	27
185	A new series of 1D polymeric nickel(II) complex cations of pyridine derivative ligands with a single $\frac{1}{4}$ -1,3 azide bridge and X-ray crystal structure and magnetic properties of polymeric $\{[Ni(4\text{-ethylpyridine})_4(N_3)](PF_6)]_n\}$. Polyhedron, 2002, 21, 1871-1876.	1.0	39
186	Benzoate as terminal ligand in the defective double-cubane, tetranuclear cobalt(II) complex $[Co_4(N_3)_2(O_2CPh)_2\{(py)_2C(OH)O\}_4] \cdot 2DMF$ with simultaneous $\frac{1}{4}$ 1,1-azido and $\frac{1}{4}$ -O bridges $[(py)_2C(OH)O=theo$ monoanion of the hydrated, gem-diol form of di-2-pyridyl ketone]. Polyhedron, 2002, 21, 2027-2032.		55
187	Syntheses, structures and magnetic properties of the dicyanamide (dca) polynuclear compounds $[Mn(ac)(terpy)(\frac{1}{4}1,5\text{-dca})]_n$, $[Mn(pdz)_2(\frac{1}{4}1,5\text{-dca})_2]_n$ and $[Mn(dca)(terpy)(MeOH)]_2(\frac{1}{4}\text{-terephthalate})$. Inorganica Chimica Acta, 2002, 340, 163-169.	1.2	58
188	Synthesis, structure and magnetic behaviour of a two-dimensional cyano-bridged complex $[\{Cu(ept)\}_3Fe(CN)_6](ClO_4)_2 \cdot 5H_2O$ [ept = N-(2-aminoethyl)-1,3-diaminopropane]. New Journal of Chemistry, 2001, 25, 655-658.	1.4	17
189	Reactivity in polynuclear transition metal chemistry as a means to obtain high-spin molecules: substitution of $\mu_4\text{-OH}^{\ominus}$ by $\frac{1}{4}\mu_4\text{-N}_3^{\ominus}$ increases nine times the ground-state S value of a nonanuclear nickel(II) cage. Chemical Communications, 2001, , 2414-2415.	2.2	157
190	Synthesis, Crystal Structure and Magnetic Properties of Novel Complex $[\frac{1}{4}\text{-}(NC)\text{-Fe}(CN)_3(NO)\text{-}\frac{1}{4}\text{-}(CN)\text{-Cu}(ept)]_n \cdot 4nH_2O$ (ept = N-(2-Aminoethyl)propane-1,3-diamine). Collection of Czechoslovak Chemical Communications, 2001, 66, 1490-1498.	1.0	4
191	Synthesis, spectral, magnetic and crystal structural characterization of two new copper(II) azido complexes: catena- $[\frac{1}{4}(N_3)Cu(pyridine)_3]_n(PF_6)_n$ and dimeric $[Cu(4\text{-ethylpyridine})(N_3)_2]_2$. Polyhedron, 2001, 20, 2971-2977.	1.0	35
192	Topological Control in Two-Dimensional Cobalt(II) Coordination Polymers by $\pi\text{-}\pi$ Stacking Interactions: Synthesis, Spectroscopic Characterization, Crystal Structure, and Magnetic Properties. Journal of Solid State Chemistry, 2001, 159, 371-378.	1.4	26
193	Defective Double-Cubane, Tetranuclear Manganese(II) and Cobalt(II) Complexes with Simultaneous $\frac{1}{4}$ 1,1-Azido and $\frac{1}{4}$ -O Bridges. European Journal of Inorganic Chemistry, 2001, 2001, 1567-1574.	1.0	105
194	Unique Single-Atom Binding of Pseudohalogeno Ligands to Four Metal Ions Induced by Their Trapping into High-Nuclearity Cages. Angewandte Chemie - International Edition, 2001, 40, 884-886.	7.2	208
195	Synthesis, Structural Characterisation, and Monte Carlo Simulation of the Magnetic Properties of the 3D-Stacked Honeycomb $Csn[\{Mn(N_3)_3\}_n]$ and the Irregular Double Chain $[\{N(C_2H_5)_4\}_n][\{Mn_2(N_3)_5(H_2O)\}_n]$. Chemistry - A European Journal, 2000, 6, 778-784.	1.7	148
196	Can a Homometallic Chain Be Ferrimagnetic?. Angewandte Chemie - International Edition, 2000, 39, 1624-1626.	7.2	134
197	Superexchange Interactions through Quasi-Linear End-to-End Azido Bridges: Structural and Magnetic Characterisation of a New Two-Dimensional Manganeseâ€“Azido System $[Mn(DENA)_2(N_3)_2]_n$ (DENA =) Tj ETQq1 1.0.7843148 BT /Ov	1.0	148
198	Synthesis, structural determination and magnetic properties of three new di- $\frac{1}{4}$ 1,3-azido copper(II) dimers derived from tridentate amines: magneto-structural correlations. Inorganica Chimica Acta, 2000, 298, 195-201.	1.2	52

#	ARTICLE	IF	CITATIONS
199	Topological Ferrimagnetic Behavior of Two New $[Mn(L)_2(N_3)_2]_n$ Chains with the New AF/AF/F Alternating Sequence (L = 3-Methylpyridine or 3,4-Dimethylpyridine). <i>Inorganic Chemistry</i> , 2000, 39, 5022-5027.	1.9	149
200	Synthesis, Structural Characterization, and Monte Carlo Simulation of the Magnetic Properties of Two New Alternating MnII Azide 2-D Honeycombs. Study of the Ferromagnetic Ordered Phase below 20 K. <i>Inorganic Chemistry</i> , 2000, 39, 4688-4695.	1.9	79
201	Structure and magnetic behaviour of $[\{ Mn_2(3-Et,4-Mepy)_6(\frac{1}{4}1,1-N_3)_2(\frac{1}{4}1,3-N_3) \}_n] [PF_6]_n$ and $[\{ Mn_2(3-ampy)_4(\frac{1}{4}1,1-N_3)_2(\frac{1}{4}1,3-N_3)(N_3)(H_2O) \cdot 4H_2O \}_n]$, two alternating ferro-antiferromagnetic one-dimensional compounds. <i>Dalton Transactions RSC</i> , 2000, , 413-416.	2.3	32
202	Two new one-dimensional systems with end-to-end single dicyanamide bridges between manganese(II) centres: structural and magnetic properties. <i>Dalton Transactions RSC</i> , 2000, , 2627-2630.	2.3	59
203	Two New One-Dimensional Compounds with End-to-End Dicyanamide as a Bridging Ligand: Syntheses and Structural Characterization of $trans-[Mn(4-bzpy)_2(N(CN)_2)_2]_n$ and $cis-[Mn(Bpy)(N(CN)_2)_2]_n$, (4-bzpy) Tj ETQq 11.10.7843146rgBT	1.1	14
204	Three New Polynuclear Copper(II) Complexes with the Symmetric $[Cu(\frac{1}{4}1,1-N_3)_2Cu]_2$ Core and Pyridine Derivatives: Syntheses, Structure, and Magnetic Behavior. <i>Inorganic Chemistry</i> , 2000, 39, 2107-2112.	1.9	121
205	Two new nickel(II) cubane compounds derived from pyridine-2-methoxide (Pym): $\{Ni_4(Pym)_4Cl_4(CH_3OH)_4\}$ and $\{Ni_4(Pym)_4(N_3)_4(CH_3OH)_4\}$. Crystal structures and magnetic properties. <i>Polyhedron</i> , 1999, 18, 909-914.	1.0	85
206	Synthesis, crystal structure and magnetic behaviour of the new linear copper(II) tetranuclear compound $[Cu_4(\frac{1}{2}2-AcO)_6(AcO)_2(DPO)_2]$ (DPO=2,3-bis(2-pyridyl)quinoxaline). <i>Inorganica Chimica Acta</i> , 1999, 286, 62-66.	1.2	29
207	Versatility of the cyanato ligand: structure and magnetic properties of four new copper(II) cyanato compounds derived from tridentate amines. <i>Inorganica Chimica Acta</i> , 1999, 286, 189-196.	1.2	36
208	Polynuclear NiII and MnII azido bridging complexes. Structural trends and magnetic behavior. <i>Coordination Chemistry Reviews</i> , 1999, 193-195, 1027-1068.	9.5	802
209	Structural Characterization and Magnetic Behaviour of the Ferro-Antiferromagnetic Alternating Manganese Azido Chain $[Mn(3-Et,4-Mepy)_2(\frac{1}{4}1-N_3)_2]_n$ (3-Et,4-Mepy = 3-Ethyl-4-methylpyridine). <i>European Journal of Inorganic Chemistry</i> , 1999, 1999, 687-691.	1.0	40
210	Constrained ferromagnetic coupling in dinuclear $\frac{1}{4}1,3$ -azido nickel(II) cryptate compounds. Crystal structure and magnetic behaviour of $[Ni_2(L_1)(N_3)(H_2O)] [CF_3SO_3]_3 \cdot 2H_2O \cdot EtOH$ $\{L_1 = N[(CH_2)_2NHCH_2(C_6H_4-m)CH_2NH(CH_2)_2]_3\}$. <i>Journal of the Chemical Society Dalton Transactions</i> , 1999, , 223-228.	1.1	72
211	Dinuclear nickel(II) complexes with a tridentate nitrito bridge and terminal thiocyanato ligands. Crystal structure and magnetic properties. <i>Journal of the Chemical Society Dalton Transactions</i> , 1999, , 3115-3119.	1.1	9
212	Synthesis, Structural Characterization, Magnetic Behavior, and Single Crystal EPR Spectra of Three New One-Dimensional Manganese Azido Systems with FM, Alternating FM-AF, and AF Coupling. <i>Inorganic Chemistry</i> , 1999, 38, 5716-5723.	1.9	177
213	catena- $(\frac{1}{4}N,S-NCS)$ $[Ni(Medien)(NCS)]$: a new 1D nickel(II) complex with bridging thiocyanato ligands in cis position. Crystal structure and magnetic properties. <i>Inorganica Chimica Acta</i> , 1998, 269, 313-316.	1.2	41
214	Synthesis, structural characterization and magnetic behaviour of two new $(\frac{1}{4}1,3-N_3)_2$ dinuclear nickel compounds: study of the magnetic transition at 235 K for the $cis-(\frac{1}{4}1,3-N_3)_2[Ni(dl-cth)]_2(ClO_4)_2$ compound. <i>Inorganica Chimica Acta</i> , 1998, 278, 43-50.	1.2	25
215	Syntheses, X-ray structures, spectroscopic and magnetic studies of two new polymeric copper (II) azide complexes; $[Cu(3-ethylpyridine)_2(N_3)_2]_n$ and $[Cu(4-ethylpyridine)_2(N_3)_2]_n$. <i>Polyhedron</i> , 1998, 17, 4265-4273.	1.0	26
216	Synthesis and crystal structure of the ferro-antiferro alternating chain containing $\frac{1}{4}1,1,3$ -azido bridging ligands $[(\frac{1}{4}1,1,3-N_3)_2 \{Cu_2(Me_2Eten)_2(N_3)_2\}]_n$. Magnetic behavior. <i>Polyhedron</i> , 1998, 18, 211-215.	1.0	32

#	ARTICLE	IF	CITATIONS
217	Synthesis, crystal structure and magnetic behaviour of $(\frac{1}{4}3\text{-CO}_3)[\text{Ni}_3(\text{Medpt})_3(\text{NCSe})_4]$, a new example of trinuclear nickel(II) complex with pentadentate carbonato bridge and strong antiferromagnetic coupling. <i>Polyhedron</i> , 1998, 18, 377-381.	1.0	17
218	Structure and magnetic behaviour of the one-dimensional $[\{\text{Ni}_2(\text{Medien})_2(\frac{1}{4}1,1\text{-N}_3)_2(\frac{1}{4}1,3\text{-N}_3)\}_n][\text{ClO}_4]_n$ compound with unusually strong ferro-antiferromagnetic alternating interactions. <i>Journal of the Chemical Society Dalton Transactions</i> , 1998, , 3905-3910.	1.1	28
219	Spin frustration in the butterfly-like tetrameric $[\text{Ni}_4(\frac{1}{4}\text{-CO}_3)_2(\text{aetpy})_8][\text{ClO}_4]_4$ [aetpy = (2-aminoethyl)pyridine] complex. Structure and magnetic properties. <i>Journal of the Chemical Society Dalton Transactions</i> , 1998, , 3473-3478.	1.1	31
220	Synthesis and Structural Characterization of the One-Dimensional $[\text{Cu}(\text{3-Clpy})_2(\text{N}_3)_2]_n$ Complex (3-Clpy) $T_j \text{ ETQq0 0 0 rgBT /Overlock 1}$ 4466-4469.	1.9	102
221	Crystal Structure and Magnetic Behavior of Two New Dinuclear Carbonato-Bridged Copper(II) Compounds. Superexchange Pathway for the Different Coordination Modes of the Carbonato Bridge in Polynuclear Copper(II) Compounds. <i>Inorganic Chemistry</i> , 1998, 37, 4190-4196.	1.9	53
222	Antiferromagnetic Alternating and Homogeneous Manganese ^{II} Azido Chains: A Structural Characterization and Magnetic Behavior of Two New One-Dimensional $[\text{Mn}(\text{L})_2(\frac{1}{4}1,3\text{-N}_3)_2]_n$ Compounds (L) $T_j \text{ ETQq0 0 0 rgBT /Overlo}$	1.9	102
223	Bis(didentate) open-chain and tetradentate pseudo-macrocyclic bridging co-ordination modes of N,N ² -bis(1,3-dimethyl-5-nitroso-2,4-dioxopyrimidin-6-yl)butane-1,4-diamine. <i>Journal of the Chemical Society Dalton Transactions</i> , 1997, , 1685-1690.	1.1	7
224	Syntheses of several kinds of one-dimensional nickel(II)-nitrito complexes: a new $\frac{1}{4}$ -nitrito nickel(II) alternating chain. <i>Journal of the Chemical Society Dalton Transactions</i> , 1997, , 531-536.	1.1	42
225	Two new tetranuclear $\frac{1}{4}4$ -carbonato copper(II) complexes. Syntheses, crystal structure and magnetic behaviour of $[(\frac{1}{4}4\text{-CO}_3)(\frac{1}{4}\text{-Br})_2\{\text{Cu}_4(\text{bapa})_4\}]\text{Br}_4$ and $[(\frac{1}{4}4\text{-CO}_3)(\frac{1}{4}\text{-Cl})_2\{\text{Cu}_4(\text{bapma})_4\}]\text{Cl}_4 \cdot 12\text{H}_2\text{O}$ [bapa = bis(aminopropyl)amine and bapma = bis(aminopropyl)methylamine]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1997, , 2315-2320.	1.1	35
226	A novel tridentate co-ordination mode for the carbonatonickel system. <i>Journal of the Chemical Society Dalton Transactions</i> , 1997, , 403-408.	1.1	20
227	A new two-dimensional manganese(II) azide polymer. Synthesis, structure and magnetic properties of $[\{\text{Mn}(\text{minc})_2(\text{N}_3)_2\}_n]$ (minc = methyl isonicotinate). <i>Journal of the Chemical Society Dalton Transactions</i> , 1997, , 4431-4434.	1.1	67
228	A New Family of High-Dimensional Molecular Magnets Built from the Manganese ^{II} Azido System. Syntheses, Structures, and Magnetic Characterization of Two New Ferro ^{II} Antiferromagnetic Two-Dimensional Complexes. <i>Inorganic Chemistry</i> , 1997, 36, 3440-3446.	1.9	113
229	Structure and Magnetic Behavior of a New 1-D Compound with Simultaneous End-On Azido and Carboxylato Bridges. Unexpected Strong Ferromagnetic Coupling for a Cu ^{II} -N ^{III} -Cu Bond Angle of 111.9 $^\circ$ as a Consequence of Ligand HOMOs Countercomplementarity. <i>Inorganic Chemistry</i> , 1997, 36, 1233-1236.	1.9	124
230	Structurally alternating copper(II) chains from oxalate and azide bridging ligands: syntheses and crystal structure of $[\text{Cu}_2(\frac{1}{4}\text{-ox})\{\text{deen}\}_2(\text{H}_2\text{O})_2(\text{ClO}_4)_2]$ and $[\{\text{Cu}_2(\frac{1}{4}\text{-N}_3)\{\text{deen}\}_2(\frac{1}{4}\text{-ox})\}_n][\text{ClO}_4]_n$ (deen = Et ₂ NCH ₂ CH ₂ NH ₂). <i>Journal of the Chemical Society Dalton Transactions</i> , 1997, , 167-172.	1.1	35
231	Magnetic Studies on $\frac{1}{4}$ -Azido Polynuclear Nickel(II) Compounds with the 222-tet Ligand. Crystal Structure of $(\frac{1}{4}\text{-N}_3)_2[\text{Ni}(\text{222-tet})_2(\text{BPh}_4)_2]$ (222-tet = Triethylenetetramine) and EXAFS Structural Characterization of the Triangular Compounds $(\frac{1}{4}\text{-N}_3)_3[\text{Ni}(\text{222-tet})_3(\text{X})_3]$ (X = PF ₆ ⁻ , ClO ₄ ⁻). <i>Inorganic Chemistry</i> , 1997, 36, 4633-4640.	1.9	54
232	Syntheses, Structure, and Magnetic Behavior of Two New Nickel(II) and Cobalt(II) Dinuclear Complexes with 1,4-Dicarboxylatopyridazine. MO Calculations of the Superexchange Pathway through the Pyridazine Bridge. <i>Inorganic Chemistry</i> , 1997, 36, 2511-2516.	1.9	63
233	Synthesis, structural characterisation and magnetic behaviour of three new dinuclear end-to-end thiocyanato bridged copper(II) compounds derived from tridentate amines. <i>Inorganica Chimica Acta</i> , 1997, 255, 7-12.	1.2	51
234	Synthesis, crystal structure and magnetic properties of the triply bridged one-dimensional polymers $[\text{Cu}_2(\text{ClO}_4)(\text{OAc})_2(\text{bppz})(\text{H}_2\text{O})_2](\text{ClO}_4)(\text{H}_2\text{O})_n$ and $[\text{Cu}_2(\text{OBz})_3(\text{bppz})](\text{ClO}_4)(\text{H}_2\text{O})_{2n}$; bppz = 2,5-bis(2-pyridyl)pyrazine. <i>Inorganica Chimica Acta</i> , 1997, 260, 189-198.	1.2	17

#	ARTICLE	IF	CITATIONS
235	Significant antiferromagnetic exchange in copper(II) and nickel(II) dinuclear complexes of the substituted pyrazine ligand 2,3,5,6-tetra(2-pyridyl)pyrazine (TPPZ): magnetic properties, synthesis and crystal structure. <i>Inorganica Chimica Acta</i> , 1997, 257, 89-97.	1.2	75
236	Three new mononuclear nickel(II) cyanate and isocyanate compounds derived from macrocyclic ligands: [Ni(TMCY)(NCO)](ClO ₄), [Ni(m-CTH)(OCN)1.5(ClO ₄)0.5]. <i>Inorganica Chimica Acta</i> , 1997, 261, 227-232.	1.2	24
237	Dinuclear copper(II) complexes containing nitroso-oximate bridges: two different copper(II)-bridging entities from the same precursor. <i>Polyhedron</i> , 1997, 16, 281-289.	1.0	22
238	Synthesis, Crystal Structure, and Magnetic Behavior of (1/4-3-CO ₃)[Cu ₃ (Medpt) ₃ (ClO ₄) ₃](ClO ₄): A New Copper(II) Carbonato-Bridged Complex with a Triangular Array. <i>Inorganic Chemistry</i> , 1996, 35, 248-251.	1.9	56
239	A Novel Pentadentate Coordination Mode for the Carbonato Bridge: Synthesis, Crystal Structure, and Magnetic Behavior of (1/4-3-CO ₃)[Ni ₃ (Medpt) ₃ (NCS) ₄], a New Trinuclear Nickel(II) Carbonato-Bridged Complex with Strong Antiferromagnetic Coupling. <i>Inorganic Chemistry</i> , 1996, 35, 3094-3098.	1.9	63
240	Synthesis and Structural Characterization of [Mn(ethyl isonicotinate) ₂ (N ₃) ₂] _n , a Two-Dimensional Alternating Ferromagnetic/Antiferromagnetic Compound. Magnetostructural Correlations for the End-to-End Pseudohalide Manganese System. <i>Inorganic Chemistry</i> , 1996, 35, 6386-6391.	1.9	140
241	Synthesis, magnetic behaviour and structural characterization of the alternating hexanuclear copper(II) compound [Cu ₆ (tmen) ₆ (μ ₃ -N ₃) ₂ (μ ₃ -C ₂ O ₄) ₃ (H ₂ O) ₂][ClO ₄] ₄ ·2H ₂ O (tmen = Me ₂ NCH ₂ CH ₂ NMe ₂). <i>Journal of the Chemical Society Dalton Transactions</i> , 1996, , 1835-1838.	1.1	23
242	Structure and magnetic behaviour of the first singly bridged nickel cyanate chain and a new dinuclear complex: an approximation to the superexchange mechanism for the nickel pseudohalide system. <i>Journal of the Chemical Society Dalton Transactions</i> , 1996, , 1013-1019.	1.1	59
243	Synthesis, structure and magnetic properties of several nickel-azido complexes with tetraamines as blocking ligands. <i>Inorganica Chimica Acta</i> , 1996, 247, 85-91.	1.2	56
244	Aqueous syntheses and crystal structures of the hexa- and pentacoordinated nickel(II) isothiocyanato derivatives of bulky triamines: (H ₂ Et ₅ dien) ₂ [Ni(NCS) ₆], [Ni(Me ₅ dien)(NCS) ₂] and [Ni(Me ₄ Etdien)(NCS) ₂]. <i>Inorganica Chimica Acta</i> , 1996, 248, 59-65.	1.2	21
245	Reactivity of pyrrole pigments, XXI: Structure and reactivity of Cu(II) and Zn(II) bilindione chelates. <i>Monatshefte für Chemie</i> , 1996, 127, 1051-1062.	0.9	10
246	Coordination behaviour of 2-guanidinobenzimidazole towards cobalt(II), nickel(II), copper(II) and zinc(II). An experimental and theoretical study. <i>Transition Metal Chemistry</i> , 1996, 21, 31-37.	0.7	25
247	Crystal Structure of [Ni(CTH)(NCS) ₂] with Linearly and Nonlinearly Coordinated Isothiocyanate Groups.. <i>Acta Chemica Scandinavica</i> , 1996, 50, 178-180.	0.7	5
248	Magnetic studies on the 1/4-oxalato nickel(II) dimers with an [NiN ₄ O ₂] environment. Crystal structure of (1/4-ox)-[Ni(323-tet)] ₂ (ClO ₄) ₂ ·2H ₂ O (323-tet=N,N'-bis(3-aminopropyl)-1,2-ethanediamine). <i>Inorganica Chimica Acta</i> , 1995, 232, 151-156.	1.2	38
249	A heterodinuclear anti oximate-bridged copper(II)-nickel(II) complex: crystal structure and magnetic properties. <i>Inorganica Chimica Acta</i> , 1995, 234, 61-65.	1.2	17
250	Magneto-structural correlations for the S = 1 alternating chain [{Ni ₂ (dpt) ₂ (1/4-μ ² -N ₃) ₂] _n (ClO ₄) _n . <i>Polyhedron</i> , 1995, 14, 2133-2138.	1.0	31
251	X-ray structure determination and magnetic behavior of the new uniform S = 1 chain [{Ni(Me ₂ [14]-1,3-dieneN ₄)(μ ² -N ₃) ₂] _n (ClO ₄) _n . Magneto-structural correlations. <i>Inorganic Chemistry</i> , 1995, 34, 1278-1281.	1.9	68
252	X-ray structure determination and magnetic behavior of the new uniform S = 1 chain [{Ni(Me ₂ [14]-1,3-dieneN ₄)(μ ² -N ₃) ₂] _n (ClO ₄) _n . Magneto-structural correlations. [Erratum to document cited in CA123:46276]. <i>Inorganic Chemistry</i> , 1995, 34, 5096-5096.	1.9	1

#	ARTICLE	IF	CITATIONS
253	Magnetic Transition and Structural Asymmetrization in the Ferromagnetic Compound $\{[Ni_2(Medpt)_2(N_3)_2] \cdot (\mu_1, \mu_2-N_3)_2\}$, an Example of a Dynamic Second-Order Jahn-Teller Effect. <i>Inorganic Chemistry</i> , 1995, 34, 1793-1798.	1.9	101
254	Ten-Membered Rings of Coppers Interconnected by 2,5-Bis(2-pyridyl)pyrazine and Acetate Groups: Synthesis, Crystal Structure, and Magnetic Properties of the Two-Dimensional Polymer catena-(Octakis(μ_2 -acetato)[2,5-bis(2-pyridyl)pyrazine]tetracopper(II)). <i>Inorganic Chemistry</i> , 1995, 34, 1946-1949.	1.9	44
255	Structure and Magnetic Behavior of a New Two-Dimensional Antiferromagnetic Manganese(II)- μ_1, μ_2 -1,3-Azido System. <i>Inorganic Chemistry</i> , 1995, 34, 5707-5708.	1.9	105
256	A mixed-bridged Cu_3 cluster with an isosceles-triangular array of copper(II) ions: synthesis, crystal structure and magnetic properties. <i>Journal of the Chemical Society Dalton Transactions</i> , 1995, , 343-348.	1.1	29
257	Polynuclear Ferromagnetic Ni(II) Complexes with Pseudohalide as Bridging Ligands. <i>Advanced Materials Research</i> , 1994, 1-2, 573-580.	0.3	3
258	Magneto-Structural Correlations in 1-D μ_2 -(1,3-Azido)Nickel(II) Complexes. <i>Advanced Materials Research</i> , 1994, 1-2, 581-586.	0.3	3
259	(Tetrakis(2-pyridylmethyl)ethylenediamine)iron(II) Perchlorate. Phase Transition Study about the First Rapidly Interconverting Ferrous Spin-Crossover Complex. <i>Journal of Solid State Chemistry</i> , 1994, 109, 315-320.	1.4	9
260	$\{[Ni(3,2,3-tet)(\mu_2-NCO)]_n\}(ClO_4)_n$: structural characterization of the first monodimensional nickel(II)-cyanato system with a description of its ferromagnetic coupling. <i>Inorganica Chimica Acta</i> , 1994, 216, 5-7.	1.2	19
261	Octahedral μ_2 -oxalato-nickel(II) dinuclear complexes with μ_2 and tridentate amines as blocking ligands: magnetostructural correlations. <i>Inorganica Chimica Acta</i> , 1994, 216, 139-145.	1.2	49
262	Crystal structure and magnetic behaviour of a new kind of one-dimensional nickel(II) thiocyanate compound $\{[NiL(SCN)(\mu_2-SCN)]_n\}$ [L = bis(3-aminopropyl)methylamine]. <i>Journal of the Chemical Society Dalton Transactions</i> , 1994, , 259-262.	1.1	53
263	Synthesis and crystal structure determination of a one-dimensional end-to-end cyanate-bridged copper(II) compound. <i>Journal of the Chemical Society Dalton Transactions</i> , 1994, , 3005-3008.	1.1	22
264	Heterodinuclear Copper(II)-Nickel(II) Complexes with Unusual Asymmetrical Bridges from a New and Versatile Dioxime Multidentate Ligand. Magnetostructural Study. <i>Inorganic Chemistry</i> , 1994, 33, 3914-3924.	1.9	54
265	Crystal Structure and Magnetic Properties of $\{[Ni_2(dpt)_2(\mu_2-ox)(\mu_2-N_3)]_n\}(PF_6)_n$: A New Strategy To Obtain S = 1 Alternating Chains. <i>Inorganic Chemistry</i> , 1994, 33, 6007-6011.	1.9	66
266			

#	ARTICLE	IF	CITATIONS
271	Two new μ -azido nickel(II) uniform chains: syntheses, structures, and magneto-structural correlations. <i>Inorganic Chemistry</i> , 1993, 32, 3727-3732.	1.9	140
272	New dinuclear penta- and hexacoordinated nickel(II) complexes with μ -azido bridges. Crystal structures of ferromagnetically coupled $(\mu$ -N ₃) ₂ [Ni(Me ₃ [12]N ₃)] ₂ (ClO ₄) ₂ ·2H ₂ O and $(\mu$ -N ₃) ₂ [Ni(232-N ₄)] ₂ (ClO ₄) ₂ . <i>Inorganic Chemistry</i> , 1993, 32, 1920-1924.	1.9	118
273	The first 1D nickel(II) complex with a single azido bridge: structure and magnetic behavior of catena- $(\mu$ -N ₃)[Ni(1,4,8,11-tetraazacyclotetradecane)](ClO ₄)·H ₂ O. <i>Inorganic Chemistry</i> , 1993, 32, 1033-1035.	1.9	83
274	The first nickel(II) alternating chain with two different end-to-end azido bridges. <i>Inorganic Chemistry</i> , 1992, 31, 1726-1728.	1.9	68
275	New nickel(II)-copper(II) heterodinuclear complexes with hexa- and pentacoordinated nickel(II) ions. Magnetostructural correlations. <i>Inorganic Chemistry</i> , 1992, 31, 2627-2633.	1.9	80
276	Magnetic behaviour of pentacoordinate di- $\frac{1}{4}$ -hydroxo complexes of copper(II), nickel(II) and cobalt(II) with 2,4,4-trimethyl-1,5,9-triazacyclododec-1-ene and 2,4,4,9-tetramethyl-1,5,9-triazacyclododec-1-ene macrocycles. <i>Polyhedron</i> , 1992, 11, 453-456.	1.0	28
277	Magnetic relationship in new NiII—CuII—NiII trinuclear complexes according to nickel(II) geometry. <i>Polyhedron</i> , 1992, 11, 857-862.	1.0	21
278	Magnetic transition in ferromagnetically coupled dimer of Ni(II) with Di- $\frac{1}{4}$ -azido bridge. <i>Journal of Magnetism and Magnetic Materials</i> , 1992, 110, 181-184.	1.0	50
279	Complexes with diimine ligands. Part III. Synthesis, structure and magnetic studies of mixed acetylacetonatecobalt(II) derivatives. <i>Inorganica Chimica Acta</i> , 1991, 181, 51-60.	1.2	35
280	Pentacoordinate nickel(II) complexes with one 1,1-dithio and one macrocyclic ligand. Electro-oxidation to nickel(III) in solution. <i>Polyhedron</i> , 1991, 10, 2025-2029.	1.0	5
281	Complexes with diimine ligands. Part II. Synthesis, structure and magnetic studies of mixed acetylacetonatenickel(II) derivatives. <i>Inorganica Chimica Acta</i> , 1990, 177, 161-166.	1.2	20
282	Synthesis, structure and electrochemical behaviour of octahedral nickel(II) complexes with one 1,1-dithio and one macrocyclic ligand. <i>Polyhedron</i> , 1990, 9, 1729-1735.	1.0	10
283	Synthesis, structure and EPR studies of mixed hexafluoroacetylacetonatecopper(II) Complexes with some Diimine Ligands. <i>Inorganica Chimica Acta</i> , 1989, 162, 97-103.	1.2	47
284	Solid state kinetic parameters and chemical mechanism of the dehydration of CoCl ₂ ·6H ₂ O: An inorganic solid state experiment. <i>Journal of Chemical Education</i> , 1988, 65, 85.	1.1	4
285	δ H and activation energy of dehydration processes calculated from dynamic TG and DTA. <i>Thermochimica Acta</i> , 1986, 101, 77-82.	1.2	0
286	Solid state kinetic parameters of the first decomposition of potassium dichlorotetracyanoplatinate(IV), dibromotetracyanoplatinate(IV) and diiodotetracyanoplatinate(IV). <i>Thermochimica Acta</i> , 1986, 104, 299-307.	1.2	2
287	Simple $g(\pm)$ non dependent method to solve the kinetic parameters by isothermal thermogravimetry. <i>Thermochimica Acta</i> , 1986, 104, 309-319.	1.2	5
288	Solid state kinetic parameters of the dehydration of CoCl ₂ ·6H ₂ O by isothermal thermogravimetric measurements. <i>Thermochimica Acta</i> , 1986, 102, 125-135.	1.2	20

#	ARTICLE	IF	CITATIONS
289	Solid state kinetic parameters and mechanism for the deaquation-anation of hexacyanocobaltate(III) of aquopentaamminecobalt(III), rhodium(III) and irid. <i>Thermochimica Acta</i> , 1986, 102, 137-148.	1.2	3
290	Dinuclear compounds with a $\frac{1}{4}$ -cyano ligand. Part XIII. Synthesis, characterization and solid state kinetics of the formation of $(\frac{1}{4}$ -cyano)(tricyanometal(II))pentaamminerhodium(III) and iridium(III) (metal(II) = nickel, palladium, platinum). Influence of the Pt- π , Pt interactions. <i>Inorganica Chimica Acta</i> , 1986, 116, 157-162.	1.2	5
291	Dinuclear complexes with a $\frac{1}{2}$ -cyano ligand. Part XI(1). Synthesis and characterization of several derivatives of cis-[Co(NH ₃)(en) ₂ (H ₂ O)] ³⁺ . <i>Transition Metal Chemistry</i> , 1985, 10, 466-469.	0.7	6
292	Dinuclear complexes with a $\frac{1}{4}$ -cyano ligand. Part X. Synthesis and characterization of several derivatives of cis-diaquaamminecobalt(III) complexes. Influence of pH. <i>Inorganica Chimica Acta</i> , 1985, 104, 143-147.	1.2	10
293	Simple numerical and graphical method to solve the kinetic parameters by dynamic thermogravimetry. <i>Thermochimica Acta</i> , 1985, 91, 107-114.	1.2	4
294	Kinetic parameters and solid state mechanism in inorganic coordination compounds. <i>Thermochimica Acta</i> , 1985, 92, 493-496.	1.2	1
295	Solid-state kinetic parameters and mechanism for the deaquation-anation of hexacyanochromate(III) and nitrosylpentacyanochromate(I) of aquopentaamminechromium(III), -cobalt(III), -rhodium(III), and -iridium(III). <i>Inorganic Chemistry</i> , 1985, 24, 1874-1878.	1.9	20
296	Kinetic parameters and solid state mechanism of the thermal dehydration of [M(H ₂ O)(NH ₃) ₅][Co(CN) ₆] [M = Co(III), Rh(III) and Ir(III)]. <i>Thermochimica Acta</i> , 1984, 76, 201-211.	1.2	14
297	Kinetics of the thermal dehydration of trans-fluoroaquo-bis(ethylenediamine)chromium(III) tetracyanomethylate(II) [metal(II) = Ni(II), Pd(II) and Pt(II)]. <i>Thermochimica Acta</i> , 1984, 80, 103-113.	1.2	8
298	Kinetic parameters and solid state mechanism of the thermal dehydration of trans CrF(H ₂ O)(en) ₂ 2, trans CrF(H ₂ O)(tn) ₂ 2H ₂ O and trans CrF(H ₂ O)(en)(tn) 2H ₂ O (en = ethylenediamine; tn = Tj ETQq0 0 0 rg 1.2/Overlap 10 Tf 50	1.2	10
299	Dinuclear complexes with a $\frac{1}{2}$ -cyano ligand. Part VIII(1). Synthesis and characterization of several derivatives of cis-[CrX(H ₂ O)(en) ₂] ²⁺ (X=Cl or Br). <i>Transition Metal Chemistry</i> , 1984, 9, 287-290.	0.7	9
300	Solid-state kinetic parameters for the deaquation-anation of the tetracyanonickelate(II), tetracyanopalladate(II), and tetracyanoplatinate(II) of aquapentaamminecobalt(III). Influence of the metal-metal interactions on the platinum complex. <i>Inorganic Chemistry</i> , 1984, 23, 2236-2242.	1.9	23
301	Thermochemical parameters of the thermal dehydration of trans-[CrF(H ₂ O)(1,3-diaminopropane) ₂][M(CN) ₄] (M = Pd, Pt). <i>Thermochimica Acta</i> , 1983, 64, 237-246.	1.2	12
302	Dinuclear complexes with a $\frac{1}{2}$ -cyano-ligand. ? Part VII. Synthesis and characterization of several derivatives of trans-[CrF(H ₂ O)(1,2-diaminopropane) ₂] ²⁺ . <i>Transition Metal Chemistry</i> , 1983, 8, 87-90.	0.7	11
303	Kinetic parameters and solid state mechanism of the thermal dehydration of trans[CrF(H ₂ O)(aaâ€²)2]K[Cr(CN) ₆]H ₂ O And trans[CrF(H ₂ O)(aaâ€²)2]K[CrNO(CN) ₅]H ₂ O (aaâ€²=ethylenediamine) Tj ETQq1 1 0.7 84	1.2	10
304	Thermochemical parameters of the thermal dehydration of trans[CrF(H ₂ O)(1,3-diaminopropane) ₂][Ni(CN) ₄]. <i>Thermochimica Acta</i> , 1982, 56, 183-191.	1.2	18
305	Synthesis and characterization of cyano-bridged compounds (IV) Chromium-nickel, chromium-palladium and chromium-platinum dinuclear complexes. <i>Journal of Inorganic and Nuclear Chemistry</i> , 1981, 43, 3113-3118.	0.5	23