

Trevor H Yeats

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5405027/trevor-h-yeats-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

27
papers

2,432
citations

18
h-index

30
g-index

30
ext. papers

3,150
ext. citations

7.3
avg, IF

5.32
L-index

#	Paper	IF	Citations
27	The formation and function of plant cuticles. <i>Plant Physiology</i> , 2013 , 163, 5-20	6.6	664
26	Arabidopsis LTPG is a glycosylphosphatidylinositol-anchored lipid transfer protein required for export of lipids to the plant surface. <i>Plant Cell</i> , 2009 , 21, 1230-8	11.6	234
25	The biochemistry and biology of extracellular plant lipid-transfer proteins (LTPs). <i>Protein Science</i> , 2008 , 17, 191-8	6.3	204
24	Fruit Softening: Revisiting the Role of Pectin. <i>Trends in Plant Science</i> , 2018 , 23, 302-310	13.1	196
23	Tissue- and cell-type specific transcriptome profiling of expanding tomato fruit provides insights into metabolic and regulatory specialization and cuticle formation. <i>Plant Cell</i> , 2011 , 23, 3893-910	11.6	162
22	Adaptive horizontal transfer of a bacterial gene to an invasive insect pest of coffee. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 4197-202	11.5	157
21	The identification of cutin synthase: formation of the plant polyester cutin. <i>Nature Chemical Biology</i> , 2012 , 8, 609-11	11.7	142
20	The Implications of Lignocellulosic Biomass Chemical Composition for the Production of Advanced Biofuels. <i>BioScience</i> , 2014 , 64, 192-201	5.7	87
19	Two oxidosqualene cyclases responsible for biosynthesis of tomato fruit cuticular triterpenoids. <i>Plant Physiology</i> , 2011 , 155, 540-52	6.6	83
18	The fruit cuticles of wild tomato species exhibit architectural and chemical diversity, providing a new model for studying the evolution of cuticle function. <i>Plant Journal</i> , 2012 , 69, 655-66	6.9	72
17	An ATP binding cassette transporter is required for cuticular wax deposition and desiccation tolerance in the moss <i>Physcomitrella patens</i> . <i>Plant Cell</i> , 2013 , 25, 4000-13	11.6	71
16	Tomato Cutin Deficient 1 (CD1) and putative orthologs comprise an ancient family of cutin synthase-like (CUS) proteins that are conserved among land plants. <i>Plant Journal</i> , 2014 , 77, 667-75	6.9	70
15	Mining the surface proteome of tomato (<i>Solanum lycopersicum</i>) fruit for proteins associated with cuticle biogenesis. <i>Journal of Experimental Botany</i> , 2010 , 61, 3759-71	7	64
14	Cloning and characterization of a lupeol synthase involved in the synthesis of epicuticular wax crystals on stem and hypocotyl surfaces of <i>Ricinus communis</i> . <i>Archives of Biochemistry and Biophysics</i> , 2006 , 448, 60-72	4.1	50
13	Cloning and characterization of oxidosqualene cyclases from <i>Kalanchoe daigremontiana</i> : enzymes catalyzing up to 10 rearrangement steps yielding friedelin and other triterpenoids. <i>Journal of Biological Chemistry</i> , 2010 , 285, 29703-12	5.4	48
12	Plant glycosylphosphatidylinositol anchored proteins at the plasma membrane-cell wall nexus. <i>Journal of Integrative Plant Biology</i> , 2018 , 60, 649-669	8.3	30
11	Cellulose Deficiency Is Enhanced on Hyper Accumulation of Sucrose by a H ⁺ -Coupled Sucrose Symporter. <i>Plant Physiology</i> , 2016 , 171, 110-24	6.6	29

10	Multivariate Genome-Wide Association Analyses Reveal the Genetic Basis of Seed Fatty Acid Composition in Oat (L.). <i>G3: Genes, Genomes, Genetics</i> , 2019 , 9, 2963-2975	3.2	20
9	Postharvest changes in LIN5-down-regulated plants suggest a role for sugar deficiency in cuticle metabolism during ripening. <i>Phytochemistry</i> , 2017 , 142, 11-20	4	16
8	Heritable temporal gene expression patterns correlate with metabolomic seed content in developing hexaploid oat seed. <i>Plant Biotechnology Journal</i> , 2020 , 18, 1211-1222	11.6	8
7	Translating insights from the seed metabolome into improved prediction for lipid-composition traits in oat (<i>Avena sativa</i> L.). <i>Genetics</i> , 2021 , 217,	4	7
6	Subcellular Spice Trade Routes: Crocin Biosynthesis in the Saffron Crocus (). <i>Plant Physiology</i> , 2018 , 177, 869-870	6.6	6
5	Multi-omics prediction of oat agronomic and seed nutritional traits across environments and in distantly related populations. <i>Theoretical and Applied Genetics</i> , 2021 , 134, 4043-4054	6	4
4	Setting and Diffusing the Cyanide Bomb in Plant Defense. <i>Plant Physiology</i> , 2018 , 178, 956-957	6.6	4
3	Improving Genomic Prediction for Seed Quality Traits in Oat (<i>Avena sativa</i> L.) Using Trait-Specific Relationship Matrices. <i>Frontiers in Genetics</i> , 2021 , 12, 643733	4.5	3
2	Multivariate Genome-wide Association Analyses Reveal the Genetic Basis of Seed Fatty Acid Composition in Oat (<i>Avena sativa</i> L.)		1
1	Buffering Lipid Synthesis by Conditional Inhibition. <i>Plant Physiology</i> , 2019 , 181, 8	6.6	