

Serena Camerini

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/541462/serena-camerini-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

49
papers

1,226
citations

19
h-index

34
g-index

50
ext. papers

1,450
ext. citations

6.6
avg, IF

3.9
L-index

#	Paper	IF	Citations
49	Structural basis of ubiquitination mediated by protein splicing in early Eukarya. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2021 , 1865, 129844	4	2
48	Re-Discovery of : Genomic and Functional Analysis of Viruses from Isolates. <i>Biomedicines</i> , 2021 , 9,	4.8	1
47	The Fatty Acid and Protein Profiles of Circulating CD81-Positive Small Extracellular Vesicles Are Associated with Disease Stage in Melanoma Patients. <i>Cancers</i> , 2021 , 13,	6.6	2
46	Natural-Killer-Derived Extracellular Vesicles: Immune Sensors and Interactors. <i>Frontiers in Immunology</i> , 2020 , 11, 262	8.4	46
45	Type E Botulinum Neurotoxin-Producing <i>Clostridium butyricum</i> Strains Are Aerotolerant during Vegetative Growth. <i>MSystems</i> , 2019 , 4,	7.6	3
44	HIPK2 Phosphorylates the Microtubule-Severing Enzyme Spastin at S268 for Abscission. <i>Cells</i> , 2019 , 8,	7.9	12
43	A novel disorder involving dyshematopoiesis, inflammation, and HLH due to aberrant CDC42 function. <i>Journal of Experimental Medicine</i> , 2019 , 216, 2778-2799	16.6	71
42	HIPK2 and extrachromosomal histone H2B are separately recruited by Aurora-B for cytokinesis. <i>Oncogene</i> , 2018 , 37, 3562-3574	9.2	6
41	Separase prevents genomic instability by controlling replication fork speed. <i>Nucleic Acids Research</i> , 2018 , 46, 267-278	20.1	18
40	The enzymatic processing of Edystroglycan by MMP-2 is controlled by two anchoring sites distinct from the active site. <i>PLoS ONE</i> , 2018 , 13, e0192651	3.7	2
39	Proteomic and functional analyses reveal pleiotropic action of the anti-tumoral compound NBDHEX in <i>Giardia duodenalis</i> . <i>International Journal for Parasitology: Drugs and Drug Resistance</i> , 2017 , 7, 147-158	4	13
38	Exogenous Alpha-Synuclein Alters Pre- and Post-Synaptic Activity by Fragmenting Lipid Rafts. <i>EBioMedicine</i> , 2016 , 7, 191-204	8.8	14
37	Mass spectrometry detection of fraudulent use of cow whey in water buffalo, sheep, or goat Italian ricotta cheese. <i>Food Chemistry</i> , 2016 , 197 Pt B, 1240-8	8.5	19
36	Anti-GAPDH Autoantibodies as a Pathogenic Determinant and Potential Biomarker of Neuropsychiatric Diseases. <i>Arthritis and Rheumatology</i> , 2016 , 68, 2708-2716	9.5	17
35	The FAD-dependent glycerol-3-phosphate dehydrogenase of <i>Giardia duodenalis</i> : an unconventional enzyme that interacts with the g14-3-3 and it is a target of the antitumoral compound NBDHEX. <i>Frontiers in Microbiology</i> , 2015 , 6, 544	5.7	16
34	Src inhibitors modulate frataxin protein levels. <i>Human Molecular Genetics</i> , 2015 , 24, 4296-305	5.6	9
33	The role of protein and peptide separation before mass spectrometry analysis in clinical proteomics. <i>Journal of Chromatography A</i> , 2015 , 1381, 1-12	4.5	51

32	MLC1 protein: a likely link between leukodystrophies and brain channelopathies. <i>Frontiers in Cellular Neuroscience</i> , 2015 , 9, 66	6.1	23
31	Autoantibodies specific to D4GDI modulate Rho GTPase mediated cytoskeleton remodeling and induce autophagy in T lymphocytes. <i>Journal of Autoimmunity</i> , 2015 , 58, 78-89	15.5	14
30	HIPK2 sustains apoptotic response by phosphorylating Che-1/AATF and promoting its degradation. <i>Cell Death and Disease</i> , 2014 , 5, e1414	9.8	11
29	Gaucher disease due to saposin C deficiency is an inherited lysosomal disease caused by rapidly degraded mutant proteins. <i>Human Molecular Genetics</i> , 2014 , 23, 5814-26	5.6	24
28	Native metastable prefibrillar oligomers are the most neurotoxic species among amyloid aggregates. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2014 , 1842, 1622-9	6.9	28
27	Megalencephalic leukoencephalopathy with subcortical cysts protein-1 modulates endosomal pH and protein trafficking in astrocytes: relevance to MLC disease pathogenesis. <i>Neurobiology of Disease</i> , 2014 , 66, 1-18	7.5	18
26	Inactivation of human salivary glutathione transferase P1-1 by hypothiocyanite: a post-translational control system in search of a role. <i>PLoS ONE</i> , 2014 , 9, e112797	3.7	14
25	The crystal structure of Giardia duodenalis 14-3-3 in the apo form: when protein post-translational modifications make the difference. <i>PLoS ONE</i> , 2014 , 9, e92902	3.7	8
24	Carbon monoxide signaling in human red blood cells: evidence for pentose phosphate pathway activation and protein deglutathionylation. <i>Antioxidants and Redox Signaling</i> , 2014 , 20, 403-16	8.4	16
23	The nebulin SH3 domain is dispensable for normal skeletal muscle structure but is required for effective active load bearing in mouse. <i>Development (Cambridge)</i> , 2014 , 141, e108-e108	6.6	
22	The nebulin SH3 domain is dispensable for normal skeletal muscle structure but is required for effective active load bearing in mouse. <i>Journal of Cell Science</i> , 2013 , 126, 5477-89	5.3	28
21	Genotype-phenotype analysis of S326C OGG1 polymorphism: a risk factor for oxidative pathologies. <i>Free Radical Biology and Medicine</i> , 2013 , 63, 401-9	7.8	23
20	HIPK2 catalytic activity and subcellular localization are regulated by activation-loop Y354 autophosphorylation. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2013 , 1833, 1443-53	4.9	40
19	Phosphorylation and nitration of tyrosine residues affect functional properties of Synaptophysin and Dynamin I, two proteins involved in exo-endocytosis of synaptic vesicles. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2013 , 1833, 110-21	4.9	26
18	Streptococcal-vimentin cross-reactive antibodies induce microvascular cardiac endothelial proinflammatory phenotype in rheumatic heart disease. <i>Clinical and Experimental Immunology</i> , 2013 , 173, 419-29	6.2	18
17	Interkingdom complementation reveals structural conservation and functional divergence of 14-3-3 proteins. <i>PLoS ONE</i> , 2013 , 8, e78090	3.7	9
16	An insight into the abundant proteome of 46BR.1G1 fibroblasts deficient of DNA ligase I. <i>Electrophoresis</i> , 2012 , 33, 307-15	3.6	3
15	Interaction network of the 14-3-3 protein in the ancient protozoan parasite Giardia duodenalis. <i>Journal of Proteome Research</i> , 2012 , 11, 2666-83	5.6	29

14 Phosphorylation of SRSF1 is modulated by replicational stress. *Nucleic Acids Research*, **2012**, 40, 1106-1720.1 26

13 Serum low-molecular-weight protein fractionation for biomarker discovery. *Methods in Molecular Biology*, **2012**, 823, 237-49 1.4 5

12 Efficient one-step chromatographic purification and functional characterization of recombinant human Saposin C. *Protein Expression and Purification*, **2011**, 78, 209-15 2 2

11 *Giardia duodenalis* 14-3-3 protein is polyglycylated by a tubulin tyrosine ligase-like member and deglycylated by two metalloproteases. *Journal of Biological Chemistry*, **2011**, 286, 4471-84 5.4 15

10 MLC1 trafficking and membrane expression in astrocytes: role of caveolin-1 and phosphorylation. *Neurobiology of Disease*, **2010**, 37, 581-95 7.5 27

9 Effects of indole-3-acetic acid on *Sinorhizobium meliloti* survival and on symbiotic nitrogen fixation and stem dry weight production. *Applied Microbiology and Biotechnology*, **2009**, 83, 727-38 5.7 39

8 Peroxynitrite induces tyrosine residue modifications in synaptophysin C-terminal domain, affecting its interaction with src. *Journal of Neurochemistry*, **2009**, 111, 859-69 6 14

7 Formation of an adduct by clenbuterol, a beta-adrenoceptor agonist drug, and serum albumin in human saliva at the acidic pH of the stomach: evidence for an aryl radical-based process. *Free Radical Biology and Medicine*, **2008**, 45, 124-35 7.8 7

6 Introduction of a novel pathway for IAA biosynthesis to rhizobia alters vetch root nodule development. *Archives of Microbiology*, **2008**, 190, 67-77 3 78

5 A method for the selective isolation and enrichment of carrier protein-bound low-molecular weight proteins and peptides in the blood. *Proteomics - Clinical Applications*, **2007**, 1, 176-84 3.1 15

4 A novel approach to identify proteins modified by nitric oxide: the HIS-TAG switch method. *Journal of Proteome Research*, **2007**, 6, 3224-31 5.6 79

3 Proteomics approaches to study the redox state of cysteine-containing proteins. *Annali Dell'istituto Superiore Di Sanita*, **2005**, 41, 451-7 1.6 4

2 Thiol-mediated protein retention in the endoplasmic reticulum: the role of ERp44. *EMBO Journal*, **2003**, 22, 5015-22 13 192

1 Optimisation of transgene action at the post-transcriptional level: high quality parthenocarpic fruits in industrial tomatoes. *BMC Biotechnology*, **2002**, 2, 1 3.5 89