
Jonas Esche

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5422360/publications.pdf Version: 2024-02-01



IONAS ESCHE

#	Article	IF	CITATIONS
1	Higher diet-dependent renal acid load associates with higher glucocorticoid secretion and potentially bioactive free glucocorticoids in healthy children. Kidney International, 2016, 90, 325-333.	5.2	46
2	Dietary Acid Load and Potassium Intake Associate with Blood Pressure and Hypertension Prevalence in a Representative Sample of the German Adult Population. Nutrients, 2018, 10, 103.	4.1	30
3	Health Care Costs of Spontaneous Aneurysmal Subarachnoid Hemorrhage for Rehabilitation, Home Care, and In-Hospital Treatment for the First Year. World Neurosurgery, 2017, 97, 495-500.	1.3	26
4	Contribution of iodized salt to total iodine and total salt intake in Germany. European Journal of Nutrition, 2020, 59, 3163-3169.	3.9	22
5	Urinary Citrate, an Index of Acid-Base Status, Predicts Bone Strength in Youths and Fracture Risk in Adult Females. Journal of Clinical Endocrinology and Metabolism, 2016, 101, 4914-4921.	3.6	21
6	Hyponatremia After Spontaneous Aneurysmal Subarachnoid Hemorrhage—A Prospective Observational Study. World Neurosurgery, 2019, 129, e538-e544.	1.3	20
7	Higher Glucocorticoid Secretion in the Physiological Range Is Associated With Lower Bone Strength at the Proximal Radius in Healthy Children: Importance of Protein Intake Adjustment. Journal of Bone and Mineral Research, 2015, 30, 240-248.	2.8	19
8	Dietary Potential Renal Acid Load Is Positively Associated with Serum Uric Acid and Odds of Hyperuricemia in the German Adult Population. Journal of Nutrition, 2018, 148, 49-55.	2.9	19
9	Dietâ€independent relevance of serum uric acid for blood pressure in a representative population sample. Journal of Clinical Hypertension, 2017, 19, 1042-1050.	2.0	7
10	Cortisol and 11 beta-hydroxysteroid dehydrogenase type 2 as potential determinants of renal citrate excretion in healthy children. Endocrine, 2020, 67, 442-448.	2.3	6
11	Increased body fatness adversely relates to 24-hour urine pH during childhood and adolescence: evidence of an adipo-renal axis. American Journal of Clinical Nutrition, 2019, 109, 1279-1287.	4.7	6
12	The DONALD study as a longitudinal sensor of nutritional developments: iodine and salt intake over more than 30Âyears in German children. European Journal of Nutrition, 2022, 61, 2143-2151.	3.9	6
13	Estimates of renal net acid excretion and their relationships with serum uric acid and hyperuricemia in a representative German population sample. European Journal of Clinical Nutrition, 2020, 74, 63-68.	2.9	5
14	Prospective relation of adolescent citrate excretion and net acid excretion capacity with blood pressure in young adulthood. American Journal of Physiology - Renal Physiology, 2018, 315, F1228-F1235.	2.7	4
15	Spontaneous Aneurysmal Subarachnoid Hemorrhage and Related Cortisol and Immunologic Alterations: Impact on Patients' Health-related Quality of Life. Journal of Neurological Surgery, Part A: Central European Neurosurgery, 2019, 80, 371-380.	0.8	3
16	Glucocorticoids and Body Fat Inversely Associate With Bone Marrow Density of the Distal Radius in Healthy Youths. Journal of Clinical Endocrinology and Metabolism, 2019, 104, 2250-2256.	3.6	3
17	Inflammatory mediators in the adipo-renal axis: leptin, adiponectin, and soluble ICAM-1. American Journal of Physiology - Renal Physiology, 2020, 319, F469-F475.	2.7	3
18	Health Economic Aspects of Aneurysmal Subarachnoid Hemorrhage: Factors Determining First Year In-Hospital Treatment Expenses. Journal of Neurological Surgery, Part A: Central European Neurosurgery, 2021, 82, 204-210.	0.8	3

#	Article	IF	CITATIONS
19	Renal biomarkers of acid excretion capacity: relationships with body fatness and blood pressure. European Journal of Clinical Nutrition, 2020, 74, 76-82.	2.9	2
20	Increased protein intake and corresponding renal acid load under a concurrent alkalizing diet regime. Physiological Reports, 2016, 4, e12851.	1.7	1