
Ricardo Gusmão

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5427239/publications.pdf Version: 2024-02-01



RICADOO CUSMÃEO

#	Article	IF	CITATIONS
1	The effects of a video-based randomized controlled trial intervention on depression stigma and help-seeking attitudes in university students. Psychiatry Research, 2022, 308, 114356.	3.3	9
2	The association between stigmatizing attitudes towards depression and help seeking attitudes in college students. PLoS ONE, 2022, 17, e0263622.	2.5	14
3	Psychometric properties of the Depression Stigma Scale in the Portuguese population and its association with gender and depressive symptomatology. Health and Quality of Life Outcomes, 2022, 20, 42.	2.4	1
4	A systematic review of older adults' request for or attitude toward euthanasia or assisted-suicide. Aging and Mental Health, 2021, 25, 420-430.	2.8	27
5	A New Viewpoint on the Etiopathogenesis of Depression: Insights From the Neurophysiology of Deep Brain Stimulation in Parkinson's Disease and Treatment-Resistant Depression. Frontiers in Psychiatry, 2021, 12, 607339.	2.6	7
6	Suicide time-series structural change analysis in Portugal (1913-2018): Impact of register bias on suicide trends. Journal of Affective Disorders, 2021, 291, 65-75.	4.1	4
7	A Naturalistic, European Multi-Center Clinical Study of Electrodermal Reactivity and Suicide Risk Among Patients With Depression. Frontiers in Psychiatry, 2021, 12, 765128.	2.6	2
8	The Association Between Changes in the University Educational Setting and Peer Relationships: Effects in Students' Depressive Symptoms During the COVID-19 Pandemic. Frontiers in Psychiatry, 2021, 12, 783776.	2.6	19
9	Tendências Seculares na Mortalidade por Doenças Cerebrovasculares em Portugal: 1902-2012. Revista De MedicinÄf InternÄf, Neurologe, Psihiatrie, Neurochirurgie, Dermato-venerologie MedicinÄf InternÄf, 2020, 27, 139-144.	0.0	0
10	Prevention of suicidal behaviour: Results of a controlled community-based intervention study in four European countries. PLoS ONE, 2019, 14, e0224602.	2.5	27
11	Effectiveness of General Practitioner training to improve suicide awareness and knowledge and skills towards depression. Journal of Affective Disorders, 2018, 227, 17-23.	4.1	21
12	A multi-centre, randomised, double-blind, placebo-controlled clinical trial of methylphenidate in the initial treatment of acute mania (MEMAP study). European Neuropsychopharmacology, 2018, 28, 185-194.	0.7	9
13	EUDOR-A multi-centre research program: A naturalistic, European Multi-centre Clinical study of EDOR Test in adult patients with primary depression. BMC Psychiatry, 2017, 17, 108.	2.6	7
14	Public attitudes toward depression and help-seeking: Impact of the OSPI-Europe depression awareness campaign in four European regions. Journal of Affective Disorders, 2017, 217, 252-259.	4.1	49
15	A cross-national study on gender differences in suicide intent. BMC Psychiatry, 2017, 17, 234.	2.6	204
16	Enhancing mental health literacy in young people. European Child and Adolescent Psychiatry, 2016, 25, 567-569.	4.7	91
17	Effectiveness of Depression–Suicidal Behaviour Gatekeeper Training among police officers in three European regions: Outcomes of the Optimising Suicide Prevention Programmes and Their Implementation in Europe (OSPI-Europe) study. International Journal of Social Psychiatry, 2016, 62, 651-660.	3.1	23
18	Exploring synergistic interactions and catalysts in complex interventions: longitudinal, mixed methods case studies of an optimised multi-level suicide prevention intervention in four european countries (Ospi-Europe). BMC Public Health, 2016, 16, 268.	2.9	44

RICARDO GUSMãO

#	Article	IF	CITATIONS
19	Contribution to the validation of the Kutcher Adolescent Depression Scale (KADS-6) in a Portuguese population. Psicologia: Reflexao E Critica, 2015, 28, 313-321.	0.9	3
20	What Are Reasons for the Large Gender Differences in the Lethality of Suicidal Acts? An Epidemiological Analysis in Four European Countries. PLoS ONE, 2015, 10, e0129062.	2.5	151
21	Depression Awareness and Self-Management Through the Internet: Protocol for an Internationally Standardized Approach. JMIR Research Protocols, 2015, 4, e99.	1.0	22
22	How to Adjust Media Recommendations on Reporting Suicidal Behavior to New Media Developments. Archives of Suicide Research, 2014, 18, 156-169.	2.3	28
23	Effectiveness of community facilitator training in improving knowledge, attitudes, and confidence in relation to depression and suicidal behavior: Results of the OSPI-Europe intervention in four European countries. Journal of Affective Disorders, 2014, 165, 142-150.	4.1	69
24	EPA-1387 – The case for a structured up-to-date suicide national prevention strategy based on the 4-level approach: lessons from EAAD and OSPI-Europe in Portugal. European Psychiatry, 2014, 29, 1.	0.2	0
25	Methylphenidate in mania project (MEMAP): study protocol of an international randomised double-blind placebo-controlled study on the initial treatment of acute mania with methylphenidate. BMC Psychiatry, 2013, 13, 71.	2.6	15
26	Developing social capital in implementing a complex intervention: a process evaluation of the early implementation of a suicide prevention intervention in four European countries. BMC Public Health, 2013, 13, 158.	2.9	35
27	Financial crisis, austerity, and health in Europe. Lancet, The, 2013, 382, 391.	13.7	16
28	Public attitudes toward depression and help-seeking in four European countries baseline survey prior to the OSPI-Europe intervention. Journal of Affective Disorders, 2013, 150, 320-329.	4.1	129
29	Financial crisis, austerity, and health in Europe. Lancet, The, 2013, 382, 391-392.	13.7	24
30	Media recommendations on reporting suicidal behaviour and suggestions for optimisation. Acta Psychiatrica Scandinavica, 2013, 128, n/a-n/a.	4.5	4
31	Antidepressant Utilization and Suicide in Europe: An Ecological Multi-National Study. PLoS ONE, 2013, 8, e66455.	2.5	109
32	Suicide after 65 years old: current data in Portugal. , 2013, , 321-334.		0
33	Drug suicide: a sex-equal cause of death in 16 European countries. BMC Public Health, 2011, 11, 61.	2.9	20
34	Best Practice Elements of Multilevel Suicide Prevention Strategies. Crisis, 2011, 32, 319-333.	1.2	214
35	Intermediate outcome criteria and evaluation of suicide prevention programmes: a review. Injury Prevention, 2010, 16, A234-A234.	2.4	0
36	Suicide registration in eight European countries: A qualitative analysis of procedures and practices. Forensic Science International, 2010, 202, 86-92.	2.2	54

RICARDO GUSMãO

#	Article	IF	CITATIONS
37	Munchausen syndrome mimicking psychiatric disease with concomitant genuine physical illness. BMJ Case Reports, 2010, 2010, bcr0720103142-bcr0720103142.	0.5	6
38	Optimizing Suicide Prevention Programs and Their Implementation in Europe (OSPI Europe): an evidence-based multi-level approach. BMC Public Health, 2009, 9, 428.	2.9	104
39	Gender issues in suicide rates, trends and methods among youths aged 15–24 in 15 European countries. Journal of Affective Disorders, 2009, 113, 216-226.	4.1	101
40	The â€~European Alliance Against Depression (EAAD)': A multifaceted, community-based action programme against depression and suicidality. World Journal of Biological Psychiatry, 2008, 9, 51-58.	2.6	101
41	Suicide methods in Europe: a gender-specific analysis of countries participating in the "European Alliance Against Depression". Journal of Epidemiology and Community Health, 2008, 62, 545-551.	3.7	221
42	Assessment of dysfunctional working models of self and others in schizophrenic patients: a summary of data collected in nine nations. Acta Psychiatrica Scandinavica, 2000, 102, 336-341.	4.5	4