

Wonil Oh

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5434545/publications.pdf>

Version: 2024-02-01

44
papers

3,414
citations

186265

28
h-index

233421

45
g-index

45
all docs

45
docs citations

45
times ranked

5024
citing authors

#	ARTICLE	IF	CITATIONS
1	Protective effects of human umbilical cord blood-derived mesenchymal stem cells against dexamethasone-induced apoptotic cell death in hair follicles. <i>International Journal of Molecular Medicine</i> , 2020, 45, 556-568.	4.0	5
2	Galectin-3 Secreted by Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Reduces Aberrant Tau Phosphorylation in an Alzheimer Disease Model. <i>Stem Cells International</i> , 2020, 2020, 1-14.	2.5	20
3	Soluble PTX3 of Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Attenuates Hyperoxic Lung Injury by Activating Macrophage Polarization in Neonatal Rat Model. <i>Stem Cells International</i> , 2020, 2020, 1-18.	2.5	14
4	Prospects for the therapeutic development of umbilical cord blood-derived mesenchymal stem cells. <i>World Journal of Stem Cells</i> , 2020, 12, 1511-1528.	2.8	19
5	Thrombospondin-1 secreted by human umbilical cord blood-derived mesenchymal stem cells rescues neurons from synaptic dysfunction in Alzheimer's disease model. <i>Scientific Reports</i> , 2018, 8, 354.	3.3	57
6	Effect of growth differentiation factor-15 secreted by human umbilical cord blood-derived mesenchymal stem cells on amyloid beta levels in <i>in vitro</i> and <i>in vivo</i> models of Alzheimer's disease. <i>Biochemical and Biophysical Research Communications</i> , 2018, 504, 933-940.	2.1	29
7	Early, but not late, treatment with human umbilical cord blood-derived mesenchymal stem cells attenuates cisplatin nephrotoxicity through immunomodulation. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 313, F984-F996.	2.7	20
8	hMSCs suppress neutrophil-dominant airway inflammation in a murine model of asthma. <i>Experimental and Molecular Medicine</i> , 2017, 49, e288-e288.	7.7	28
9	Optimization of culture conditions for rapid clinical-scale expansion of human umbilical cord blood-derived mesenchymal stem cells. <i>Clinical and Translational Medicine</i> , 2017, 6, 38.	4.0	19
10	Effect of Single and Double Administration of Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Following Focal Cerebral Ischemia in Rats. <i>Experimental Neurobiology</i> , 2017, 26, 55-65.	1.6	19
11	Downregulation of Melanoma Cell Adhesion Molecule (MCAM/CD146) Accelerates Cellular Senescence in Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells. <i>Stem Cells Translational Medicine</i> , 2016, 5, 427-439.	3.3	56
12	Human umbilical cord blood-derived mesenchymal stem cells improve functional recovery through thrombospondin1, pantraxin3, and vascular endothelial growth factor in the ischemic rat brain. <i>Journal of Neuroscience Research</i> , 2015, 93, 1814-1825.	2.9	33
13	Autocrine Action of Thrombospondin-2 Determines the Chondrogenic Differentiation Potential and Suppresses Hypertrophic Maturation of Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells. <i>Stem Cells</i> , 2015, 33, 3291-3303.	3.2	39
14	The Effect of Donor-Dependent Administration of Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells following Focal Cerebral Ischemia in Rats. <i>Experimental Neurobiology</i> , 2015, 24, 358-365.	1.6	13
15	The Effect of Umbilical Cord Blood Derived Mesenchymal Stem Cells in Monocrotaline-induced Pulmonary Artery Hypertension Rats. <i>Journal of Korean Medical Science</i> , 2015, 30, 576.	2.5	20
16	GDF-15 Secreted from Human Umbilical Cord Blood Mesenchymal Stem Cells Delivered Through the Cerebrospinal Fluid Promotes Hippocampal Neurogenesis and Synaptic Activity in an Alzheimer's Disease Model. <i>Stem Cells and Development</i> , 2015, 24, 2378-2390.	2.1	93
17	Conditioned Media from Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Inhibits Melanogenesis by Promoting Proteasomal Degradation of MITF. <i>PLoS ONE</i> , 2015, 10, e0128078.	2.5	21
18	Effect of preemptive treatment with human umbilical cord blood-derived mesenchymal stem cells on the development of renal ischemia-reperfusion injury in mice. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 307, F1149-F1161.	2.7	36

#	ARTICLE	IF	CITATIONS
19	Low immunogenicity of allogeneic human umbilical cord blood-derived mesenchymal stem cells in vitro and in vivo. <i>Biochemical and Biophysical Research Communications</i> , 2014, 446, 983-989.	2.1	152
20	Transplantation of Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells or Their Conditioned Medium Prevents Bone Loss in Ovariectomized Nude Mice. <i>Tissue Engineering - Part A</i> , 2013, 19, 685-696.	3.1	46
21	Comparative Analysis of Human Mesenchymal Stem Cells from Bone Marrow, Adipose Tissue, and Umbilical Cord Blood as Sources of Cell Therapy. <i>International Journal of Molecular Sciences</i> , 2013, 14, 17986-18001.	4.1	504
22	Thrombospondin-2 secreted by human umbilical cord blood-derived mesenchymal stem cells promotes chondrogenic differentiation. <i>Stem Cells</i> , 2013, 31, 2136-2148.	3.2	102
23	Cryopreserved Cord Blood Progenitors and Their Cell Adhesion Molecules Are Increased by Coculture With Osteoblasts and Parathyroid Hormone. <i>Journal of Pediatric Hematology/Oncology</i> , 2013, 35, e229-e233.	0.6	2
24	Human umbilical cord blood-derived mesenchymal stem cells improve neuropathology and cognitive impairment in an Alzheimer's disease mouse model through modulation of neuroinflammation. <i>Neurobiology of Aging</i> , 2012, 33, 588-602.	3.1	240
25	Human umbilical cord blood-derived mesenchymal stem cell transplantation attenuates severe brain injury by permanent middle cerebral artery occlusion in newborn rats. <i>Pediatric Research</i> , 2012, 72, 277-284.	2.3	112
26	Umbilical cord blood mesenchymal stem cells protect amyloid- β 242 neurotoxicity via paracrine. <i>World Journal of Stem Cells</i> , 2012, 4, 110.	2.8	28
27	Therapeutic effects of human umbilical cord blood-derived mesenchymal stem cells after intrathecal administration by lumbar puncture in a rat model of cerebral ischemia. <i>Stem Cell Research and Therapy</i> , 2011, 2, 38.	5.5	110
28	Galectin-3 Secreted by Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Reduces Amyloid-Beta42 Neurotoxicity In Vitro. <i>Biophysical Journal</i> , 2011, 100, 415a.	0.5	2
29	CXC chemokine receptor 1 enhances the ability of human umbilical cord blood-derived mesenchymal stem cells to migrate toward gliomas. <i>Biochemical and Biophysical Research Communications</i> , 2011, 407, 741-746.	2.1	68
30	Intratracheal Transplantation of Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Dose-Dependently Attenuates Hyperoxia-Induced Lung Injury in Neonatal Rats. <i>Cell Transplantation</i> , 2011, 20, 1843-1854.	2.5	130
31	GD2 expression is closely associated with neuronal differentiation of human umbilical cord blood-derived mesenchymal stem cells. <i>Cellular and Molecular Life Sciences</i> , 2010, 67, 1845-1858.	5.4	47
32	Galectin-3 secreted by human umbilical cord blood-derived mesenchymal stem cells reduces amyloid- β 242 neurotoxicity in vitro. <i>FEBS Letters</i> , 2010, 584, 3601-3608.	2.8	44
33	Microporation is a valuable transfection method for efficient gene delivery into human umbilical cord blood-derived mesenchymal stem cells. <i>BMC Biotechnology</i> , 2010, 10, 38.	3.3	37
34	Irradiation Enhances the Tumor Tropism and Therapeutic Potential of Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Secreting Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells in Glioma Therapy. <i>Stem Cells</i> , 2010, 28, 2217-2228.	3.2	117
35	Application of human umbilical cord blood-derived mesenchymal stem cells in disease models. <i>World Journal of Stem Cells</i> , 2010, 2, 34.	2.8	75
36	Overexpression of CXC Chemokine Receptors Is Required for the Superior Glioma-Tracking Property of Umbilical Cord Blood-Derived Mesenchymal Stem Cells. <i>Stem Cells and Development</i> , 2009, 18, 511-520.	2.1	84

#	ARTICLE	IF	CITATIONS
37	Down-regulation of CD105 is associated with multi-lineage differentiation in human umbilical cord blood-derived mesenchymal stem cells. <i>Biochemical and Biophysical Research Communications</i> , 2009, 381, 676-681.	2.1	90
38	Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Attenuate Hyperoxia-Induced Lung Injury in Neonatal Rats. <i>Cell Transplantation</i> , 2009, 18, 869-886.	2.5	219
39	Brain-derived neurotrophic factor stimulates the neural differentiation of human umbilical cord blood-derived mesenchymal stem cells and survival of differentiated cells through MAPK/ERK and PI3K/Akt-dependent signaling pathways. <i>Journal of Neuroscience Research</i> , 2008, 86, 2168-2178.	2.9	146
40	Immunological properties of umbilical cord blood-derived mesenchymal stromal cells. <i>Cellular Immunology</i> , 2008, 251, 116-123.	3.0	122
41	Gene Therapy Using TRAIL-Secreting Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells against Intracranial Glioma. <i>Cancer Research</i> , 2008, 68, 9614-9623.	0.9	257
42	Efficient Intracytoplasmic Labeling of Human Umbilical Cord Blood Mesenchymal Stromal Cells with Ferumoxides. <i>Cell Transplantation</i> , 2007, 16, 849-857.	2.5	29
43	Effect of Aging on the Pluripotential Capacity of Human Bone Marrow Derived Mesenchymal Stem Cells. <i>The Journal of the Korean Orthopaedic Association</i> , 2007, 42, 701.	0.1	1
44	Mesenchymal stem cells feeder layer from human umbilical cord blood for ex vivo expanded growth and proliferation of hematopoietic progenitor cells. <i>Annals of Hematology</i> , 2006, 85, 212-225.	1.8	107