

Morten Fenger-Grn

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5436840/morten-fenger-gron-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

71
papers

1,836
citations

23
h-index

40
g-index

72
ext. papers

2,307
ext. citations

5.5
avg, IF

4.86
L-index

#	Paper	IF	Citations
71	Long-term risk of dementia among people with traumatic brain injury in Denmark: a population-based observational cohort study. <i>Lancet Psychiatry</i> , 2018, 5, 424-431	23.3	148
70	Effect of depression and diabetes mellitus on the risk for dementia: a national population-based cohort study. <i>JAMA Psychiatry</i> , 2015, 72, 612-9	14.5	111
69	Association between Mental Disorders and Subsequent Medical Conditions. <i>New England Journal of Medicine</i> , 2020, 382, 1721-1731	59.2	103
68	Drug exposure and risk of microscopic colitis: a nationwide Danish case-control study with 5751 cases. <i>Inflammatory Bowel Diseases</i> , 2014, 20, 1702-7	4.5	86
67	The Association Between Perceived Stress and Mortality Among People With Multimorbidity: A Prospective Population-Based Cohort Study. <i>American Journal of Epidemiology</i> , 2016, 184, 199-210	3.8	77
66	Long-term Risk of Dementia in Persons With Schizophrenia: A Danish Population-Based Cohort Study. <i>JAMA Psychiatry</i> , 2015, 72, 1095-101	14.5	74
65	Impact of fish predation on cladoceran body weight distribution and zooplankton grazing in lakes during winter. <i>Freshwater Biology</i> , 2004, 49, 432-447	3.1	71
64	The epidemiology of microscopic colitis: a 10-year pathology-based nationwide Danish cohort study. <i>Scandinavian Journal of Gastroenterology</i> , 2015, 50, 393-8	2.4	68
63	Sick leave and work disability in primary care patients with recent-onset multiple medically unexplained symptoms and persistent somatoform disorders: a 10-year follow-up of the FIP study. <i>General Hospital Psychiatry</i> , 2015, 37, 53-9	5.6	56
62	Small artery structure during antihypertensive therapy is an independent predictor of cardiovascular events in essential hypertension. <i>Journal of Hypertension</i> , 2013, 31, 791-7	1.9	56
61	Body fat, body fat distribution, lean body mass and atrial fibrillation and flutter. A Danish cohort study. <i>Obesity</i> , 2014, 22, 1546-52	8	55
60	Graft patency after off-pump coronary artery bypass surgery is inferior even with identical heparinization protocols: results from the Danish On-pump Versus Off-pump Randomization Study (DOORS). <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2014, 148, 1812-1819.e2	1.5	52
59	Use of general practice, diagnostic investigations and hospital services before and after cancer diagnosis - a population-based nationwide registry study of 127,000 incident adult cancer patients. <i>BMC Health Services Research</i> , 2012, 12, 224	2.9	52
58	Diagnostic intervals before and after implementation of cancer patient pathways - a GP survey and registry based comparison of three cohorts of cancer patients. <i>BMC Cancer</i> , 2015, 15, 308	4.8	48
57	Lean Body Mass Is the Predominant Anthropometric Risk Factor for Atrial Fibrillation. <i>Journal of the American College of Cardiology</i> , 2017, 69, 2488-2497	15.1	45
56	Freestanding midwifery unit versus obstetric unit: a matched cohort study of outcomes in low-risk women. <i>BMJ Open</i> , 2011, 1, e000262	3	44
55	The impact of birthplace on women's birth experiences and perceptions of care. <i>Social Science and Medicine</i> , 2012, 74, 973-81	5.1	43

54	Serious Mental Illness and Risk for Hospitalizations and Rehospitalizations for Ambulatory Care-sensitive Conditions in Denmark: A Nationwide Population-based Cohort Study. <i>Medical Care</i> , 2016 , 54, 90-7	3.1	41
53	Risk of suicide, deliberate self-harm and psychiatric illness after the loss of a close relative: A nationwide cohort study. <i>World Psychiatry</i> , 2017 , 16, 193-199	14.4	30
52	Statins and Risk of Intracerebral Haemorrhage in a Stroke-Free Population: A Nationwide Danish Propensity Score Matched Cohort Study. <i>EClinicalMedicine</i> , 2019 , 8, 78-84	11.3	28
51	Reproducibility of depth of extramural tumor spread and distance to circumferential resection margin at rectal MRI: enhancement of clinical guidelines for neoadjuvant therapy. <i>American Journal of Roentgenology</i> , 2011 , 197, 1360-6	5.4	28
50	Total skin electron beam therapy for cutaneous T-cell lymphoma: a nationwide cohort study from Denmark. <i>Acta Oncologica</i> , 2011 , 50, 1199-205	3.2	25
49	Variation in general practice prostate-specific antigen testing and prostate cancer outcomes: an ecological study. <i>International Journal of Cancer</i> , 2015 , 136, 435-42	7.5	24
48	Closure of secundum atrial septal defects in the adult and elderly patients. <i>European Journal of Cardio-thoracic Surgery</i> , 2013 , 43, 752-7	3	23
47	Use of primary care during the year before childhood cancer diagnosis: a nationwide population-based matched comparative study. <i>PLoS ONE</i> , 2013 , 8, e59098	3.7	23
46	21 Days head-down bed rest induces weakening of cell-mediated immunity - Some spaceflight findings confirmed in a ground-based analog. <i>Cytokine</i> , 2012 , 59, 403-9	4	21
45	Subsequent cancers, mortality, and causes of death in patients with mycosis fungoides and parapsoriasis: a Danish nationwide, population-based cohort study. <i>Journal of the American Academy of Dermatology</i> , 2014 , 71, 529-35	4.5	19
44	The role of general practice in routes to diagnosis of lung cancer in Denmark: a population-based study of general practice involvement, diagnostic activity and diagnostic intervals. <i>BMC Health Services Research</i> , 2015 , 15, 21	2.9	18
43	Long-term risk of atrial fibrillation after the death of a partner. <i>Open Heart</i> , 2016 , 3, e000367	3	18
42	Atrial Fibrillation and Risk of Cancer: A Danish Population-Based Cohort Study. <i>Journal of the American Heart Association</i> , 2018 , 7, e009543	6	18
41	Correlation of mortality with lesion level in patients with myelomeningocele: a population-based study. <i>Journal of Neurosurgery: Pediatrics</i> , 2017 , 19, 227-231	2.1	17
40	Statins and Risk of Intracerebral Hemorrhage in Individuals With a History of Stroke. <i>Stroke</i> , 2020 , 51, 1111-1119	6.7	17
39	Low breastfeeding rates and body mass index in Danish children of women with gestational diabetes mellitus. <i>International Breastfeeding Journal</i> , 2015 , 10, 26	3.8	17
38	Depression and risk of hospitalisations and rehospitalisations for ambulatory care-sensitive conditions in Denmark: a population-based cohort study. <i>BMJ Open</i> , 2015 , 5, e009878	3	16
37	Depression, antidepressants, and the risk of non-valvular atrial fibrillation: A nationwide Danish matched cohort study. <i>European Journal of Preventive Cardiology</i> , 2019 , 26, 187-195	3.9	16

36	The effect of an active implementation of a disease management programme for chronic obstructive pulmonary disease on healthcare utilization--a cluster-randomised controlled trial. <i>BMC Health Services Research</i> , 2013 , 13, 385	2.9	15
35	Decreased incidence of myelomeningocele at birth: effect of folic acid recommendations or prenatal diagnostics?. <i>Childs Nervous System</i> , 2011 , 27, 1951-5	1.7	15
34	Association between perceived stress, multimorbidity and primary care health services: a Danish population-based cohort study. <i>BMJ Open</i> , 2018 , 8, e018323	3	14
33	Freestanding midwifery units versus obstetric units: does the effect of place of birth differ with level of social disadvantage?. <i>BMC Public Health</i> , 2012 , 12, 478	4.1	14
32	The effect of direct access to CT scan in early lung cancer detection: an unblinded, cluster-randomised trial. <i>BMC Cancer</i> , 2015 , 15, 934	4.8	12
31	OPCAB surgery is cost-effective for elderly patients. <i>Scandinavian Cardiovascular Journal</i> , 2013 , 47, 185-92		12
30	Caffeine Supplementation Had No Effect on Endurance Capacity in Elderly Subjects Who Had Abstained from Caffeine-Containing Nutrition for 8 Hours. <i>Journal of Caffeine Research</i> , 2011 , 1, 109-116		11
29	Primary Care Use before Cancer Diagnosis in Adolescents and Young Adults - A Nationwide Register Study. <i>PLoS ONE</i> , 2016 , 11, e0155933	3.7	10
28	Tumour stage and implementation of standardised cancer patient pathways: a comparative cohort study. <i>British Journal of General Practice</i> , 2016 , 66, e434-43	1.6	9
27	Patient-experienced effect of an active implementation of a disease management programme for COPD - a randomised trial. <i>BMC Family Practice</i> , 2013 , 14, 147	2.6	9
26	Algorithm linking patients and general practices in Denmark using the Danish National Health Service Register. <i>Clinical Epidemiology</i> , 2016 , 8, 273-83	5.9	9
25	Does perceived stress increase the risk of atrial fibrillation? A population-based cohort study in Denmark. <i>American Heart Journal</i> , 2017 , 188, 26-34	4.9	8
24	The association of unipolar depression with thirty-day mortality after hospitalization for infection: A population-based cohort study in Denmark. <i>Journal of Psychosomatic Research</i> , 2016 , 89, 32-8	4.1	8
23	Frequency of health care utilization in the year prior to completed suicide: A Danish nationwide matched comparative study. <i>PLoS ONE</i> , 2019 , 14, e0214605	3.7	7
22	Use of primary health care prior to a postpartum psychiatric episode. <i>Scandinavian Journal of Primary Health Care</i> , 2015 , 33, 127-33	2.7	7
21	Perceived Stress, Multimorbidity, and Risk for Hospitalizations for Ambulatory Care-sensitive Conditions: A Population-based Cohort Study. <i>Medical Care</i> , 2017 , 55, 131-139	3.1	7
20	Healthcare Contacts after Myocardial Infarction According to Mental Health and Socioeconomic Position: A Population-Based Cohort Study. <i>PLoS ONE</i> , 2015 , 10, e0134557	3.7	7
19	Socioeconomic position and contact to general practice among persons with diabetes. <i>Primary Care Diabetes</i> , 2012 , 6, 313-8	2.4	7

18	A cluster-randomised, parallel group, controlled intervention study of genetic prostate cancer risk assessment and use of PSA tests in general practice--the ProCaRis study: study protocol. <i>BMJ Open</i> , 2013 , 3,	3	7
17	Cancer Mortality in People Treated with Antidepressants before Cancer Diagnosis: A Population Based Cohort Study. <i>PLoS ONE</i> , 2015 , 10, e0138134	3.7	7
16	Body mass and atrial fibrillation risk: Status of the epidemiology concerning the influence of fat versus lean body mass. <i>Trends in Cardiovascular Medicine</i> , 2020 , 30, 205-211	6.9	7
15	Early treatment with talk therapy or antidepressants in severely bereaved people and risk of suicidal behavior and psychiatric illness: an instrumental variable analysis. <i>Clinical Epidemiology</i> , 2018 , 10, 1013-1026	5.9	6
14	Role for machine learning in sex-specific prediction of successful electrical cardioversion in atrial fibrillation?. <i>Open Heart</i> , 2020 , 7,	3	6
13	Selective serotonin reuptake inhibitors and risk of epilepsy after traumatic brain injury - A population based cohort study. <i>PLoS ONE</i> , 2019 , 14, e0219137	3.7	5
12	Use of hyperlinks in electronic test result communication: a survey study in general practice. <i>BMC Medical Informatics and Decision Making</i> , 2012 , 12, 114	3.6	5
11	Long-term subjective results of tension-free vaginal tape operation for female urinary stress incontinence. <i>International Urogynecology Journal</i> , 2012 , 23, 585-8	2	5
10	Depression and Uptake of Oral Anticoagulation Therapy in Patients With Atrial Fibrillation: A Danish Nationwide Cohort Study. <i>Medical Care</i> , 2020 , 58, 216-224	3.1	5
9	Inequality in oral anticoagulation use and clinical outcomes in atrial fibrillation: a Danish nationwide perspective. <i>European Heart Journal Quality of Care & Clinical Outcomes</i> , 2018 , 4, 189-199	4.6	4
8	Association Between Bipolar Disorder or Schizophrenia and Oral Anticoagulation Use in Danish Adults With Incident or Prevalent Atrial Fibrillation. <i>JAMA Network Open</i> , 2021 , 4, e2110096	10.4	3
7	Health care utilization in general practice after HPV vaccination-A Danish nationwide register-based cohort study. <i>PLoS ONE</i> , 2017 , 12, e0184658	3.7	2
6	Classification Error of the Thresholded Independence Rule. <i>Scandinavian Journal of Statistics</i> , 2015 , 42, 32-42	0.8	1
5	A Randomised Trial Examining Cardiovascular Morbidity and All-Cause Mortality 24 years Following General Health Checks: the Ebeltoft Health Promotion Project (EHPP). <i>BMJ Open</i> , 2019 , 9, e030400	3	1
4	Death of a Partner and Risks of Ischemic Stroke and Intracerebral Hemorrhage: A Nationwide Danish Matched Cohort Study. <i>Journal of the American Heart Association</i> , 2020 , 9, e018763	6	1
3	Potentially inappropriate medications (PIMs): frequency and extent of GP-related variation in PIMs: a register-based cohort study. <i>BMJ Open</i> , 2021 , 11, e046756	3	1
2	Reply to the comment to the paper (CNS DOI 10.1007/s00381-010-1113-2) A retrospective study of infections after primary VP shunt placement in the newborn with myelomeningocele without prophylactic antibiotics by Farid Radmanesh. MD, MPH. <i>Childs Nervous System</i> , 2011 , 27, 343-344	1.7	
1	Traumatic brain injury and dementia - Authors' reply. <i>Lancet Psychiatry</i> , 2018 , 5, 783	23.3	

