

Satoru Joshita

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5441803/publications.pdf>

Version: 2024-02-01

107
papers

3,385
citations

159358
30
h-index

168136
53
g-index

107
all docs

107
docs citations

107
times ranked

4869
citing authors

#	ARTICLE	IF	CITATIONS
1	Astrocyte-derived interleukin-33 promotes microglial synapse engulfment and neural circuit development. <i>Science</i> , 2018, 359, 1269-1273.	6.0	422
2	Genome-wide Association Study Identifies TNFSF15 and POU2AF1 as Susceptibility Loci for Primary Biliary Cirrhosis in the Japanese Population. <i>American Journal of Human Genetics</i> , 2012, 91, 721-728.	2.6	251
3	Humanized TREM2 mice reveal microglia-intrinsic and -extrinsic effects of R47H polymorphism. <i>Journal of Experimental Medicine</i> , 2018, 215, 745-760.	4.2	182
4	Pretreatment prediction of virological response to peginterferon plus ribavirin therapy in chronic hepatitis C patients using viral and host factors. <i>Hepatology</i> , 2008, 48, 1753-1760.	3.6	129
5	Serum Wisteria floribunda Agglutinin-Positive Mac-2-Binding Protein Level Predicts Liver Fibrosis and Prognosis in Primary Biliary Cirrhosis. <i>American Journal of Gastroenterology</i> , 2015, 110, 857-864.	0.2	115
6	Clinical significance of immunoglobulin G4-associated autoimmune hepatitis. <i>Journal of Gastroenterology</i> , 2011, 46, 48-55.	2.3	113
7	Long-term outcome of Japanese patients with type 1 autoimmune hepatitis. <i>Hepatology</i> , 2012, 56, 668-676.	3.6	95
8	Down-regulation of SREBP-1c is associated with the development of burned-out NASH. <i>Journal of Hepatology</i> , 2010, 53, 724-731.	1.8	89
9	Bezafibrate Improves GLOBE and UKâ€PBC Scores and Longâ€Term Outcomes in Patients With Primary Biliary Cholangitis. <i>Hepatology</i> , 2019, 70, 2035-2046.	3.6	83
10	Human leukocyte antigen class II molecules confer both susceptibility and progression in Japanese patients with primary biliary cirrhosis. <i>Hepatology</i> , 2012, 55, 506-511.	3.6	73
11	Serum Fragmented Cytokeratin 18 Levels Reflect the Histologic Activity Score of Nonalcoholic Fatty Liver Disease More Accurately Than Serum Alanine Aminotransferase Levels. <i>Journal of Clinical Gastroenterology</i> , 2010, 44, 440-447.	1.1	68
12	An international genome-wide meta-analysis of primary biliary cholangitis: Novel risk loci and candidate drugs. <i>Journal of Hepatology</i> , 2021, 75, 572-581.	1.8	62
13	Steatogenesis in adult-onset type II citrullinemia is associated with down-regulation of PPARÎ±. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 473-481.	1.8	57
14	Human Leukocyte Antigen Class II Haplotypes Affect Clinical Characteristics and Progression of Type 1 Autoimmune Hepatitis in Japan. <i>PLoS ONE</i> , 2014, 9, e100565.	1.1	54
15	Mild drinking habit is a risk factor for hepatocarcinogenesis in non-alcoholic fatty liver disease with advanced fibrosis. <i>World Journal of Gastroenterology</i> , 2018, 24, 1440-1450.	1.4	54
16	Granulocyte-Colony Stimulating Factor-Producing Pancreatic Adenosquamous Carcinoma Showing Aggressive Clinical Course. <i>Internal Medicine</i> , 2009, 48, 687-691.	0.3	51
17	Association of Serum Cytokine Levels With Treatment Response to Pegylated Interferon and Ribavirin Therapy in Genotype 1 Chronic Hepatitis C Patients. <i>Journal of Infectious Diseases</i> , 2011, 203, 1087-1095.	1.9	50
18	Association of Serum Autotaxin Levels with Liver Fibrosis in Patients with Chronic Hepatitis C. <i>Scientific Reports</i> , 2017, 7, 46705.	1.6	49

#	ARTICLE	IF	CITATIONS
19	Genome-wide association studies identify PRKCB as a novel genetic susceptibility locus for primary biliary cholangitis in the Japanese population. <i>Human Molecular Genetics</i> , 2017, 26, ddw406.	1.4	46
20	Serum <i>Wisteria floribunda</i> agglutinin-2 positive human Mac-2 binding protein may predict liver fibrosis and progression to hepatocellular carcinoma in patients with chronic hepatitis B virus infection. <i>Hepatology Research</i> , 2017, 47, 226-233.	1.8	42
21	Serum autotaxin levels are correlated with hepatic fibrosis and ballooning in patients with non-alcoholic fatty liver disease. <i>World Journal of Gastroenterology</i> , 2018, 24, 1239-1249.	1.4	39
22	Association analysis of cytotoxic T-lymphocyte antigen 4 gene polymorphisms with primary biliary cirrhosis in Japanese patients. <i>Journal of Hepatology</i> , 2010, 53, 537-541.	1.8	38
23	Association of IL28B gene polymorphism with development of hepatocellular carcinoma in Japanese patients with chronic hepatitis C virus infection. <i>Human Immunology</i> , 2012, 73, 298-300.	1.2	38
24	Genetics and epigenetics in the pathogenesis of primary biliary cholangitis. <i>Clinical Journal of Gastroenterology</i> , 2018, 11, 11-18.	0.4	36
25	Association between endotoxemia and histological features of nonalcoholic fatty liver disease. <i>World Journal of Gastroenterology</i> , 2017, 23, 712.	1.4	36
26	Clinical features of autoimmune hepatitis with acute presentation: a Japanese nationwide survey. <i>Journal of Gastroenterology</i> , 2018, 53, 1079-1088.	2.3	35
27	Incidence and prevalence of autoimmune hepatitis in the Ueda area, Japan. <i>Hepatology Research</i> , 2016, 46, 878-883.	1.8	34
28	Serum autotaxin is a useful liver fibrosis marker in patients with chronic hepatitis B virus infection. <i>Hepatology Research</i> , 2018, 48, 275-285.	1.8	34
29	Accurate and simple method for quantification of hepatic fat content using magnetic resonance imaging: a prospective study in biopsy-proven nonalcoholic fatty liver disease. <i>Journal of Gastroenterology</i> , 2010, 45, 1263-1271.	2.3	32
30	Genetic polymorphisms in CTLA4 and SLC4A2 are differentially associated with the pathogenesis of primary biliary cirrhosis in Japanese patients. <i>Journal of Gastroenterology</i> , 2011, 46, 1203-1212.	2.3	32
31	AST/platelet ratio index associates with progression to hepatic failure and correlates with histological fibrosis stage in Japanese patients with primary biliary cirrhosis. <i>Journal of Hepatology</i> , 2014, 61, 1443-1445.	1.8	32
32	Serum Autotaxin Is a Useful Disease Progression Marker in Patients with Primary Biliary Cholangitis. <i>Scientific Reports</i> , 2018, 8, 8159.	1.6	32
33	Serum interleukin (IL)-10 and IL-12 levels and IL28B gene polymorphisms: pretreatment prediction of treatment failure in chronic hepatitis C. <i>Antiviral Therapy</i> , 2011, 16, 1073-1080.	0.6	30
34	Clinicopathological characteristics of non-B non-C hepatocellular carcinoma without past hepatitis B virus infection. <i>Hepatology Research</i> , 2017, 47, 405-418.	1.8	30
35	A Patient with Advanced Hepatocellular Carcinoma Treated with Sorafenib Tosylate Showed Massive Tumor Lysis with Avoidance of Tumor Lysis syndrome. <i>Internal Medicine</i> , 2010, 49, 991-994.	0.3	29
36	Development from simple steatosis to liver cirrhosis and hepatocellular carcinoma: a 27-year follow-up case. <i>Clinical Journal of Gastroenterology</i> , 2008, 1, 116-121.	0.4	27

#	ARTICLE	IF	CITATIONS
37	Genetic Association of PTPN22 Polymorphisms with Autoimmune Hepatitis and Primary Biliary Cholangitis in Japan. <i>Scientific Reports</i> , 2016, 6, 29770.	1.6	27
38	Serum thrombospondin 2 is a novel predictor for the severity in the patients with NAFLD. <i>Liver International</i> , 2021, 41, 505-514.	1.9	25
39	A Case of Granulocyte-Colony Stimulating Factor-Producing Hepatocellular Carcinoma Confirmed by Immunohistochemistry. <i>Journal of Korean Medical Science</i> , 2010, 25, 476.	1.1	24
40	Serum Cell Death Biomarkers for Prediction of Liver Fibrosis and Poor Prognosis in Primary Biliary Cirrhosis. <i>PLoS ONE</i> , 2015, 10, e0131658.	1.1	24
41	Biochemical and plasma lipid responses to pemafibrate in patients with primary biliary cholangitis. <i>Hepatology Research</i> , 2019, 49, 1236-1243.	1.8	24
42	POGLUT1, the putative effector gene driven by rs2293370 in primary biliary cholangitis susceptibility locus chromosome 3q13.33. <i>Scientific Reports</i> , 2019, 9, 102.	1.6	23
43	Association between serum soluble CD14 and IL8 levels and clinical outcome in primary biliary cholangitis. <i>Liver International</i> , 2017, 37, 897-905.	1.9	22
44	A cis-eQTL of HLA-DPB1 Affects Susceptibility to Type 1 Autoimmune Hepatitis. <i>Scientific Reports</i> , 2018, 8, 11924.	1.6	22
45	Association between KIR-HLA combination and ulcerative colitis and Crohn's disease in a Japanese population. <i>PLoS ONE</i> , 2018, 13, e0195778.	1.1	21
46	Past history of hepatocellular carcinoma is an independent risk factor of treatment failure in patients with chronic hepatitis C virus infection receiving direct-acting antivirals. <i>Journal of Viral Hepatitis</i> , 2018, 25, 1462-1471.	1.0	21
47	A case of well-differentiated cholangiolocellular carcinoma visualized with contrast-enhanced ultrasonography using Sonazoid. <i>Hepatology Research</i> , 2009, 39, 207-212.	1.8	20
48	A Case of Pyogenic Liver Abscess Infected with <i>Fusobacterium necrophorum</i> Depicted by Microscopy and Confirmed by Tissue Culture. <i>Internal Medicine</i> , 2011, 50, 1815-1819.	0.3	20
49	Serum levels of interleukin-22 and hepatitis B core-related antigen are associated with treatment response to entecavir therapy in chronic hepatitis B. <i>Hepatology Research</i> , 2014, 44, E172-80.	1.8	20
50	Association of autoimmune hepatitis with Src homology 2 adaptor protein 3 gene polymorphisms in Japanese patients. <i>Journal of Human Genetics</i> , 2017, 62, 963-967.	1.1	20
51	Cytokine profiles affecting the pathogenesis of autoimmune hepatitis in Japanese patients. <i>Hepatology Research</i> , 2011, 41, 350-357.	1.8	19
52	A2BP1 as a novel susceptible gene for primary biliary cirrhosis in Japanese patients. <i>Human Immunology</i> , 2010, 71, 520-524.	1.2	18
53	Serum chemokine levels are associated with the outcome of pegylated interferon and ribavirin therapy in patients with chronic hepatitis C. <i>Hepatology Research</i> , 2011, 41, 587-593.	1.8	18
54	Genetic Contribution to the Pathogenesis of Primary Biliary Cholangitis. <i>Journal of Immunology Research</i> , 2017, 2017, 1-6.	0.9	17

#	ARTICLE	IF	CITATIONS
55	Clinical utility of FibroScan as a noninvasive diagnostic test for primary biliary cholangitis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2020, 35, 1208-1214.	1.4	17
56	KIR, HLA, and IL28B Variant Predict Response to Antiviral Therapy in Genotype 1 Chronic Hepatitis C Patients in Japan. <i>PLoS ONE</i> , 2013, 8, e83381.	1.1	17
57	Liver Dysfunction and Thrombocytopenia Diagnosed as Intravascular Large B-cell Lymphoma Using a Timely and Accurate Transjugular Liver Biopsy. <i>Internal Medicine</i> , 2013, 52, 1903-1908.	0.3	16
58	KIR/HLA genotypes confer susceptibility and progression in patients with autoimmune hepatitis. <i>JHEP Reports</i> , 2019, 1, 353-360.	2.6	16
59	STAT4 Gene Polymorphisms Are Associated with Susceptibility and ANA Status in Primary Biliary Cirrhosis. <i>Disease Markers</i> , 2014, 2014, 1-8.	0.6	15
60	Liver stiffness-spleen size-to-platelet ratio risk score detects esophageal varices in chronic liver disease. <i>SpringerPlus</i> , 2016, 5, 998.	1.2	15
61	aMAP score prediction of hepatocellular carcinoma occurrence and incidence-free rate after a sustained virologic response in chronic hepatitis C. <i>Hepatology Research</i> , 2021, 51, 933-942.	1.8	15
62	Efficacy and safety of eradication therapy for elderly patients with helicobacter pylori infection. <i>Medicine (United States)</i> , 2019, 98, e16619.	0.4	14
63	A validation study of the Ursodeoxycholic Acid Response Score in Japanese patients with primary biliary cholangitis. <i>Liver International</i> , 2020, 40, 1926-1933.	1.9	14
64	Miglitol attenuates nonalcoholic steatohepatitis in diabetic patients. <i>Hepatology Research</i> , 2018, 48, 1092-1098.	1.8	13
65	Changes in serum levels of autotaxin with direct-acting antiviral therapy in patients with chronic hepatitis C. <i>PLoS ONE</i> , 2018, 13, e0195632.	1.1	13
66	Liver stiffness-spleen size-to-platelet ratio risk score identifies esophageal varices in Japanese patients with chronic hepatitis C. <i>Hepatology Research</i> , 2016, 46, 884-889.	1.8	12
67	Changes in the serum level of hepatitis B virus (HBV) surface antigen over the natural course of HBV infection. <i>Journal of Gastroenterology</i> , 2012, 47, 1006-1013.	2.3	11
68	KIR3DL1-HLA-Bw4 combination and IL28B polymorphism predict response to Peg-IFN and ribavirin with and without telaprevir in chronic hepatitis C. <i>Human Immunology</i> , 2014, 75, 822-826.	1.2	11
69	Effectiveness of Glecaprevir/Pibrentasvir for Hepatitis C: Real-World Experience and Clinical Features of Retreatment Cases. <i>Biomedicines</i> , 2020, 8, 74.	1.4	11
70	KIR2DL2 combined with HLA-C1 confers risk of hepatitis C virus-related hepatocellular carcinoma in younger patients. <i>Oncotarget</i> , 2018, 9, 19650-19661.	0.8	11
71	Cutaneous sarcoidosis in a chronic hepatitis C patient receiving pegylated interferon and ribavirin therapy. <i>Hepatology Research</i> , 2013, 43, 801-807.	1.8	10
72	Genetic polymorphism in IFNL4 and response to pegylated interferon and ribavirin in Japanese chronic hepatitis C patients. <i>Tissue Antigens</i> , 2014, 83, 45-48.	1.0	10

#	ARTICLE	IF	CITATIONS
73	<i>Lymphocyte Antigen 75</i> Polymorphisms Are Associated with Disease Susceptibility and Phenotype in Japanese Patients with Inflammatory Bowel Disease. <i>Disease Markers</i> , 2016, 2016, 1-7.	0.6	10
74	Cimetidine/lactulose therapy ameliorates erythropoietic protoporphyria-related liver injury. <i>Clinical Journal of Gastroenterology</i> , 2017, 10, 452-458.	0.4	9
75	Renal impairment is associated with increased risk of mortality in patients with cirrhosis. <i>Medicine (United States)</i> , 2019, 98, e14475.	0.4	9
76	2â€Step PLT16â€AST44 method: Simplified liver fibrosis detection system in patients with nonâ€alcoholic fatty liver disease. <i>Hepatology Research</i> , 2022, 52, 352-363.	1.8	8
77	Association between lower air pressure and the onset of ischemic colitis: a caseâ€control study. <i>European Journal of Gastroenterology and Hepatology</i> , 2017, 29, 1071-1078.	0.8	7
78	Long-term luseogliflozin therapy improves histological activity of non-alcoholic steatohepatitis accompanied by type 2 diabetes mellitus. <i>Clinical Journal of Gastroenterology</i> , 2020, 13, 83-89.	0.4	7
79	The ursodeoxycholic acid response score predicts pathological features in primary biliary cholangitis. <i>Hepatology Research</i> , 2021, 51, 80-89.	1.8	7
80	Clinical practice guidelines for autoimmune hepatitis. <i>Hepatology Research</i> , 2022, 52, 571-585.	1.8	7
81	Association analysis of toll-like receptor 4 polymorphisms in Japanese primary biliary cirrhosis. <i>Human Immunology</i> , 2013, 74, 219-222.	1.2	6
82	Investigation of the Effect of KIRâ€HLA Pairs on Hepatocellular Carcinoma in Hepatitis C Virus Cirrhotic Patients. <i>Cancers</i> , 2021, 13, 3267.	1.7	6
83	Salmonella Enteritidis cholecystitis with chronic granulomatous disease. <i>IDCases</i> , 2018, 12, 49-52.	0.4	5
84	Virological Factors Associated with the Occurrence of HBV Reactivation in Patients with Resolved HBV Infection Analyzed through Ultradeep Sequencing. <i>Journal of Infectious Diseases</i> , 2019, 221, 400-407.	1.9	5
85	A case of liver abscess co-infected with <i>Desulfovibrio desulfuricans</i> and <i>Escherichia coli</i> and review of the literature. <i>Journal of Infection and Chemotherapy</i> , 2018, 24, 393-397.	0.8	4
86	Quantitative analysis of serum chemokines associated with treatment failure of direct-acting antivirals in chronic hepatitis C. <i>Cytokine</i> , 2018, 111, 357-363.	1.4	4
87	A Case of Adult T-Cell Leukemia/Lymphoma Complicated with Bilateral Chylothorax. <i>Case Reports in Oncological Medicine</i> , 2019, 2019, 1-5.	0.2	4
88	Association analysis of KIR/HLA genotype with liver cirrhosis, hepatocellular carcinoma, and NUC freedom in chronic hepatitis B patients. <i>Scientific Reports</i> , 2021, 11, 21424.	1.6	4
89	Zygomycosis Presenting as Acute Myocardial Infarction during Hematological Malignancies. <i>Internal Medicine</i> , 2008, 47, 839-842.	0.3	3
90	Early detection of interstitial pneumonia by monitoring KLâ€6 in a chronic hepatitis C patient undergoing pegylated interferon and ribavirin therapy. <i>Hepatology Research</i> , 2011, 41, 904-909.	1.8	3

#	ARTICLE	IF	CITATIONS
91	Laparoscopic findings of congenital hepatic fibrosis: A case report and review of the published work. <i>Hepatology Research</i> , 2014, 44, 818-824.	1.8	3
92	A case report of pancreatic panniculitis due to acute pancreatitis with intraductal papillary mucinous neoplasm. <i>BMC Gastroenterology</i> , 2020, 20, 286.	0.8	3
93	Clinical impact of normal alanine aminotransferase on direct-acting antiviral outcome in patients with chronic hepatitis C virus infection. <i>JGH Open</i> , 2020, 4, 574-581.	0.7	3
94	Primary Hepatic Extranodal Marginal Zone Lymphoma of Mucosa-Associated Lymphoid Tissue in a Patient with Chronic Hepatitis B Virus Infection: Case Report and Summary of the Literature. <i>Medicina (Lithuania)</i> , 2021, 57, 280.	0.8	3
95	Quantitative and qualitative lipid improvement with chronic hepatitis C virus eradication using direct-acting antivirals. <i>Hepatology Research</i> , 2021, 51, 758-766.	1.8	3
96	Risk factors for relapse of autoimmune hepatitis in Japan: A nationwide survey. <i>Hepatology Research</i> , 2022, 52, 597-602.	1.8	3
97	Characteristics and prediction of hepatitis B e-antigen negative hepatitis following seroconversion in patients with chronic hepatitis B. <i>Hepatology Research</i> , 2014, 44, E45-53.	1.8	2
98	Emergence of anti-mitochondrial M2 antibody in patient with angioimmunoblastic T-cell lymphoma. <i>Clinical Journal of Gastroenterology</i> , 2018, 11, 302-308.	0.4	2
99	Improvement of porphyria cutanea tarda following treatment of hepatitis C virus by direct-acting antivirals: A case report. <i>Journal of Dermatology</i> , 2019, 46, e149-e151.	0.6	2
100	Protocol: Prospective observational study aiming for micro-elimination of hepatitis C virus in Nagawa town: The Nagawa Project. <i>PLoS ONE</i> , 2021, 16, e0256711.	1.1	2
101	Editorial: FAST score—a new predictive marker for HCC after SVR. <i>Alimentary Pharmacology and Therapeutics</i> , 2020, 52, 1222-1223.	1.9	2
102	Primitive Neuroectodermal Tumor as a Differential Diagnosis of CD56-Positive Tumors in Adults. <i>Internal Medicine</i> , 2009, 48, 1267-1272.	0.3	1
103	Polymorphism at rs9264942 is associated with HLA-C expression and inflammatory bowel disease in the Japanese. <i>Scientific Reports</i> , 2020, 10, 12424.	1.6	1
104	Protocol: Prospective observational study investigating the prevalence and clinical outcome of portopulmonary hypertension in Japanese patients with chronic liver disease. <i>PLoS ONE</i> , 2021, 16, e0249435.	1.1	1
105	An Autopsy Case of Primary Biliary Cholangitis with Histological Submassive Hepatic Necrosis Caused by Acute Hepatitis E Virus Infection. <i>Internal Medicine</i> , 2021, 60, 1863-1870.	0.3	1
106	A Patency Capsule Remained Intact in the Colon over 210 Hours. <i>Case Reports in Gastrointestinal Medicine</i> , 2017, 2017, 1-3.	0.2	0
107	The levels of IL-1 β and soluble IL-1 receptors in patients with IgG4-related periaortitis/periarteritis. <i>Advances in Medical Sciences</i> , 2022, 67, 257-261.	0.9	0