Eiichi Tanaka

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5443205/publications.pdf

Version: 2024-02-01

19	518	9	18
papers	citations	h-index	g-index
19	19	19	510 citing authors
all docs	docs citations	times ranked	

#	Article	IF	CITATIONS
1	Drug treatment algorithm and recommendations from the 2020 update of the Japan College of Rheumatology clinical practice guidelines for the management of rheumatoid arthritis—secondary publication. Modern Rheumatology, 2023, 33, 21-35.	1.8	35
2	Systematic review and meta-analysis of biosimilar for the treatment of rheumatoid arthritis informing the 2020 update of the Japan College of Rheumatology clinical practice guidelines for the management of rheumatoid arthritis. Modern Rheumatology, 2022, 32, 74-86.	1.8	11
3	Factors associated with frailty in Japanese patients with rheumatoid arthritis: results from the Institute of Rheumatology Rheumatoid Arthritis cohort study. Clinical Rheumatology, 2022, 41, 405-410.	2.2	8
4	Impact of concomitant chronic kidney disease on hospitalised infections and remission in patients with rheumatoid arthritis: results from the IORRA cohort. Modern Rheumatology, 2022, 32, 875-884.	1.8	1
5	A simple screening test to assess risk of falls in Japanese patients with rheumatoid arthritis: Results from the IORRA cohort study. Modern Rheumatology, 2021, 31, 506-509.	1.8	2
6	Adherence to denosumab for the treatment of osteoporosis in Japanese patients with rheumatoid arthritis: Results from the IORRA cohort study. Modern Rheumatology, 2021, 31, 510-513.	1.8	2
7	Polypharmacy, declined walking speed, bent back, and disability associated with a history of falls in Japanese patients with rheumatoid arthritis: Results from the IORRA cohort study. Modern Rheumatology, 2021, 31, 764-767.	1.8	5
8	Vitamin D deficiency is a risk factor for new fractures in Japanese postmenopausal women with rheumatoid arthritis: results from the IORRA cohort study. Archives of Osteoporosis, 2021, 16, 119.	2.4	8
9	A large observational cohort study of rheumatoid arthritis, IORRA: Providing context for today's treatment options. Modern Rheumatology, 2020, 30, 1-6.	1.8	83
10	Sustained discontinuation of infliximab with a raising-dose strategy after obtaining remission in patients with rheumatoid arthritis: the RRRR study, a randomised controlled trial. Annals of the Rheumatic Diseases, 2020, 79, 94-102.	0.9	13
11	A retrospective analysis of the relationship between anti-cyclic citrullinated peptide antibody and the effectiveness of abatacept in rheumatoid arthritis patients. Scientific Reports, 2020, 10, 19717.	3.3	3
12	Factors associated with osteoporosis medication use in Japanese patients with rheumatoid arthritis: Results from the Institute of Rheumatology Rheumatoid Arthritis cohort study. Osteoporosis and Sarcopenia, 2020, 6, 82-87.	1.9	6
13	Factors associated with decreasing serum 25(OH)D among Japanese patients with rheumatoid arthritis: Results from the IORRA cohort study. Modern Rheumatology, 2019, 29, 430-435.	1.8	7
14	2017 Clinical practice guidelines of the Japan Research Committee of the Ministry of Health, Labour, and Welfare for Intractable Vasculitis for the management of ANCA-associated vasculitis. Modern Rheumatology, 2019, 29, 20-30.	1.8	47
15	Risk of opportunistic infections in patients with antineutrophil cytoplasmic antibodyâ€associated vasculitis, using a Japanese health insurance database. International Journal of Rheumatic Diseases, 2019, 22, 1978-1984.	1.9	16
16	Prevalence of and factors associated with vitamin D deficiency in 4,793 Japanese patients with rheumatoid arthritis. Clinical Rheumatology, 2013, 32, 1081-1087.	2.2	38
17	Morphology of the epithelium of the lower rectum and the anal canal in the adult human. Medical Molecular Morphology, 2012, 45, 72-79.	1.0	21
18	Intracavitary brachytherapy for carcinoma of the uterine cervixâ€"Comparison of HDR (Ir-192) and MDR (Cs-137)â€". Radiation Medicine, 2006, 24, 50-57.	0.8	6

#	Article	IF	CITATIONS
19	Validation of a Japanese version of the Stanford Health Assessment Questionnaire in 3,763 patients with rheumatoid arthritis. Arthritis and Rheumatism, 2003, 49, 784-788.	6.7	206