Junichi Matsumoto

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5450005/publications.pdf

Version: 2024-02-01

10	304	9	10
papers	citations	h-index	g-index
10	10	10	462
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Dipeptidyl peptidase-4 inhibitor improved exercise capacity and mitochondrial biogenesis in mice with heart failure via activation of glucagon-like peptide-1 receptor signalling. Cardiovascular Research, 2016, 111, 338-347.	3.8	64
2	Protein acetylation in skeletal muscle mitochondria is involved in impaired fatty acid oxidation and exercise intolerance in heart failure. Journal of Cachexia, Sarcopenia and Muscle, 2018, 9, 844-859.	7.3	46
3	Empagliflozin restores lowered exercise endurance capacity via the activation of skeletal muscle fatty acid oxidation in a murine model of heart failure. European Journal of Pharmacology, 2020, 866, 172810.	3.5	43
4	The experimental model of transition from compensated cardiac hypertrophy to failure created by transverse aortic constriction in mice. IJC Heart and Vasculature, 2016, 11, 24-28.	1.1	33
5	Brain-Derived Neurotrophic Factor Improves Limited Exercise Capacity in Mice With Heart Failure. Circulation, 2018, 138, 2064-2066.	1.6	32
6	Inhibition of xanthine oxidase in the acute phase of myocardial infarction prevents skeletal muscle abnormalities and exercise intolerance. Cardiovascular Research, 2021, 117, 805-819.	3.8	25
7	Cardiac-specific loss of mitoNEET expression is linked with age-related heart failure. Communications Biology, 2021, 4, 138.	4.4	20
8	Brain-Derived Neurotrophic Factor Improves Impaired Fatty Acid Oxidation Via the Activation of Adenosine Monophosphate-Activated Protein Kinase-ɑ – Proliferator-Activated Receptor-r Coactivator-1ɑ Signaling in Skeletal Muscle of Mice With Heart Failure. Circulation: Heart Failure, 2021, 14, e005890.	3.9	18
9	Deletion of NAD(P)H Oxidase 2 Prevents Angiotensin II-Induced Skeletal Muscle Atrophy. BioMed Research International, 2018, 2018, 1-10.	1.9	13
10	Angiotensin-converting-enzyme inhibitor prevents skeletal muscle fibrosis in myocardial infarction mice. Skeletal Muscle, 2020, 10, 11.	4.2	10