# Tine Rask Licht

#### List of Publications by Citations

Source: https://exaly.com/author-pdf/5458612/tine-rask-licht-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

101<br/>papers5,380<br/>citations42<br/>h-index72<br/>g-index104<br/>ext. papers6,852<br/>ext. citations7.2<br/>avg, IF5.98<br/>L-index

#	Paper	IF	Citations
101	Microbial tryptophan catabolites in health and disease. <i>Nature Communications</i> , <b>2018</b> , 9, 3294	17.4	532
100	A catalog of the mouse gut metagenome. <i>Nature Biotechnology</i> , <b>2015</b> , 33, 1103-8	44.5	295
99	Establishment of intestinal microbiota during early life: a longitudinal, explorative study of a large cohort of Danish infants. <i>Applied and Environmental Microbiology</i> , <b>2014</b> , 80, 2889-900	4.8	289
98	Colonic transit time is related to bacterial metabolism and mucosal turnover in the gut. <i>Nature Microbiology</i> , <b>2016</b> , 1, 16093	26.6	204
97	Choice of bacterial DNA extraction method from fecal material influences community structure as evaluated by metagenomic analysis. <i>Microbiome</i> , <b>2014</b> , 2, 19	16.6	167
96	Aberrant intestinal microbiota in individuals with prediabetes. <i>Diabetologia</i> , <b>2018</b> , 61, 810-820	10.3	163
95	Whole grain-rich diet reduces body weight and systemic low-grade inflammation without inducing major changes of the gut microbiome: a randomised cross-over trial. <i>Gut</i> , <b>2019</b> , 68, 83-93	19.2	162
94	Freezing fecal samples prior to DNA extraction affects the Firmicutes to Bacteroidetes ratio determined by downstream quantitative PCR analysis. <i>FEMS Microbiology Letters</i> , <b>2012</b> , 329, 193-7	2.9	155
93	Plasmid transfer in the animal intestine and other dynamic bacterial populations: the role of community structure and environment. <i>Microbiology (United Kingdom)</i> , <b>1999</b> , 145 ( Pt 9), 2615-2622	2.9	128
92	Antibiotic Treatment Affects Intestinal Permeability and Gut Microbial Composition in Wistar Rats Dependent on Antibiotic Class. <i>PLoS ONE</i> , <b>2015</b> , 10, e0144854	3.7	123
91	Infant Gut Microbiota Development Is Driven by Transition to Family Foods Independent of Maternal Obesity. <i>MSphere</i> , <b>2016</b> , 1,	5	118
90	Intake of whole apples or clear apple juice has contrasting effects on plasma lipids in healthy volunteers. <i>European Journal of Nutrition</i> , <b>2013</b> , 52, 1875-89	5.2	103
89	Microbial enterotypes, inferred by the prevotella-to-bacteroides ratio, remained stable during a 6-month randomized controlled diet intervention with the new nordic diet. <i>Applied and Environmental Microbiology</i> , <b>2014</b> , 80, 1142-9	4.8	101
88	Qualified presumption of safety (QPS): a generic risk assessment approach for biological agents notified to the European Food Safety Authority (EFSA). <i>Trends in Food Science and Technology</i> , <b>2010</b> , 21, 425-435	15.3	99
87	Transfer of gut microbiota from lean and obese mice to antibiotic-treated mice. <i>Scientific Reports</i> , <b>2014</b> , 4, 5922	4.9	93
86	First Foods and Gut Microbes. Frontiers in Microbiology, 2017, 8, 356	5.7	93
85	Selective pressure affects transfer and establishment of a Lactobacillus plantarum resistance plasmid in the gastrointestinal environment. <i>Journal of Antimicrobial Chemotherapy</i> , <b>2008</b> , 61, 845-52	5.1	82

# (2014-2017)

84	High-fat feeding rather than obesity drives taxonomical and functional changes in the gut microbiota in mice. <i>Microbiome</i> , <b>2017</b> , 5, 43	16.6	77
83	Effects of apples and specific apple components on the cecal environment of conventional rats: role of apple pectin. <i>BMC Microbiology</i> , <b>2010</b> , 10, 13	4.5	75
82	Effects of Lactococcus lactis on composition of intestinal microbiota: role of nisin. <i>Applied and Environmental Microbiology</i> , <b>2006</b> , 72, 239-44	4.8	72
81	Tailored enzymatic production of oligosaccharides from sugar beet pectin and evidence of differential effects of a single DP chain length difference on human faecal microbiota composition after in vitro fermentation. <i>Process Biochemistry</i> , <b>2011</b> , 46, 1039-1049	4.8	71
80	Differential bacterial capture and transport preferences facilitate co-growth on dietary xylan in the human gut. <i>Nature Microbiology</i> , <b>2018</b> , 3, 570-580	26.6	70
79	Having older siblings is associated with gut microbiota development during early childhood. <i>BMC Microbiology</i> , <b>2015</b> , 15, 154	4.5	69
78	A low-gluten diet induces changes in the intestinal microbiome of healthy Danish adults. <i>Nature Communications</i> , <b>2018</b> , 9, 4630	17.4	69
77	Glyphosate has limited short-term effects on commensal bacterial community composition in the gut environment due to sufficient aromatic amino acid levels. <i>Environmental Pollution</i> , <b>2018</b> , 233, 364-3	76 <sup>3</sup>	66
76	Dietary xylooligosaccharide downregulates IFN-land the low-grade inflammatory cytokine IL-1 systemically in mice. <i>Journal of Nutrition</i> , <b>2013</b> , 143, 533-40	4.1	64
75	Effect of Antibiotics on Gut Microbiota, Gut Hormones and Glucose Metabolism. <i>PLoS ONE</i> , <b>2015</b> , 10, e0142352	3.7	61
74	Introducing GUt low-density array (GULDA): a validated approach for qPCR-based intestinal microbial community analysis. <i>FEMS Microbiology Letters</i> , <b>2012</b> , 337, 38-47	2.9	61
73	Maximal release of highly bifidogenic soluble dietary fibers from industrial potato pulp by minimal enzymatic treatment. <i>Applied Microbiology and Biotechnology</i> , <b>2011</b> , 90, 873-84	5.7	60
72	Plasmid transfer from Pseudomonas putida to the indigenous bacteria on alfalfa sprouts: characterization, direct quantification, and in situ location of transconjugant cells. <i>Applied and Environmental Microbiology</i> , <b>2003</b> , 69, 5536-42	4.8	57
71	Feruloylated and nonferuloylated arabino-oligosaccharides from sugar beet pectin selectively stimulate the growth of Bifidobacterium spp. in human fecal in vitro fermentations. <i>Journal of Agricultural and Food Chemistry</i> , <b>2011</b> , 59, 6511-9	5.7	56
70	In vitro fermentation of sugar beet arabino-oligosaccharides by fecal microbiota obtained from patients with ulcerative colitis to selectively stimulate the growth of Bifidobacterium spp. and Lactobacillus spp. <i>Applied and Environmental Microbiology</i> , <b>2011</b> , 77, 8336-44	4.8	55
69	Some putative prebiotics increase the severity of Salmonella enterica serovar Typhimurium infection in mice. <i>BMC Microbiology</i> , <b>2009</b> , 9, 245	4.5	53
68	Klebsiella pneumoniae capsule expression is necessary for colonization of large intestines of streptomycin-treated mice. <i>Infection and Immunity</i> , <b>1999</b> , 67, 6152-6	3.7	53
67	Dietary xylo-oligosaccharide stimulates intestinal bifidobacteria and lactobacilli but has limited effect on intestinal integrity in rats. <i>BMC Research Notes</i> , <b>2014</b> , 7, 660	2.3	51

66	Effect of tetracycline on transfer and establishment of the tetracycline-inducible conjugative transposon Tn916 in the guts of gnotobiotic rats. <i>Applied and Environmental Microbiology</i> , <b>2004</b> , 70, 75	8- <del>6</del> :8	50
65	Fatty acid composition and phospholipid types used in infant formulas modifies the establishment of human gut bacteria in germ-free mice. <i>Scientific Reports</i> , <b>2017</b> , 7, 3975	4.9	49
64	Characterization of transferable tetracycline resistance genes in Enterococcus faecalis isolated from raw food. <i>FEMS Microbiology Letters</i> , <b>2005</b> , 243, 15-9	2.9	49
63	Dietary carbohydrate source influences molecular fingerprints of the rat faecal microbiota. <i>BMC Microbiology</i> , <b>2006</b> , 6, 98	4.5	48
62	Oxygen restriction increases the infective potential of Listeria monocytogenes in vitro in Caco-2 cells and in vivo in guinea pigs. <i>BMC Microbiology</i> , <b>2007</b> , 7, 55	4.5	47
61	Inhibition of Escherichia coli precursor-16S rRNA processing by mouse intestinal contents. <i>Environmental Microbiology</i> , <b>1999</b> , 1, 23-32	5.2	44
60	Estimation of growth rates of Escherichia coli BJ4 in streptomycin-treated and previously germfree mice by in situ rRNA hybridization. <i>Vaccine Journal</i> , <b>1999</b> , 6, 434-6		44
59	Environmental spread of microbes impacts the development of metabolic phenotypes in mice transplanted with microbial communities from humans. <i>ISME Journal</i> , <b>2017</b> , 11, 676-690	11.9	41
58	Nature of bacterial colonization influences transcription of mucin genes in mice during the first week of life. <i>BMC Research Notes</i> , <b>2012</b> , 5, 402	2.3	39
57	Transfer of the pheromone-inducible plasmid pCF10 among Enterococcus faecalis microorganisms colonizing the intestine of mini-pigs. <i>Applied and Environmental Microbiology</i> , <b>2002</b> , 68, 187-93	4.8	38
56	Administration of two probiotic strains during early childhood does not affect the endogenous gut microbiota composition despite probiotic proliferation. <i>BMC Microbiology</i> , <b>2017</b> , 17, 175	4.5	37
55	In vivo detection and quantification of tetracycline by use of a whole-cell biosensor in the rat intestine. <i>Antimicrobial Agents and Chemotherapy</i> , <b>2004</b> , 48, 1112-7	5.9	36
54	A comparative analysis of the intestinal metagenomes present in guinea pigs (Cavia porcellus) and humans (Homo sapiens). <i>BMC Genomics</i> , <b>2012</b> , 13, 514	4.5	35
53	Construction of a multiple fluorescence labelling system for use in co-invasion studies of Listeria monocytogenes. <i>BMC Microbiology</i> , <b>2006</b> , 6, 86	4.5	35
52	Comparison of methods and animal models commonly used for investigation of fecal microbiota: effects of time, host and gender. <i>Journal of Microbiological Methods</i> , <b>2006</b> , 66, 87-95	2.8	35
51	Effect of a long-term high-protein diet on survival, obesity development, and gut microbiota in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , <b>2016</b> , 310, E886-99	6	34
50	Prebiotics for prevention of gut infections. <i>Trends in Food Science and Technology</i> , <b>2012</b> , 23, 70-82	15.3	32
49	Microbiotas from UC patients display altered metabolism and reduced ability of LAB to colonize mucus. <i>Scientific Reports</i> , <b>2013</b> , 3, 1110	4.9	30

### (2020-2005)

48	Conjugative Gene Transfer in the Gastrointestinal Environment. <i>Advances in Applied Microbiology</i> , <b>2005</b> , 58C, 77-95	4.9	29
47	Microbiota composition of simultaneously colonized mice housed under either a gnotobiotic isolator or individually ventilated cage regime. <i>Scientific Reports</i> , <b>2017</b> , 7, 42245	4.9	28
46	Lipid hydrolysis products affect the composition of infant gut microbial communities in vitro. <i>British Journal of Nutrition</i> , <b>2015</b> , 114, 63-74	3.6	25
45	Selection of bacteria originating from a human intestinal microbiota in the gut of previously germ-free rats. <i>FEMS Microbiology Letters</i> , <b>2007</b> , 277, 205-9	2.9	25
44	Bifidobacterium species associated with breastfeeding produce aromatic lactic acids in the infant gut. <i>Nature Microbiology</i> , <b>2021</b> , 6, 1367-1382	26.6	25
43	Xylo-oligosaccharides inhibit pathogen adhesion to enterocytes in vitro. <i>Research in Microbiology</i> , <b>2012</b> , 163, 22-7	4	24
42	Lactobacillus acidophilus induces a slow but more sustained chemokine and cytokine response in naMe foetal enterocytes compared to commensal Escherichia coli. <i>BMC Immunology</i> , <b>2010</b> , 11, 2	3.7	23
41	Certain dietary carbohydrates promote Listeria infection in a guinea pig model, while others prevent it. <i>International Journal of Food Microbiology</i> , <b>2010</b> , 140, 218-24	5.8	23
40	Lactobacillus plantarum inhibits growth of Listeria monocytogenes in an in vitro continuous flow gut model, but promotes invasion of L. monocytogenes in the gut of gnotobiotic rats. <i>International Journal of Food Microbiology</i> , <b>2006</b> , 108, 10-4	5.8	22
39	A combined metabolomic and phylogenetic study reveals putatively prebiotic effects of high molecular weight arabino-oligosaccharides when assessed by in vitro fermentation in bacterial communities derived from humans. <i>Anaerobe</i> , <b>2014</b> , 28, 68-77	2.8	21
38	Pediocin PA-1 and a pediocin producing Lactobacillus plantarum strain do not change the HMA rat microbiota. <i>International Journal of Food Microbiology</i> , <b>2009</b> , 130, 251-7	5.8	21
37	Effects of Gliadin consumption on the Intestinal Microbiota and Metabolic Homeostasis in Mice Fed a High-fat Diet. <i>Scientific Reports</i> , <b>2017</b> , 7, 44613	4.9	19
36	Comparison of three Listeria monocytogenes strains in a guinea-pig model simulating food-borne exposure. <i>FEMS Microbiology Letters</i> , <b>2009</b> , 291, 88-94	2.9	19
35	Conjugative transfer facilitates stable maintenance of IncP-1 plasmid pKJK5 in Escherichia coli cells colonizing the gastrointestinal tract of the germfree rat. <i>Applied and Environmental Microbiology</i> , <b>2007</b> , 73, 341-3	4.8	19
34	A single exposure to a sublethal pediocin concentration initiates a resistance-associated temporal cell envelope and general stress response in Listeria monocytogenes. <i>Environmental Microbiology</i> , <b>2015</b> , 17, 1134-51	5.2	18
33	Gut Colonization Is Accelerated by Presence of Older Siblings. <i>MSphere</i> , <b>2017</b> , 2,	5	18
32	A functional cra gene is required for Salmonella enterica serovar typhimurium virulence in BALB/c mice. <i>Infection and Immunity</i> , <b>2000</b> , 68, 3772-5	3.7	18
31	Human microbiota-transplanted C57BL/6 mice and offspring display reduced establishment of key bacteria and reduced immune stimulation compared to mouse microbiota-transplantation.  Scientific Reports, 2020, 10, 7805	4.9	17

30	Listeria monocytogenes strains encoding premature stop codons in inlA invade mice and guinea pig fetuses in orally dosed dams. <i>Journal of Medical Microbiology</i> , <b>2013</b> , 62, 1799-1806	3.2	17
29	Evidence of increased spread and establishment of plasmid RP4 in the intestine under sub-inhibitory tetracycline concentrations. <i>FEMS Microbiology Ecology</i> , <b>2003</b> , 44, 217-23	4.3	17
28	Monitoring bacterial growth activity in biofilms from laboratory flow chambers, plant rhizosphere, and animal intestine. <i>Methods in Enzymology</i> , <b>2001</b> , 337, 21-42	1.7	16
27	Lactobacillus acidophilus NCFM affects vitamin E acetate metabolism and intestinal bile acid signature in monocolonized mice. <i>Gut Microbes</i> , <b>2014</b> , 5, 296-303	8.8	15
26	Xylo-oligosaccharides and inulin affect genotoxicity and bacterial populations differently in a human colonic simulator challenged with soy protein. <i>Nutrients</i> , <b>2013</b> , 5, 3740-56	6.7	14
25	In vitro growth of four individual human gut bacteria on oligosaccharides produced by chemoenzymatic synthesis. <i>Food and Function</i> , <b>2013</b> , 4, 784-93	6.1	13
24	Impact of the gut microbiota on chemical risk assessment. Current Opinion in Toxicology, 2019, 15, 109-1	I 1434	13
23	Breastmilk-promoted bifidobacteria produce aromatic amino acids in the infant gut		12
22	Short-Term Amoxicillin-Induced Perturbation of the Gut Microbiota Promotes Acute Intestinal Immune Regulation in Brown Norway Rats. <i>Frontiers in Microbiology</i> , <b>2020</b> , 11, 496	5.7	11
21	Metabolic footprint of Lactobacillus acidophilus NCFM at different pH. <i>Metabolomics</i> , <b>2012</b> , 8, 244-252	4.7	10
20	Intestinal Enterococcus abundance correlates inversely with excessive weight gain and increased plasma leptin in breastfed infants. <i>FEMS Microbiology Ecology</i> , <b>2020</b> , 96,	4.3	9
19	Determining Gut Microbial Dysbiosis: a Review of Applied Indexes for Assessment of Intestinal Microbiota Imbalances. <i>Applied and Environmental Microbiology</i> , <b>2021</b> , 87,	4.8	9
18	Neonatal microbial colonization in mice promotes prolonged dominance of CD11b(+)Gr-1(+) cells and accelerated establishment of the CD4(+) T cell population in the spleen. <i>Immunity, Inflammation and Disease,</i> <b>2015</b> , 3, 309-20	2.4	8
17	Settlers of our inner surface - factors shaping the gut microbiota from birth to toddlerhood. <i>FEMS Microbiology Reviews</i> , <b>2021</b> , 45,	15.1	8
16	Antibiotic treatment of rat dams affects bacterial colonization and causes decreased weight gain in pups. <i>Communications Biology</i> , <b>2018</b> , 1, 145	6.7	8
15	Effect of pheromone induction on transfer of the Enterococcus faecalis plasmid pCF10 in intestinal mucus ex vivo. <i>FEMS Microbiology Letters</i> , <b>2001</b> , 204, 305-9	2.9	6
14	Colon-Specific Delivery of Bioactive Agents Using Genipin-Cross-Linked Chitosan Coated Microcontainers. <i>ACS Applied Bio Materials</i> , <b>2021</b> , 4, 752-762	4.1	6
13	Maternal milk microbiota and oligosaccharides contribute to the infant gut microbiota assembly.  ISME Communications, 2021, 1,		5

#### LIST OF PUBLICATIONS

12	Multi-layer PLGA-pullulan-PLGA electrospun nanofibers for probiotic delivery. <i>Food Hydrocolloids</i> , <b>2022</b> , 123, 107112	10.6	4
11	Faecal microbiota transplantation for eradication of co-infection with and extensively drug-resistant KPC-producing. <i>Scandinavian Journal of Gastroenterology</i> , <b>2020</b> , 55, 626-630	2.4	3
10	Effect of the vitamin B12-binding protein haptocorrin present in human milk on a panel of commensal and pathogenic bacteria. <i>BMC Research Notes</i> , <b>2011</b> , 4, 208	2.3	3
9	Data integration for prediction of weight loss in randomized controlled dietary trials. <i>Scientific Reports</i> , <b>2020</b> , 10, 20103	4.9	2
8	Partially Hydrolysed Whey Has Superior Allergy Preventive Capacity Compared to Intact Whey Regardless of Amoxicillin Administration in Brown Norway Rats. <i>Frontiers in Immunology</i> , <b>2021</b> , 12, 7055	;8 <del>31</del>	2
7	Amoxicillin does not affect the development of cows milk allergy in a Brown Norway rat model <i>Scandinavian Journal of Immunology</i> , <b>2022</b> , e13148	3.4	1
6	The intestinal microbiome is a co-determinant of the postprandial plasma glucose response. <i>PLoS ONE</i> , <b>2020</b> , 15, e0238648	3.7	1
5	Casein glycomacropeptide is well tolerated in healthy adults and changes neither high-sensitive C-reactive protein, gut microbiota nor faecal butyrate: a restricted randomised trial. <i>British Journal of Nutrition</i> , <b>2021</b> , 125, 1374-1385	3.6	1
4	Optimizing oral delivery of next generation probiotics. <i>Trends in Food Science and Technology</i> , <b>2022</b> , 119, 101-109	15.3	O
3	Inter-microbial relations shape the preterm gut. <i>Nature Microbiology</i> , <b>2021</b> , 6, 423-424	26.6	
2	Systems-wide effects of short-term feed deprivation in obese mice. Scientific Reports, 2021, 11, 5716	4.9	
1	Amendments: Author Correction: A catalog of the mouse gut metagenome. <i>Nature Biotechnology</i> , <b>2019</b> , 37, 102	44.5	