

Tine Rask Licht

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5458612/tine-rask-licht-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

101
papers

5,380
citations

42
h-index

72
g-index

104
ext. papers

6,852
ext. citations

7.2
avg, IF

5.98
L-index

#	Paper	IF	Citations
101	Amoxicillin does not affect the development of cow's milk allergy in a Brown Norway rat model.. <i>Scandinavian Journal of Immunology</i> , 2022 , e13148	3.4	1
100	Optimizing oral delivery of next generation probiotics. <i>Trends in Food Science and Technology</i> , 2022 , 119, 101-109	15.3	0
99	Multi-layer PLGA-pullulan-PLGA electrospun nanofibers for probiotic delivery. <i>Food Hydrocolloids</i> , 2022 , 123, 107112	10.6	4
98	Bifidobacterium species associated with breastfeeding produce aromatic lactic acids in the infant gut. <i>Nature Microbiology</i> , 2021 , 6, 1367-1382	26.6	25
97	Colon-Specific Delivery of Bioactive Agents Using Genipin-Cross-Linked Chitosan Coated Microcontainers. <i>ACS Applied Bio Materials</i> , 2021 , 4, 752-762	4.1	6
96	Inter-microbial relations shape the preterm gut. <i>Nature Microbiology</i> , 2021 , 6, 423-424	26.6	
95	Systems-wide effects of short-term feed deprivation in obese mice. <i>Scientific Reports</i> , 2021 , 11, 5716	4.9	
94	Determining Gut Microbial Dysbiosis: a Review of Applied Indexes for Assessment of Intestinal Microbiota Imbalances. <i>Applied and Environmental Microbiology</i> , 2021 , 87,	4.8	9
93	Maternal milk microbiota and oligosaccharides contribute to the infant gut microbiota assembly. <i>ISME Communications</i> , 2021 , 1,		5
92	Casein glycomacropeptide is well tolerated in healthy adults and changes neither high-sensitive C-reactive protein, gut microbiota nor faecal butyrate: a restricted randomised trial. <i>British Journal of Nutrition</i> , 2021 , 125, 1374-1385	3.6	1
91	Settlers of our inner surface - factors shaping the gut microbiota from birth to toddlerhood. <i>FEMS Microbiology Reviews</i> , 2021 , 45,	15.1	8
90	Partially Hydrolysed Whey Has Superior Allergy Preventive Capacity Compared to Intact Whey Regardless of Amoxicillin Administration in Brown Norway Rats. <i>Frontiers in Immunology</i> , 2021 , 12, 705543	8.4	2
89	Human microbiota-transplanted C57BL/6 mice and offspring display reduced establishment of key bacteria and reduced immune stimulation compared to mouse microbiota-transplantation. <i>Scientific Reports</i> , 2020 , 10, 7805	4.9	17
88	Short-Term Amoxicillin-Induced Perturbation of the Gut Microbiota Promotes Acute Intestinal Immune Regulation in Brown Norway Rats. <i>Frontiers in Microbiology</i> , 2020 , 11, 496	5.7	11
87	Faecal microbiota transplantation for eradication of co-infection with and extensively drug-resistant KPC-producing. <i>Scandinavian Journal of Gastroenterology</i> , 2020 , 55, 626-630	2.4	3
86	Intestinal Enterococcus abundance correlates inversely with excessive weight gain and increased plasma leptin in breastfed infants. <i>FEMS Microbiology Ecology</i> , 2020 , 96,	4.3	9
85	The intestinal microbiome is a co-determinant of the postprandial plasma glucose response. <i>PLoS ONE</i> , 2020 , 15, e0238648	3.7	1

84	Data integration for prediction of weight loss in randomized controlled dietary trials. <i>Scientific Reports</i> , 2020 , 10, 20103	4.9	2
83	Amendments: Author Correction: A catalog of the mouse gut metagenome. <i>Nature Biotechnology</i> , 2019 , 37, 102	44.5	
82	Impact of the gut microbiota on chemical risk assessment. <i>Current Opinion in Toxicology</i> , 2019 , 15, 109-113	13	13
81	Whole grain-rich diet reduces body weight and systemic low-grade inflammation without inducing major changes of the gut microbiome: a randomised cross-over trial. <i>Gut</i> , 2019 , 68, 83-93	19.2	162
80	Aberrant intestinal microbiota in individuals with prediabetes. <i>Diabetologia</i> , 2018 , 61, 810-820	10.3	163
79	Differential bacterial capture and transport preferences facilitate co-growth on dietary xylan in the human gut. <i>Nature Microbiology</i> , 2018 , 3, 570-580	26.6	70
78	Microbial tryptophan catabolites in health and disease. <i>Nature Communications</i> , 2018 , 9, 3294	17.4	532
77	Glyphosate has limited short-term effects on commensal bacterial community composition in the gut environment due to sufficient aromatic amino acid levels. <i>Environmental Pollution</i> , 2018 , 233, 364-376	2.3	66
76	A low-gluten diet induces changes in the intestinal microbiome of healthy Danish adults. <i>Nature Communications</i> , 2018 , 9, 4630	17.4	69
75	Antibiotic treatment of rat dams affects bacterial colonization and causes decreased weight gain in pups. <i>Communications Biology</i> , 2018 , 1, 145	6.7	8
74	Microbiota composition of simultaneously colonized mice housed under either a gnotobiotic isolator or individually ventilated cage regime. <i>Scientific Reports</i> , 2017 , 7, 42245	4.9	28
73	High-fat feeding rather than obesity drives taxonomical and functional changes in the gut microbiota in mice. <i>Microbiome</i> , 2017 , 5, 43	16.6	77
72	Effects of Gliadin consumption on the Intestinal Microbiota and Metabolic Homeostasis in Mice Fed a High-fat Diet. <i>Scientific Reports</i> , 2017 , 7, 44613	4.9	19
71	Administration of two probiotic strains during early childhood does not affect the endogenous gut microbiota composition despite probiotic proliferation. <i>BMC Microbiology</i> , 2017 , 17, 175	4.5	37
70	Fatty acid composition and phospholipid types used in infant formulas modifies the establishment of human gut bacteria in germ-free mice. <i>Scientific Reports</i> , 2017 , 7, 3975	4.9	49
69	Environmental spread of microbes impacts the development of metabolic phenotypes in mice transplanted with microbial communities from humans. <i>ISME Journal</i> , 2017 , 11, 676-690	11.9	41
68	Gut Colonization Is Accelerated by Presence of Older Siblings. <i>MSphere</i> , 2017 , 2,	5	18
67	First Foods and Gut Microbes. <i>Frontiers in Microbiology</i> , 2017 , 8, 356	5.7	93

66	Colonic transit time is related to bacterial metabolism and mucosal turnover in the gut. <i>Nature Microbiology</i> , 2016 , 1, 16093	26.6	204
65	Effect of a long-term high-protein diet on survival, obesity development, and gut microbiota in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016 , 310, E886-99	6	34
64	Infant Gut Microbiota Development Is Driven by Transition to Family Foods Independent of Maternal Obesity. <i>MSphere</i> , 2016 , 1,	5	118
63	A catalog of the mouse gut metagenome. <i>Nature Biotechnology</i> , 2015 , 33, 1103-8	44.5	295
62	A single exposure to a sublethal pediocin concentration initiates a resistance-associated temporal cell envelope and general stress response in <i>Listeria monocytogenes</i> . <i>Environmental Microbiology</i> , 2015 , 17, 1134-51	5.2	18
61	Lipid hydrolysis products affect the composition of infant gut microbial communities in vitro. <i>British Journal of Nutrition</i> , 2015 , 114, 63-74	3.6	25
60	Neonatal microbial colonization in mice promotes prolonged dominance of CD11b(+)Gr-1(+) cells and accelerated establishment of the CD4(+) T cell population in the spleen. <i>Immunity, Inflammation and Disease</i> , 2015 , 3, 309-20	2.4	8
59	Effect of Antibiotics on Gut Microbiota, Gut Hormones and Glucose Metabolism. <i>PLoS ONE</i> , 2015 , 10, e0142352	3.7	61
58	Antibiotic Treatment Affects Intestinal Permeability and Gut Microbial Composition in Wistar Rats Dependent on Antibiotic Class. <i>PLoS ONE</i> , 2015 , 10, e0144854	3.7	123
57	Having older siblings is associated with gut microbiota development during early childhood. <i>BMC Microbiology</i> , 2015 , 15, 154	4.5	69
56	Transfer of gut microbiota from lean and obese mice to antibiotic-treated mice. <i>Scientific Reports</i> , 2014 , 4, 5922	4.9	93
55	Microbial enterotypes, inferred by the prevotella-to-bacteroides ratio, remained stable during a 6-month randomized controlled diet intervention with the new nordic diet. <i>Applied and Environmental Microbiology</i> , 2014 , 80, 1142-9	4.8	101
54	Establishment of intestinal microbiota during early life: a longitudinal, explorative study of a large cohort of Danish infants. <i>Applied and Environmental Microbiology</i> , 2014 , 80, 2889-900	4.8	289
53	Choice of bacterial DNA extraction method from fecal material influences community structure as evaluated by metagenomic analysis. <i>Microbiome</i> , 2014 , 2, 19	16.6	167
52	A combined metabolomic and phylogenetic study reveals putatively prebiotic effects of high molecular weight arabino-oligosaccharides when assessed by <i>in vitro</i> fermentation in bacterial communities derived from humans. <i>Anaerobe</i> , 2014 , 28, 68-77	2.8	21
51	<i>Lactobacillus acidophilus</i> NCFM affects vitamin E acetate metabolism and intestinal bile acid signature in monocolonized mice. <i>Gut Microbes</i> , 2014 , 5, 296-303	8.8	15
50	Dietary xylo-oligosaccharide stimulates intestinal bifidobacteria and lactobacilli but has limited effect on intestinal integrity in rats. <i>BMC Research Notes</i> , 2014 , 7, 660	2.3	51
49	Intake of whole apples or clear apple juice has contrasting effects on plasma lipids in healthy volunteers. <i>European Journal of Nutrition</i> , 2013 , 52, 1875-89	5.2	103

48	In vitro growth of four individual human gut bacteria on oligosaccharides produced by chemoenzymatic synthesis. <i>Food and Function</i> , 2013 , 4, 784-93	6.1	13
47	<i>Listeria monocytogenes</i> strains encoding premature stop codons in inlA invade mice and guinea pig fetuses in orally dosed dams. <i>Journal of Medical Microbiology</i> , 2013 , 62, 1799-1806	3.2	17
46	Dietary xylooligosaccharide downregulates IFN- γ and the low-grade inflammatory cytokine IL-1 β systemically in mice. <i>Journal of Nutrition</i> , 2013 , 143, 533-40	4.1	64
45	Microbiotas from UC patients display altered metabolism and reduced ability of LAB to colonize mucus. <i>Scientific Reports</i> , 2013 , 3, 1110	4.9	30
44	Xylo-oligosaccharides and inulin affect genotoxicity and bacterial populations differently in a human colonic simulator challenged with soy protein. <i>Nutrients</i> , 2013 , 5, 3740-56	6.7	14
43	Xylo-oligosaccharides inhibit pathogen adhesion to enterocytes in vitro. <i>Research in Microbiology</i> , 2012 , 163, 22-7	4	24
42	Prebiotics for prevention of gut infections. <i>Trends in Food Science and Technology</i> , 2012 , 23, 70-82	15.3	32
41	A comparative analysis of the intestinal metagenomes present in guinea pigs (<i>Cavia porcellus</i>) and humans (<i>Homo sapiens</i>). <i>BMC Genomics</i> , 2012 , 13, 514	4.5	35
40	Nature of bacterial colonization influences transcription of mucin genes in mice during the first week of life. <i>BMC Research Notes</i> , 2012 , 5, 402	2.3	39
39	Introducing GUT low-density array (GULDA): a validated approach for qPCR-based intestinal microbial community analysis. <i>FEMS Microbiology Letters</i> , 2012 , 337, 38-47	2.9	61
38	Metabolic footprint of <i>Lactobacillus acidophilus</i> NCFM at different pH. <i>Metabolomics</i> , 2012 , 8, 244-252	4.7	10
37	Freezing fecal samples prior to DNA extraction affects the Firmicutes to Bacteroidetes ratio determined by downstream quantitative PCR analysis. <i>FEMS Microbiology Letters</i> , 2012 , 329, 193-7	2.9	155
36	Feruloylated and nonferuloylated arabino-oligosaccharides from sugar beet pectin selectively stimulate the growth of <i>Bifidobacterium</i> spp. in human fecal in vitro fermentations. <i>Journal of Agricultural and Food Chemistry</i> , 2011 , 59, 6511-9	5.7	56
35	Effect of the vitamin B12-binding protein haptocorrin present in human milk on a panel of commensal and pathogenic bacteria. <i>BMC Research Notes</i> , 2011 , 4, 208	2.3	3
34	Maximal release of highly bifidogenic soluble dietary fibers from industrial potato pulp by minimal enzymatic treatment. <i>Applied Microbiology and Biotechnology</i> , 2011 , 90, 873-84	5.7	60
33	Tailored enzymatic production of oligosaccharides from sugar beet pectin and evidence of differential effects of a single DP chain length difference on human faecal microbiota composition after in vitro fermentation. <i>Process Biochemistry</i> , 2011 , 46, 1039-1049	4.8	71
32	In vitro fermentation of sugar beet arabino-oligosaccharides by fecal microbiota obtained from patients with ulcerative colitis to selectively stimulate the growth of <i>Bifidobacterium</i> spp. and <i>Lactobacillus</i> spp. <i>Applied and Environmental Microbiology</i> , 2011 , 77, 8336-44	4.8	55
31	Qualified presumption of safety (QPS): a generic risk assessment approach for biological agents notified to the European Food Safety Authority (EFSA). <i>Trends in Food Science and Technology</i> , 2010 , 21, 425-435	15.3	99

30	Lactobacillus acidophilus induces a slow but more sustained chemokine and cytokine response in naïve foetal enterocytes compared to commensal Escherichia coli. <i>BMC Immunology</i> , 2010 , 11, 2	3.7	23
29	Effects of apples and specific apple components on the cecal environment of conventional rats: role of apple pectin. <i>BMC Microbiology</i> , 2010 , 10, 13	4.5	75
28	Certain dietary carbohydrates promote Listeria infection in a guinea pig model, while others prevent it. <i>International Journal of Food Microbiology</i> , 2010 , 140, 218-24	5.8	23
27	Some putative prebiotics increase the severity of Salmonella enterica serovar Typhimurium infection in mice. <i>BMC Microbiology</i> , 2009 , 9, 245	4.5	53
26	Pediocin PA-1 and a pediocin producing Lactobacillus plantarum strain do not change the HMA rat microbiota. <i>International Journal of Food Microbiology</i> , 2009 , 130, 251-7	5.8	21
25	Comparison of three Listeria monocytogenes strains in a guinea-pig model simulating food-borne exposure. <i>FEMS Microbiology Letters</i> , 2009 , 291, 88-94	2.9	19
24	Selective pressure affects transfer and establishment of a Lactobacillus plantarum resistance plasmid in the gastrointestinal environment. <i>Journal of Antimicrobial Chemotherapy</i> , 2008 , 61, 845-52	5.1	82
23	Selection of bacteria originating from a human intestinal microbiota in the gut of previously germ-free rats. <i>FEMS Microbiology Letters</i> , 2007 , 277, 205-9	2.9	25
22	Oxygen restriction increases the infective potential of Listeria monocytogenes in vitro in Caco-2 cells and in vivo in guinea pigs. <i>BMC Microbiology</i> , 2007 , 7, 55	4.5	47
21	Conjugative transfer facilitates stable maintenance of IncP-1 plasmid pKJK5 in Escherichia coli cells colonizing the gastrointestinal tract of the germfree rat. <i>Applied and Environmental Microbiology</i> , 2007 , 73, 341-3	4.8	19
20	Lactobacillus plantarum inhibits growth of Listeria monocytogenes in an in vitro continuous flow gut model, but promotes invasion of L. monocytogenes in the gut of gnotobiotic rats. <i>International Journal of Food Microbiology</i> , 2006 , 108, 10-4	5.8	22
19	Construction of a multiple fluorescence labelling system for use in co-invasion studies of Listeria monocytogenes. <i>BMC Microbiology</i> , 2006 , 6, 86	4.5	35
18	Dietary carbohydrate source influences molecular fingerprints of the rat faecal microbiota. <i>BMC Microbiology</i> , 2006 , 6, 98	4.5	48
17	Effects of Lactococcus lactis on composition of intestinal microbiota: role of nisin. <i>Applied and Environmental Microbiology</i> , 2006 , 72, 239-44	4.8	72
16	Comparison of methods and animal models commonly used for investigation of fecal microbiota: effects of time, host and gender. <i>Journal of Microbiological Methods</i> , 2006 , 66, 87-95	2.8	35
15	Characterization of transferable tetracycline resistance genes in Enterococcus faecalis isolated from raw food. <i>FEMS Microbiology Letters</i> , 2005 , 243, 15-9	2.9	49
14	Conjugative Gene Transfer in the Gastrointestinal Environment. <i>Advances in Applied Microbiology</i> , 2005 , 58C, 77-95	4.9	29
13	In vivo detection and quantification of tetracycline by use of a whole-cell biosensor in the rat intestine. <i>Antimicrobial Agents and Chemotherapy</i> , 2004 , 48, 1112-7	5.9	36

12	Effect of tetracycline on transfer and establishment of the tetracycline-inducible conjugative transposon Tn916 in the guts of gnotobiotic rats. <i>Applied and Environmental Microbiology</i> , 2004 , 70, 758-64	4.8	50
11	Evidence of increased spread and establishment of plasmid RP4 in the intestine under sub-inhibitory tetracycline concentrations. <i>FEMS Microbiology Ecology</i> , 2003 , 44, 217-23	4.3	17
10	Plasmid transfer from <i>Pseudomonas putida</i> to the indigenous bacteria on alfalfa sprouts: characterization, direct quantification, and in situ location of transconjugant cells. <i>Applied and Environmental Microbiology</i> , 2003 , 69, 5536-42	4.8	57
9	Transfer of the pheromone-inducible plasmid pCF10 among <i>Enterococcus faecalis</i> microorganisms colonizing the intestine of mini-pigs. <i>Applied and Environmental Microbiology</i> , 2002 , 68, 187-93	4.8	38
8	Effect of pheromone induction on transfer of the <i>Enterococcus faecalis</i> plasmid pCF10 in intestinal mucus ex vivo. <i>FEMS Microbiology Letters</i> , 2001 , 204, 305-9	2.9	6
7	Monitoring bacterial growth activity in biofilms from laboratory flow chambers, plant rhizosphere, and animal intestine. <i>Methods in Enzymology</i> , 2001 , 337, 21-42	1.7	16
6	A functional <i>cra</i> gene is required for <i>Salmonella enterica</i> serovar typhimurium virulence in BALB/c mice. <i>Infection and Immunity</i> , 2000 , 68, 3772-5	3.7	18
5	Inhibition of <i>Escherichia coli</i> precursor-16S rRNA processing by mouse intestinal contents. <i>Environmental Microbiology</i> , 1999 , 1, 23-32	5.2	44
4	Estimation of growth rates of <i>Escherichia coli</i> BJ4 in streptomycin-treated and previously germfree mice by in situ rRNA hybridization. <i>Vaccine Journal</i> , 1999 , 6, 434-6		44
3	<i>Klebsiella pneumoniae</i> capsule expression is necessary for colonization of large intestines of streptomycin-treated mice. <i>Infection and Immunity</i> , 1999 , 67, 6152-6	3.7	53
2	Plasmid transfer in the animal intestine and other dynamic bacterial populations: the role of community structure and environment. <i>Microbiology (United Kingdom)</i> , 1999 , 145 (Pt 9), 2615-2622	2.9	128
1	Breastmilk-promoted bifidobacteria produce aromatic amino acids in the infant gut		12