
Makoto Ihara

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5460101/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Ligand-gated ion channels as targets of neuroactive insecticides. Bioscience, Biotechnology and Biochemistry, 2022, 86, 157-164.	1.3	4
2	Competitive chrodrimanin B interactions with rat brain GABAA receptors revealed by radioligand binding assays. Pesticide Biochemistry and Physiology, 2022, 183, 105074.	3.6	0
3	Effects of cofactors RIC-3, TMX3 and UNC-50, together with distinct subunit ratios on the agonist actions of imidacloprid on Drosophila melanogaster $D\hat{1}\pm 1/D\hat{1}^21$ nicotinic acetylcholine receptors expressed in Xenopus laevis oocytes. Pesticide Biochemistry and Physiology, 2022, 187, 105177.	3.6	3
4	A dual-target molecular mechanism of pyrethrum repellency against mosquitoes. Nature Communications, 2021, 12, 2553.	12.8	31
5	Identification of multiple odorant receptors essential for pyrethrum repellency in Drosophila melanogaster. PLoS Genetics, 2021, 17, e1009677.	3.5	10
6	Neonicotinoid Insecticides: Molecular Targets, Resistance, and Toxicity. Annual Review of Pharmacology and Toxicology, 2020, 60, 241-255.	9.4	168
7	The mechanism of loop C-neonicotinoid interactions at insect nicotinic acetylcholine receptor α1 subunit predicts resistance emergence in pests. Scientific Reports, 2020, 10, 7529.	3.3	9
8	Metabolome Analysis Identified Okaramines in the Soybean Rhizosphere as a Legacy of Hairy Vetch. Frontiers in Genetics, 2020, 11, 114.	2.3	13
9	Cofactor-enabled functional expression of fruit fly, honeybee, and bumblebee nicotinic receptors reveals picomolar neonicotinoid actions. Proceedings of the National Academy of Sciences of the United States of America, 2020, 117, 16283-16291.	7.1	61
10	STAT3 inhibitory activity of naphthoquinones isolated from Tabebuia avellanedae. Bioorganic and Medicinal Chemistry, 2020, 28, 115347.	3.0	14
11	Biosynthesis and Structure–Activity Relationship Studies of Okaramines That Target Insect Glutamate-Gated Chloride Channels. ACS Chemical Biology, 2018, 13, 561-566.	3.4	29
12	Combined effects of mutations in loop C and the loop D-E-G triangle on neonicotinoid interactions with Drosophila Dα1/chicken β2 hybrid nAChRs. Pesticide Biochemistry and Physiology, 2018, 151, 47-52.	3.6	11
13	Loops D, E and G in the <i>Drosophila</i> Dα1 subunit contribute to high neonicotinoid sensitivity of Dα1â€chicken β2 nicotinic acetylcholine receptor. British Journal of Pharmacology, 2018, 175, 1999-2012.	5.4	22
14	Neonicotinoids: molecular mechanisms of action, insights into resistance and impact on pollinators. Current Opinion in Insect Science, 2018, 30, 86-92.	4.4	85
15	The fungal alkaloid Okaramine-B activates an L-glutamate-gated chloride channel from Ixodes scapularis, a tick vector of Lyme disease. International Journal for Parasitology: Drugs and Drug Resistance, 2018, 8, 350-360.	3.4	8
16	Availability of NHS-biotin labeling to identify free protein lysine revealed by experiment and MD simulation. Analytical Biochemistry, 2018, 557, 46-58.	2.4	6
17	Action of six pyrethrins purified from the botanical insecticide pyrethrum on cockroach sodium channels expressed in Xenopus oocytes. Pesticide Biochemistry and Physiology, 2018, 151, 82-89.	3.6	25
18	Okaramine insecticidal alkaloids show similar activity on both exon 3c and exon 3b variants of glutamate-gated chloride channels of the larval silkworm, Bombyx mori. NeuroToxicology, 2017, 60, 240-244.	3.0	17

Makoto Ihara

#	Article	IF	CITATIONS
19	Comparative ecotoxicity of imidacloprid and dinotefuran to aquatic insects in rice mesocosms. Ecotoxicology and Environmental Safety, 2017, 138, 122-129.	6.0	42
20	Flupyrimin: A Novel Insecticide Acting at the Nicotinic Acetylcholine Receptors. Journal of Agricultural and Food Chemistry, 2017, 65, 7865-7873.	5.2	40
21	An L319F mutation in transmembrane region 3 (TM3) selectively reduces sensitivity to okaramine B of the <i>Bombyx mori</i> Â <scp>l</scp> -glutamate-gated chloride channel. Bioscience, Biotechnology and Biochemistry, 2017, 81, 1861-1867.	1.3	10
22	Splice Variants of pH-Sensitive Chloride Channel Identify a Key Determinant of Ivermectin Sensitivity in the Larvae of the Silkworm <i>Bombyx mori</i> . Molecular Pharmacology, 2017, 92, 491-499.	2.3	4
23	Modes of Action, Resistance and Toxicity of Insecticides Targeting Nicotinic Acetylcholine Receptors. Current Medicinal Chemistry, 2017, 24, 2925-2934.	2.4	74
24	Mechanisms of Action, Resistance and Toxicity of Insecticides Targeting GABA Receptors. Current Medicinal Chemistry, 2017, 24, 2935-2945.	2.4	58
25	Selective regulation of pyrethrin biosynthesis by the specific blend of wound induced volatiles in <i>Tanacetum cinerariifolium</i> . Plant Signaling and Behavior, 2016, 11, e1149675.	2.4	7
26	lvermectin modulation of pH-sensitive chloride channels in the silkworm larvae of Bombyx mori. Pesticide Biochemistry and Physiology, 2016, 126, 1-5.	3.6	11
27	Probing new components (loop G and the αâ€"î± interface) of neonicotinoid binding sites on nicotinic acetylcholine receptors. Pesticide Biochemistry and Physiology, 2015, 121, 47-52.	3.6	20
28	Meroterpenoid Chrodrimanins Are Selective and Potent Blockers of Insect GABA-Gated Chloride Channels. PLoS ONE, 2015, 10, e0122629.	2.5	22
29	Studies on an Acetylcholine Binding Protein Identify a Basic Residue in Loop G on the <i>β</i> 1 Strand as a New Structural Determinant of Neonicotinoid Actions. Molecular Pharmacology, 2014, 86, 736-746.	2.3	35
30	General flexible nature of the cytosolic regions of fungal transient receptor potential (TRP) channels, revealed by expression screening using GFPâ€fusion techniques. Protein Science, 2014, 23, 923-931.	7.6	1
31	A single amino acid polymorphism in the Drosophila melanogaster Dα1 (ALS) subunit enhances neonicotinoid efficacy at Dα1-chicken β2 hybrid nicotinic acetylcholine receptor expressed in Xenopus laevis oocytes. Bioscience, Biotechnology and Biochemistry, 2014, 78, 543-549.	1.3	4
32	Exon 3 Splicing and Mutagenesis Identify Residues Influencing Cell Surface Density of Heterologously Expressed Silkworm (<i>Bombyx mori</i>) Glutamate-Gated Chloride Channels. Molecular Pharmacology, 2014, 86, 686-695.	2.3	24
33	GluCl a target of indole alkaloid okaramines: a 25 year enigma solved. Scientific Reports, 2014, 4, 6190.	3.3	41
34	Molecular Bases of Multimodal Regulation of a Fungal Transient Receptor Potential (TRP) Channel. Journal of Biological Chemistry, 2013, 288, 15303-15317.	3.4	19
35	GFPâ€based evaluation system of recombinant expression through the secretory pathway in insect cells and its application to the extracellular domains of class C GPCRs. Protein Science, 2011, 20, 1720-1734.	7.6	20
36	High-resolution Native-PAGE for membrane proteins capable of fluorescence detection and hydrodynamic state evaluation. Analytical Biochemistry, 2011, 412, 217-223.	2.4	11

Makoto Ihara

#	Article	IF	CITATIONS
37	Crystal structures of Lymnaea stagnalis AChBP in complex with neonicotinoid insecticides imidacloprid and clothianidin. Invertebrate Neuroscience, 2008, 8, 71-81.	1.8	135
38	Cy3-3-acylcholine: A fluorescent analogue of acetylcholine for single molecule detection. Bioorganic and Medicinal Chemistry Letters, 2008, 18, 1106-1109.	2.2	5
39	Potentiating and blocking actions of neonicotinoids on the response to acetylcholine of the neuronal .ALPHA.4.BETA.2 nicotinic acetylcholine receptor. Journal of Pesticide Sciences, 2008, 33, 146-151.	1.4	18
40	Blocking actions of alkylene-tethered bis-neonicotinoids on nicotinic acetylcholine receptors expressed by terminal abdominal ganglion neurons of Periplaneta americana. Neuroscience Letters, 2007, 425, 137-140.	2.1	16
41	Mechanisms of action of insecticides on ligand-gated ion channels. Journal of Pesticide Sciences, 2007, 32, 278-280.	1.4	2
42	Proinsecticide candidates N-(5-methyl-2-oxo-1,3-dioxol-4-yl)methyl derivatives of imidacloprid and 1-chlorothiazolylmethyl-2-nitroimino-imidazolidine. Bioorganic and Medicinal Chemistry Letters, 2007, 17, 4500-4503.	2.2	13
43	A hypothesis to account for the selective and diverse actions of neonicotinoid insecticides at their molecular targets, nicotinic acetylcholine receptors: catch and release in hydrogen bond networks. Invertebrate Neuroscience, 2007, 7, 47-51.	1.8	22
44	Actions of imidacloprid, clothianidin and related neonicotinoids on nicotinic acetylcholine receptors of American cockroach neurons and their relationships with insecticidal potency. Journal of Pesticide Sciences, 2006, 31, 35-40.	1.4	45
45	Neonicotinoid insecticides display partial and super agonist actions on native insect nicotinic acetylcholine receptors. Journal of Neurochemistry, 2006, 99, 608-615.	3.9	127
46	Isolation and identification of histamine-release inhibitors from Pistacia weinmannifolia J. Pisson ex. Franch. Journal of Natural Medicines, 2006, 60, 138-140.	2.3	7
47	Role in the Selectivity of Neonicotinoids of Insect-Specific Basic Residues in Loop D of the Nicotinic Acetylcholine Receptor Agonist Binding Site. Molecular Pharmacology, 2006, 70, 1255-1263.	2.3	114
48	Differential blocking actions of 4′-ethynyl-4-n-propylbicycloorthobenzoate (EBOB) and γ-hexachlorocyclohexane (γ-HCH) on γ-aminobutyric acid- and glutamate-induced responses of American cockroach neurons. Invertebrate Neuroscience, 2005, 5, 157-164.	1.8	28
49	Neonicotinoids Show Selective and Diverse Actions on Their Nicotinic Receptor Targets: Electrophysiology, Molecular Biology, and Receptor Modeling Studies. Bioscience, Biotechnology and Biochemistry, 2005, 69, 1442-1452.	1.3	175
50	Insect-vertebrate chimeric nicotinic acetylcholine receptors identify a region, loop B to the N-terminus of the Drosophila Dα2 subunit, which contributes to neonicotinoid sensitivity. Neuroscience Letters, 2005, 385, 168-172.	2.1	37
51	Super Agonist Actions of Clothianidin and Related Compounds on the SADβ2 Nicotinic Acetylcholine Receptor Expressed inXenopus laevisOocytes. Bioscience, Biotechnology and Biochemistry, 2004, 68, 761-763.	1.3	57
52	Action of nereistoxin on recombinant neuronal nicotinic acetylcholine receptors expressed in Xenopus laevis oocytes. Invertebrate Neuroscience, 2003, 5, 29-35.	1.8	23
53	Diverse actions of neonicotinoids on chicken α7, α4β2 and Drosophila–chicken SADβ2 and ALSβ2 hybrid nicotinic acetylcholine receptors expressed in Xenopus laevis oocytes. Neuropharmacology, 2003, 45, 133-144.	4.1	102
54	Insecticidal and Neural Activities of Candidate Photoaffinity Probes for Neonicotinoid Binding Sites. Bioscience, Biotechnology and Biochemistry, 2001, 65, 1534-1541.	1.3	11

#	Article	IF	CITATIONS
55	Role of loop D of the $\hat{1}\pm7$ nicotinic acetylcholine receptor in its interaction with the insecticide imidacloprid and related neonicotinoids. British Journal of Pharmacology, 2000, 130, 981-986.	5.4	66
56	PREFERENTIAL INHIBITION OF NUCLEOSOME ASSEMBLY BY ULTRAVIOLETâ€INDUCED (6â€4)PHOTOPRODUCTS. Photochemistry and Photobiology, 1995, 61, 459-462.	2.5	11
57	SIMULTANEOUS ESTABLISHMENT OF MONOCLONAL ANTIBODIES SPECIFIC FOR EITHER CYCLOBUTANE PYRIMIDINE DIMER OR (6â€4)PHOTOPRODUCT FROM THE SAME MOUSE IMMUNIZED WITH ULTRAVIOLETâ€IRRADIATED DNA. Photochemistry and Photobiology, 1991, 54, 225-232.	2.5	413