

Antonio Cigliano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5462633/publications.pdf>

Version: 2024-02-01

67
papers

3,266
citations

185998

28
h-index

155451

55
g-index

76
all docs

76
docs citations

76
times ranked

5169
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | TAZ is indispensable for c-MYC-induced hepatocarcinogenesis. <i>Journal of Hepatology</i> , 2022, 76, 123-134. | 1.8 | 28 |
| 2 | Nuclear localization dictates hepatocarcinogenesis suppression by glycine N-methyltransferase. <i>Translational Oncology</i> , 2022, 15, 101239. | 1.7 | 4 |
| 3 | RASSF1A independence and early galectin-1 upregulation in PIK3CA-induced hepatocarcinogenesis: new therapeutic venues. <i>Molecular Oncology</i> , 2022, 16, 1091-1118. | 2.1 | 8 |
| 4 | CD90 is regulated by notch1 and hallmarks a more aggressive intrahepatic cholangiocarcinoma phenotype. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022, 41, 65. | 3.5 | 7 |
| 5 | The Hippo pathway effector TAZ induces intrahepatic cholangiocarcinoma in mice and is ubiquitously activated in the human disease. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022, 41, . | 3.5 | 10 |
| 6 | Cabozantinib-based combination therapy for the treatment of hepatocellular carcinoma. <i>Gut</i> , 2021, 70, 1746-1757. | 6.1 | 60 |
| 7 | Distinct and Overlapping Roles of Hippo Effectors YAP and TAZ During Human and Mouse Hepatocarcinogenesis. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2021, 11, 1095-1117. | 2.3 | 21 |
| 8 | Current challenges to underpinning the genetic basis for cholangiocarcinoma. <i>Expert Review of Gastroenterology and Hepatology</i> , 2021, 15, 511-526. | 1.4 | 3 |
| 9 | Overexpression of Mothers Against Decapentaplegic Homolog 7 Activates the Yes-Associated Protein/NOTCH Cascade and Promotes Liver Carcinogenesis in Mice and Humans. <i>Hepatology</i> , 2021, 74, 248-263. | 3.6 | 22 |
| 10 | Cholesterol biosynthesis supports the growth of hepatocarcinoma lesions depleted of fatty acid synthase in mice and humans. <i>Gut</i> , 2020, 69, 177-186. | 6.1 | 121 |
| 11 | Inhibition of MELK Protooncogene as an Innovative Treatment for Intrahepatic Cholangiocarcinoma. <i>Medicina (Lithuania)</i> , 2020, 56, 1. | 0.8 | 13 |
| 12 | Pivotal Role of Fatty Acid Synthase in c-MYC Driven Hepatocarcinogenesis. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8467. | 1.8 | 20 |
| 13 | Transcriptomic and Proteomic Analysis of Clear Cell Foci (CCF) in the Human Non-Cirrhotic Liver Identifies Several Differentially Expressed Genes and Proteins with Functions in Cancer Cell Biology and Glycogen Metabolism. <i>Molecules</i> , 2020, 25, 4141. | 1.7 | 3 |
| 14 | Crenigacestat, a selective NOTCH1 inhibitor, reduces intrahepatic cholangiocarcinoma progression by blocking VEGFA/DLL4/MMP13 axis. <i>Cell Death and Differentiation</i> , 2020, 27, 2330-2343. | 5.0 | 39 |
| 15 | The Hippo Effector Transcriptional Coactivator with PDZ-Binding Motif Cooperates with Oncogenic β -Catenin to Induce Hepatoblastoma Development in Mice and Humans. <i>American Journal of Pathology</i> , 2020, 190, 1397-1413. | 1.9 | 13 |
| 16 | Combined CDK4/6 and Pan-mTOR Inhibition Is Synergistic Against Intrahepatic Cholangiocarcinoma. <i>Clinical Cancer Research</i> , 2019, 25, 403-413. | 3.2 | 56 |
| 17 | SNAI1 Promotes the Cholangiocellular Phenotype, but not Epithelial-Mesenchymal Transition, in a Murine Hepatocellular Carcinoma Model. <i>Cancer Research</i> , 2019, 79, 5563-5574. | 0.4 | 12 |
| 18 | Modification of the base excision repair enzyme MBD4 by the small ubiquitin-like molecule SUMO1. <i>DNA Repair</i> , 2019, 82, 102687. | 1.3 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Frizzled-10 Extracellular Vesicles Plasma Concentration Is Associated with Tumoral Progression in Patients with Colorectal and Gastric Cancer. <i>Journal of Oncology</i> , 2019, 2019, 1-12. | 0.6 | 24 |
| 20 | Loss of Fbxw7 synergizes with activated Akt signaling to promote c-Myc dependent cholangiocarcinogenesis. <i>Journal of Hepatology</i> , 2019, 71, 742-752. | 1.8 | 44 |
| 21 | The mTORC2-Akt1 Cascade Is Crucial for c-Myc to Promote Hepatocarcinogenesis in Mice and Humans. <i>Hepatology</i> , 2019, 70, 1600-1613. | 3.6 | 70 |
| 22 | Functional role of SGK3 in PI3K/Pten driven liver tumor development. <i>BMC Cancer</i> , 2019, 19, 343. | 1.1 | 17 |
| 23 | MEK inhibition suppresses K-Ras wild-type cholangiocarcinoma in vitro and in vivo via inhibiting cell proliferation and modulating tumor microenvironment. <i>Cell Death and Disease</i> , 2019, 10, 120. | 2.7 | 10 |
| 24 | TEA Domain Transcription Factor 4 Is the Major Mediator of Yes-Associated Protein Oncogenic Activity in Mouse and Human Hepatoblastoma. <i>American Journal of Pathology</i> , 2019, 189, 1077-1090. | 1.9 | 25 |
| 25 | Pathogenetic, Prognostic, and Therapeutic Role of Fatty Acid Synthase in Human Hepatocellular Carcinoma. <i>Frontiers in Oncology</i> , 2019, 9, 1412. | 1.3 | 44 |
| 26 | TGF- β 2 as Multifaceted Orchestrator in HCC Progression: Signaling, EMT, Immune Microenvironment, and Novel Therapeutic Perspectives. <i>Seminars in Liver Disease</i> , 2019, 39, 053-069. | 1.8 | 78 |
| 27 | MicroRNA-203 impacts on the growth, aggressiveness and prognosis of hepatocellular carcinoma by targeting <i>MAT2A</i> and <i>MAT2B</i> genes. <i>Oncotarget</i> , 2019, 10, 2835-2854. | 0.8 | 18 |
| 28 | A novel preclinical model of cholangiocarcinoma based on human aberrant FBXW7 expression.. <i>Journal of Clinical Oncology</i> , 2019, 37, e15624-e15624. | 0.8 | 0 |
| 29 | Hippo Cascade Controls Lineage Commitment of Liver Tumors in Mice and Humans. <i>American Journal of Pathology</i> , 2018, 188, 995-1006. | 1.9 | 29 |
| 30 | Efficacy of MEK inhibition in a K-Ras-driven cholangiocarcinoma preclinical model. <i>Cell Death and Disease</i> , 2018, 9, 31. | 2.7 | 23 |
| 31 | Loss of Pten synergizes with c-Met to promote hepatocellular carcinoma development via mTORC2 pathway. <i>Experimental and Molecular Medicine</i> , 2018, 50, e417-e417. | 3.2 | 39 |
| 32 | Oncogenic potential of N-terminal deletion and S45Y mutant β -catenin in promoting hepatocellular carcinoma development in mice. <i>BMC Cancer</i> , 2018, 18, 1093. | 1.1 | 17 |
| 33 | Focal adhesion kinase activation limits efficacy of Dasatinib in c-Myc driven hepatocellular carcinoma. <i>Cancer Medicine</i> , 2018, 7, 6170-6181. | 1.3 | 11 |
| 34 | Oncogene-dependent addiction to carbohydrate-responsive element binding protein in hepatocellular carcinoma. <i>Cell Cycle</i> , 2018, 17, 1496-1512. | 1.3 | 14 |
| 35 | Both <i>de novo</i> synthesized and exogenous fatty acids support the growth of hepatocellular carcinoma cells. <i>Liver International</i> , 2017, 37, 80-89. | 1.9 | 60 |
| 36 | Oncogene dependent requirement of fatty acid synthase in hepatocellular carcinoma. <i>Cell Cycle</i> , 2017, 16, 499-507. | 1.3 | 45 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Role of the Notch signaling in cholangiocarcinoma. <i>Expert Opinion on Therapeutic Targets</i> , 2017, 21, 471-483. | 1.5 | 27 |
| 38 | A functional mammalian target of rapamycin complex 1 signaling is indispensable for c-Myc-driven hepatocarcinogenesis. <i>Hepatology</i> , 2017, 66, 167-181. | 3.6 | 119 |
| 39 | Pan-mTOR inhibitor MLN0128 is effective against intrahepatic cholangiocarcinoma in mice. <i>Journal of Hepatology</i> , 2017, 67, 1194-1203. | 1.8 | 77 |
| 40 | Tankyrase inhibitors suppress hepatocellular carcinoma cell growth via modulating the Hippo cascade. <i>PLoS ONE</i> , 2017, 12, e0184068. | 1.1 | 35 |
| 41 | Deregulated c-Myc requires a functional HSF1 for experimental and human hepatocarcinogenesis. <i>Oncotarget</i> , 2017, 8, 90638-90650. | 0.8 | 17 |
| 42 | Inhibition of HSF1 suppresses the growth of hepatocarcinoma cell lines <i>in vitro</i> and AKT-driven hepatocarcinogenesis in mice. <i>Oncotarget</i> , 2017, 8, 54149-54159. | 0.8 | 24 |
| 43 | Central role of mTORC1 downstream of YAP/TAZ in hepatoblastoma development. <i>Oncotarget</i> , 2017, 8, 73433-73447. | 0.8 | 26 |
| 44 | Hepatocellular glycogenotic foci after combined intraportal pancreatic islet transplantation and knockout of the carbohydrate responsive element binding protein in diabetic mice. <i>Oncotarget</i> , 2017, 8, 104315-104329. | 0.8 | 7 |
| 45 | The Epidermal Growth Factor Receptor (EGFR) Inhibitor Gefitinib Reduces but Does Not Prevent Tumorigenesis in Chemical and Hormonal Induced Hepatocarcinogenesis Rat Models. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1618. | 1.8 | 4 |
| 46 | PI3K/AKT/mTOR-dependent stabilization of oncogenic far-upstream element binding proteins in hepatocellular carcinoma cells. <i>Hepatology</i> , 2016, 63, 813-826. | 3.6 | 52 |
| 47 | Quantification of liver proton-density fat fraction in 7.1T preclinical MR systems: Impact of the fitting technique. <i>Journal of Magnetic Resonance Imaging</i> , 2016, 44, 1425-1431. | 1.9 | 0 |
| 48 | Jagged 1 is a major Notch ligand along cholangiocarcinoma development in mice and humans. <i>Oncogenesis</i> , 2016, 5, e274-e274. | 2.1 | 28 |
| 49 | Co-activation of AKT and c-Met triggers rapid hepatocellular carcinoma development via the mTORC1/FASN pathway in mice. <i>Scientific Reports</i> , 2016, 6, 20484. | 1.6 | 100 |
| 50 | Differential requirement for de novo lipogenesis in cholangiocarcinoma and hepatocellular carcinoma of mice and humans. <i>Hepatology</i> , 2016, 63, 1900-1913. | 3.6 | 82 |
| 51 | Inactivation of fatty acid synthase impairs hepatocarcinogenesis driven by AKT in mice and humans. <i>Journal of Hepatology</i> , 2016, 64, 333-341. | 1.8 | 115 |
| 52 | 4EBP1/eIF4E and p70S6K/RPS6 axes play critical and distinct roles in hepatocarcinogenesis driven by AKT and N-Ras proto-oncogenes in mice. <i>Hepatology</i> , 2015, 61, 200-213. | 3.6 | 63 |
| 53 | Co-activation of PIK3CA and Yap promotes development of hepatocellular and cholangiocellular tumors in mouse and human liver. <i>Oncotarget</i> , 2015, 6, 10102-10115. | 0.8 | 61 |
| 54 | SKP2 cooperates with N-Ras or AKT to induce liver tumor development in mice. <i>Oncotarget</i> , 2015, 6, 2222-2234. | 0.8 | 27 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 55 | PI3K/AKT/mTOR pathway plays a major pathogenetic role in glycogen accumulation and tumor development in renal distal tubules of rats and men. <i>Oncotarget</i> , 2015, 6, 13036-13048. | 0.8 | 42 |
| 56 | Activation of β -Catenin and Yap1 in Human Hepatoblastoma and Induction of Hepatocarcinogenesis in Mice. <i>Gastroenterology</i> , 2014, 147, 690-701. | 0.6 | 249 |
| 57 | Molecular and metabolic changes in human liver clear cell foci resemble the alterations occurring in rat hepatocarcinogenesis. <i>Journal of Hepatology</i> , 2013, 58, 1147-1156. | 1.8 | 26 |
| 58 | Functional crosstalk between AKT/mTOR and Ras/MAPK pathways in hepatocarcinogenesis: Implications for the treatment of human liver cancer. <i>Cell Cycle</i> , 2013, 12, 1999-2010. | 1.3 | 82 |
| 59 | Association between Human Plasma Chondroitin Sulfate Isomers and Carotid Atherosclerotic Plaques. <i>Biochemistry Research International</i> , 2012, 2012, 1-6. | 1.5 | 13 |
| 60 | Fine Structure of Glycosaminoglycans from Fresh and Decellularized Porcine Cardiac Valves and Pericardium. <i>Biochemistry Research International</i> , 2012, 2012, 1-10. | 1.5 | 51 |
| 61 | Inactivation of Spry2 accelerates AKT-driven hepatocarcinogenesis via activation of MAPK and PKM2 pathways. <i>Journal of Hepatology</i> , 2012, 57, 577-583. | 1.8 | 45 |
| 62 | Thymine DNA Glycosylase Is Essential for Active DNA Demethylation by Linked Deamination-Base Excision Repair. <i>Cell</i> , 2011, 146, 67-79. | 13.5 | 700 |
| 63 | Differential distribution of structural components and hydration in aortic and pulmonary heart valve conduits: Impact of detergent-based cell removal. <i>Acta Biomaterialia</i> , 2010, 6, 4675-4688. | 4.1 | 24 |
| 64 | Plasma levels of C-reactive protein, leptin and glycosaminoglycans during spontaneous menstrual cycle: differences between ovulatory and anovulatory cycles. <i>Archives of Gynecology and Obstetrics</i> , 2010, 282, 207-213. | 0.8 | 35 |
| 65 | Evaluation of human serum albumin sulfhydryl groups oxidation in plasma and atherosclerotic plaque extracts. <i>Journal of Biological Research (Italy)</i> , 2010, 83, . | 0.0 | 0 |
| 66 | Glycosaminoglycans and Fabry's disease. <i>Journal of Biological Research (Italy)</i> , 2010, 83, . | 0.0 | 0 |
| 67 | A proteomic approach to differentiate histologically classified stable and unstable plaques from human carotid arteries. <i>Atherosclerosis</i> , 2009, 203, 112-118. | 0.4 | 120 |