

Ramit Ravona-Springer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5477836/publications.pdf>

Version: 2024-02-01

99
papers

1,007
citations

516561

16
h-index

501076

28
g-index

118
all docs

118
docs citations

118
times ranked

1702
citing authors

#	ARTICLE	IF	CITATIONS
1	The effects of cardiovascular risk factors on cognitive compromise. <i>Dialogues in Clinical Neuroscience</i> , 2009, 11, 201-212.	1.8	89
2	Changes in Glycemic Control are Associated with Changes in Cognition in Non-Diabetic Elderly. <i>Journal of Alzheimer's Disease</i> , 2012, 30, 299-309.	1.2	65
3	Diabetes Is Associated with Increased Rate of Cognitive Decline in Questionably Demented Elderly. <i>Dementia and Geriatric Cognitive Disorders</i> , 2010, 29, 68-74.	0.7	55
4	The Association of Duration of Type 2 Diabetes with Cognitive Performance is Modulated by Long-Term Glycemic Control. <i>American Journal of Geriatric Psychiatry</i> , 2014, 22, 1055-1059.	0.6	54
5	Trajectories in Glycemic Control over Time Are Associated with Cognitive Performance in Elderly Subjects with Type 2 Diabetes. <i>PLoS ONE</i> , 2014, 9, e97384.	1.1	53
6	The Israel Diabetes and Cognitive Decline (IDCD) study: Design and baseline characteristics. <i>Alzheimer's and Dementia</i> , 2014, 10, 769-778.	0.4	52
7	The association of diabetes and dementia and possible implications for nondiabetic populations. <i>Expert Review of Neurotherapeutics</i> , 2011, 11, 1609-1617.	1.4	40
8	A Systematic Review and Meta-Analysis of the Association between Helicobacter pylori Infection and Dementia. <i>Journal of Alzheimer's Disease</i> , 2016, 52, 1431-1442.	1.2	29
9	Younger Age at Crisis Following Parental Death in Male Children and Adolescents Is Associated With Higher Risk for Dementia at Old Age. <i>Alzheimer Disease and Associated Disorders</i> , 2012, 26, 68-73.	0.6	27
10	Hemoglobin A1c Variability Predicts Symptoms of Depression in Elderly Individuals With Type 2 Diabetes. <i>Diabetes Care</i> , 2017, 40, 1187-1193.	4.3	27
11	Haptoglobin 1-1 Genotype Is Associated With Poorer Cognitive Functioning in the Elderly With Type 2 Diabetes. <i>Diabetes Care</i> , 2013, 36, 3139-3145.	4.3	26
12	Long-term Variability in Glycemic Control Is Associated With White Matter Hyperintensities in APOE4 Genotype Carriers With Type 2 Diabetes. <i>Diabetes Care</i> , 2016, 39, 1056-1059.	4.3	24
13	Impaired Cerebral Hemodynamics and Cognitive Performance in Patients with Atherothrombotic Disease. <i>Journal of Alzheimer's Disease</i> , 2015, 46, 137-144.	1.2	22
14	Waist circumference is correlated with poorer cognition in elderly type 2 diabetes women. <i>Alzheimer's and Dementia</i> , 2016, 12, 925-929.	0.4	22
15	Computerized Cognitive Training for Older Adults at Higher Dementia Risk due to Diabetes: Findings From a Randomized Controlled Trial. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2020, 75, 747-754.	1.7	20
16	The Role of Cardiovascular Risk Factors in Alzheimer's Disease. <i>CNS Spectrums</i> , 2003, 8, 824-831.	0.7	18
17	Decreased Motor Function Is Associated with Poorer Cognitive Function in Elderly with Type 2 Diabetes. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2014, 4, 103-112.	0.6	18
18	Association of the Haptoglobin Gene Polymorphism With Cognitive Function and Decline in Elderly African American Adults With Type 2 Diabetes. <i>JAMA Network Open</i> , 2018, 1, e184458.	2.8	17

#	ARTICLE	IF	CITATIONS
19	The associations between objective and subjective health among older adults with type 2 diabetes: The moderating role of personality. <i>Journal of Psychosomatic Research</i> , 2019, 117, 41-47.	1.2	17
20	Shorter Adult Height is Associated with Poorer Cognitive Performance in Elderly Men with Type II Diabetes. <i>Journal of Alzheimer's Disease</i> , 2015, 44, 927-935.	1.2	16
21	Exposure to the Holocaust and World War II Concentration Camps during Late Adolescence and Adulthood is not Associated with Increased Risk for Dementia at Old Age. <i>Journal of Alzheimer's Disease</i> , 2011, 23, 709-716.	1.2	15
22	The ApoE4 genotype modifies the relationship of long-term glycemic control with cognitive functioning in elderly with type 2 diabetes. <i>European Neuropsychopharmacology</i> , 2014, 24, 1303-1308.	0.3	15
23	Glycemic control, inflammation, and cognitive function in older patients with type 2 diabetes. <i>International Journal of Geriatric Psychiatry</i> , 2015, 30, 1093-1100.	1.3	15
24	The <i>CADM2</i> gene is associated with processing speed performance – evidence among elderly with type 2 diabetes. <i>World Journal of Biological Psychiatry</i> , 2019, 20, 577-583.	1.3	15
25	Haptoglobin genotype modulates the relationships of glycaemic control with cognitive function in elderly individuals with type 2 diabetes. <i>Diabetologia</i> , 2015, 58, 736-744.	2.9	14
26	Potential contribution of the Alzheimer's disease risk locus BIN1 to episodic memory performance in cognitively normal Type 2 diabetes elderly. <i>European Neuropsychopharmacology</i> , 2016, 26, 787-795.	0.3	14
27	Is the distinction between Alzheimer's disease and vascular dementia possible and relevant?. <i>Dialogues in Clinical Neuroscience</i> , 2003, 5, 7-15.	1.8	14
28	Haptoglobin 1-1 Genotype Modulates the Association of Glycemic Control With Hippocampal Volume in Elderly Individuals With Type 2 Diabetes. <i>Diabetes</i> , 2017, 66, 2927-2932.	0.3	13
29	Considerations in psychotropic treatments in dementia – can polypharmacy be avoided?. <i>International Journal of Neuropsychopharmacology</i> , 2014, 17, 1107-1117.	1.0	12
30	Depressive Symptoms Are Associated with Cognitive Function in the Elderly with Type 2 Diabetes. <i>Journal of Alzheimer's Disease</i> , 2018, 65, 683-692.	1.2	12
31	Age Modulates the Association of Caffeine Intake With Cognition and With Gray Matter in Elderly Diabetics. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2019, 74, 683-688.	1.7	12
32	Higher BMI is associated with smaller regional brain volume in older adults with type 2 diabetes. <i>Diabetologia</i> , 2020, 63, 2446-2451.	2.9	12
33	Long-term trajectories of BMI predict carotid stiffness and plaque volume in type 2 diabetes older adults: a cohort study. <i>Cardiovascular Diabetology</i> , 2020, 19, 138.	2.7	12
34	Increase in Number of Depression Symptoms Over Time is Related to Worse Cognitive Outcomes in Older Adults With Type 2 Diabetes. <i>American Journal of Geriatric Psychiatry</i> , 2021, 29, 1-11.	0.6	12
35	The Association of Age With Rate of Cognitive Decline in Elderly Individuals Residing in Supporting Care Facilities. <i>Alzheimer Disease and Associated Disorders</i> , 2011, 25, 312-316.	0.6	11
36	The Israel Registry for Alzheimer's Prevention (IRAP) Study: Design and Baseline Characteristics. <i>Journal of Alzheimer's Disease</i> , 2020, 78, 777-788.	1.2	11

#	ARTICLE	IF	CITATIONS
37	Deterioration in Motor Function Over Time in Older Adults With Type 2 Diabetes is Associated with Accelerated Cognitive Decline. <i>Endocrine Practice</i> , 2020, 26, 1143-1152.	1.1	11
38	Specific Dimensions of Depression Have Different Associations With Cognitive Decline in Older Adults With Type 2 Diabetes. <i>Diabetes Care</i> , 2021, 44, 655-662.	4.3	10
39	Repetitive Thinking as a Psychological Cognitive Style in Midlife Is Associated with Lower Risk for Dementia Three Decades Later. <i>Dementia and Geriatric Cognitive Disorders</i> , 2009, 28, 513-520.	0.7	9
40	Statin Use is Associated with Better Cognitive Function in Elderly with Type 2 Diabetes. <i>Journal of Alzheimer's Disease</i> , 2015, 47, 55-59.	1.2	9
41	Psychosis as a Treatment Target in Dementia: A Roadmap for Designing Interventions. <i>Journal of Alzheimer's Disease</i> , 2022, 88, 1203-1228.	1.2	7
42	Computerized cognitive training for older diabetic adults at risk of dementia: Study protocol for a randomized controlled trial. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2017, 3, 636-650.	1.8	6
43	Differences in Semantic Memory Encoding Strategies in Young, Healthy Old and MCI Patients. <i>Frontiers in Aging Neuroscience</i> , 2019, 11, 306.	1.7	6
44	Sleep Monitoring Using WatchPAT Device to Predict Recurrence of Major Depression in Patients at High Risk for Major Depression Disorder Recurrence: A Case Report. <i>Frontiers in Psychiatry</i> , 2021, 12, 572660.	1.3	6
45	TCF7L2 polymorphisms are associated with amygdalar volume in elderly individuals with Type 2 Diabetes. <i>Scientific Reports</i> , 2019, 9, 15818.	1.6	5
46	Personality traits and cognitive function in old-adults with type-2 diabetes. <i>Aging and Mental Health</i> , 2019, 23, 1317-1325.	1.5	5
47	Ethnicity/culture modulates the relationships of the haptoglobin (Hp) 1-1 phenotype with cognitive function in older individuals with type 2 diabetes. <i>International Journal of Geriatric Psychiatry</i> , 2016, 31, 494-501.	1.3	4
48	Association of the CD2AP locus with cognitive functioning among middle-aged individuals with a family history of Alzheimer's disease. <i>Neurobiology of Aging</i> , 2021, 101, 50-56.	1.5	4
49	Satisfaction with Current Status at Work and Lack of Motivation to Improve It During Midlife is Associated with Increased Risk for Dementia in Subjects who Survived Thirty-Seven Years Later. <i>Journal of Alzheimer's Disease</i> , 2013, 36, 769-780.	1.2	3
50	C-reactive protein in midlife is associated with depressive symptoms two decades later among men with coronary heart disease. <i>Nordic Journal of Psychiatry</i> , 2020, 74, 226-233.	0.7	3
51	Vitamin E Intake Is Associated with Lower Brain Volume in Haptoglobin 1-1 Elderly with Type 2 Diabetes. <i>Journal of Alzheimer's Disease</i> , 2020, 74, 649-658.	1.2	2
52	Physical fitness mediates the association between age and cognition in healthy adults. <i>Aging Clinical and Experimental Research</i> , 2021, 33, 1359-1366.	1.4	2
53	Trajectories of depression symptoms over time differ by APOE4 genotype in older adults with type 2 diabetes. <i>International Journal of Geriatric Psychiatry</i> , 2021, 36, 1567-1575.	1.3	2
54	A spectrum of contributions of type 2 diabetes and related metabolic characteristics to dementia. <i>European Neuropsychopharmacology</i> , 2014, 24, 1945-1946.	0.3	1

#	ARTICLE	IF	CITATIONS
55	F3-01-04: SOCIOECONOMIC STATUS AND CHOLESTEROL LEVELS IN MIDLIFE AND RISK OF DEMENTIA 35 YEARS LATER. , 2014, 10, P203-P203.		1
56	O2-09-04: TRAJECTORIES IN GLYCEMIC CONTROL OVER TIME ARE ASSOCIATED WITH COGNITIVE PERFORMANCE IN ELDERLY SUBJECTS WITH TYPE 2 DIABETES. , 2014, 10, P184-P185.		1
57	Arterial Wall Function is Associated with Cognitive Performance Primarily in Elderly with Type 2 Diabetes. Journal of Alzheimer's Disease, 2015, 44, 687-693.	1.2	1
58	P2-252: Structural Brain Abnormalities Associated with Depression in Elderly with Type 2 Diabetes Differ by Haptoglobin Genotype. Alzheimer's and Dementia, 2016, 12, P722.	0.4	1
59	[P4-366]: DIETARY REDUCTION OF ADVANCED GLYCATION END PRODUCTS TO PREVENT COGNITIVE DECLINE IN HIGH-RISK DIABETICS: A PILOT RANDOMIZED TRIAL. Alzheimer's and Dementia, 2017, 13, P1432.	0.4	1
60	The Association of Depressive Symptoms With Brain Volume Is Stronger Among Diabetic Elderly Carriers of the Haptoglobin 1-1 Genotype Compared to Non-carriers. Frontiers in Endocrinology, 2019, 10, 68.	1.5	1
61	Trajectories of depression symptoms over time differ by APOE4 carriership in older adults with type 2 diabetes. Alzheimer's and Dementia, 2020, 16, e045304.	0.4	1
62	Effective cognitive screening tools for Alzheimer's disease in the primary care setting: the role of the visual paired associative learning task. International Psychogeriatrics, 2021, 33, 1111-1114.	0.6	1
63	Consumption of ultra-processed food and cognitive decline among older adults with type 2 diabetes. Alzheimer's and Dementia, 2021, 17, .	0.4	1
64	P4-239: THE ASSOCIATION OF THE TCF7L2 DIABETES SUSCEPTIBILITY GENE WITH BRAIN FUNCTION AMONG ELDERLY SUBJECTS WITH TYPE 2 DIABETES. , 2014, 10, P874-P874.		0
65	P3-343: ARTERIAL WALL FUNCTION AND COGNITIVE PERFORMANCE IN DIABETIC AND NON-DIABETIC ELDERLY SUBJECTS. , 2014, 10, P756-P756.		0
66	O3-07-05: DIABETES AND ALZHEIMER'S-ASSOCIATED GENES AFFECT DIFFERENTLY THE ASSOCIATION OF GLYCEMIC CONTROL WITH COGNITIVE FUNCTION. , 2014, 10, P222-P223.		0
67	O3-07-04: MIDLIFE RISK FACTORS FOR MORTALITY AND DEMENTIA AT VERY OLD AGE: THE ISRAEL ISCHEMIC HEART DISEASE STUDY. , 2014, 10, P222-P222.		0
68	P4-011: The Alzheimer's disease risk loci bin1 is associated with poorer episodic memory performance in cognitively normal type 2 diabetes elderly. , 2015, 11, P769-P770.		0
69	P1-266: Trajectories of glycemic control are associated with depressive symptoms beyond cognitive function in nondemented very elderly with type 2 diabetes. , 2015, 11, P457-P457.		0
70	P1-202: Haptoglobin genotype modulates the relationships of glycemic control with white matter hyperintensities in elderly with type 2 diabetes. , 2015, 11, P427-P427.		0
71	P3-250: Haptoglobin genotype modulates the relationships of glycaemic control with cognitive function in elderly individuals with type 2 diabetes. , 2015, 11, P726-P726.		0
72	P3-251: Ethnicity/culture modulates the relationships of the haptoglobin (Hp) 1-1 phenotype with cognitive function in elderly with type 2 diabetes. , 2015, 11, P727-P727.		0

#	ARTICLE	IF	CITATIONS
73	P1425: The Effects of Computerized Cognitive Training on Cognitive and Non-Cognitive Outcomes Among Older Adults with Diabetes: Protocol for a Double-Blind Randomized Controlled Trial. <i>Alzheimer's and Dementia</i> , 2016, 12, P598.	0.4	0
74	P3375: Neuropsychiatric Symptoms are Associated with Cognitive Function in Elderly with Type 2 Diabetes. <i>Alzheimer's and Dementia</i> , 2016, 12, P993.	0.4	0
75	P4-155: The Relationship of Depressive Symptoms with Cognitive Function is Different for Men and Women with Type 2 Diabetes. , 2016, 12, P1074-P1074.		0
76	O41206: Adherence to Type 2 Diabetes Medications is Associated with Better Cognitive Function. <i>Alzheimer's and Dementia</i> , 2016, 12, P365.	0.4	0
77	[P1137]: TYPE 2 DIABETES GENETIC RISK VARIANTS WITHIN <i>TCF7L2</i> ARE ASSOCIATED WITH SMALLER AMYGDALAR VOLUME AMONG DIABETIC ELDERLY PATIENTS. <i>Alzheimer's and Dementia</i> , 2017, 13, P294.	0.4	0
78	[P4340]: THE ISRAEL REGISTRY FOR ALZHEIMER'S PREVENTION (IRAP) STUDY OF MIDDLE-AGED OFFSPRING OF ALZHEIMER'S DISEASE PATIENTS WITH 20 YEARS OF EXQUISITE HEALTH CHARACTERIZATION. <i>Alzheimer's and Dementia</i> , 2017, 13, P1419.	0.4	0
79	O11402: THE HAPTOGLOBIN 1 GENOTYPE IS ASSOCIATED WITH COGNITIVE FUNCTION AND DECLINE IN AFRICAN AMERICAN PERSONS WITH TYPE 2 DIABETES: THE ACCORD-MIND STUDY. <i>Alzheimer's and Dementia</i> , 2018, 14, P257.	0.4	0
80	O30702: THE EFFECTS OF COMPUTERISED COGNITIVE TRAINING ON COGNITIVE OUTCOMES AND SELF-MANAGEMENT IN DIABETIC OLDER ADULTS AT RISK OF DEMENTIA: RESULTS FROM A DOUBLE-BLIND RCT. <i>Alzheimer's and Dementia</i> , 2018, 14, P1029.	0.4	0
81	P2250: ASSOCIATION OF STRUCTURAL RETINAL MARKERS WITH COGNITIVE FUNCTION IN ASYMPTOMATIC INDIVIDUALS AT HIGH RISK FOR ALZHEIMER'S DISEASE. <i>Alzheimer's and Dementia</i> , 2018, 14, P769.	0.4	0
82	P3627: PERSONALITY TRAITS AND COGNITIVE FUNCTION IN OLD ADULTS WITH TYPE 2 DIABETES. <i>Alzheimer's and Dementia</i> , 2018, 14, P1372.	0.4	0
83	P4592: RETINAL THICKNESS CHANGES IN ASYMPTOMATIC MIDDLE-AGED INDIVIDUALS AT HIGH RISK FOR ALZHEIMERS DISEASE. <i>Alzheimer's and Dementia</i> , 2019, 15, P1550.	0.4	0
84	Depression is more strongly associated with cognition in elderly women than men with type 2 diabetes. <i>International Psychogeriatrics</i> , 2019, 31, 591-595.	0.6	0
85	Long-term adiposity is associated with poorer cognitive performance in middle-aged adults at high risk of Alzheimer's disease. <i>Alzheimer's and Dementia</i> , 2020, 16, e041829.	0.4	0
86	HAPPCAP-AD (Human Application Combined Approach for Prevention of Alzheimer's Disease): A novel feasibility study of a personalized midlife intervention to prevent AD. <i>Alzheimer's and Dementia</i> , 2020, 16, e044234.	0.4	0
87	The Israel Registry for Alzheimer's Prevention (IRAP) study of middle-aged offspring of Alzheimer's disease patients. <i>Alzheimer's and Dementia</i> , 2020, 16, e045222.	0.4	0
88	The COMT rs4680 polymorphism is associated with rate of cognitive decline in older adults with type 2 diabetes. <i>Alzheimer's and Dementia</i> , 2020, 16, e045227.	0.4	0
89	Greater apathy is associated with faster rate of cognitive decline in older adults with type 2 diabetes. <i>Alzheimer's and Dementia</i> , 2020, 16, e045641.	0.4	0
90	Distinct Dimensions of Depression Are Associated With Different Brain-Related Biomarkers. <i>Diabetes Care</i> , 2022, , .	4.3	0

#	ARTICLE	IF	CITATIONS
91	The relationship of regional abdominal adiposity and adiposity-related factors with cognitive functioning among middle-aged individuals at high Alzheimer's dementia risk. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.4	0
92	Regional abdominal adiposity is associated with BMI-related brain regions in middle-aged adults at high risk of Alzheimer's dementia. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.4	0
93	Amyloid pathology, small-vessel disease, atrophy, and cognition in normal adults with type 2 diabetes. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.4	0
94	Lower motor function prior to the Covid-19 imposed lockdown predicts emotional distress in older adults with T2D. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.4	0
95	APOE ϵ 4 genotype is associated with thicker retinal layers in asymptomatic middle-aged adults at high Alzheimer's disease risk. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.4	0
96	Regional abdominal adiposity and related factors are associated with brain volumes and cognitive functioning in middle-aged adults at high AD risk. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.4	0
97	No evidence for association of Alzheimer's disease risk loci with the rate of cognitive decline among elderly with type 2 diabetes. <i>Alzheimer's and Dementia</i> , 2021, 17, e050081.	0.4	0
98	Gait speed is associated with changes in number of depression symptoms in older adults with type 2 diabetes. <i>Alzheimer's and Dementia</i> , 2021, 17, .	0.4	0
99	Current and long-term trajectories of high BMI are associated with poorer cognitive functioning in middle-aged adults at high Alzheimer's risk due to a maternal family history. <i>Alzheimer's and Dementia</i> , 2021, 17, e051464.	0.4	0