Ramit Ravona-Springer

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5477836/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Distinct Dimensions of Depression Are Associated With Different Brain-Related Biomarkers. Diabetes Care, 2022, , .	8.6	0
2	Psychosis as a Treatment Target in Dementia: A Roadmap for Designing Interventions. Journal of Alzheimer's Disease, 2022, 88, 1203-1228.	2.6	7
3	Physical fitness mediates the association between age and cognition in healthy adults. Aging Clinical and Experimental Research, 2021, 33, 1359-1366.	2.9	2
4	Increase in Number of Depression Symptoms Over Time is Related to Worse Cognitive Outcomes in Older Adults With Type 2 Diabetes. American Journal of Geriatric Psychiatry, 2021, 29, 1-11.	1.2	12
5	Association of the CD2AP locus with cognitive functioning among middle-aged individuals with a family history of Alzheimer's disease. Neurobiology of Aging, 2021, 101, 50-56.	3.1	4
6	Specific Dimensions of Depression Have Different Associations With Cognitive Decline in Older Adults With Type 2 Diabetes. Diabetes Care, 2021, 44, 655-662.	8.6	10
7	Effective cognitive screening tools for Alzheimer's disease in the primary care setting: the role of the visual paired associative learning task. International Psychogeriatrics, 2021, 33, 1111-1114.	1.0	1
8	Trajectories of depression symptoms over time differ by APOE4 genotype in older adults with type 2 diabetes. International Journal of Geriatric Psychiatry, 2021, 36, 1567-1575.	2.7	2
9	Sleep Monitoring Using WatchPAT Device to Predict Recurrence of Major Depression in Patients at High Risk for Major Depression Disorder Recurrence: A Case Report. Frontiers in Psychiatry, 2021, 12, 572660.	2.6	6
10	The relationship of regional abdominal adiposity and adiposityâ€related factors with cognitive functioning among middleâ€aged individuals at high Alzheimer's dementia risk. Alzheimer's and Dementia, 2021, 17, .	0.8	0
11	Regional abdominal adiposity is associated with BMIâ€related brain regions in middleâ€aged adults at high risk of Alzheimer's dementia. Alzheimer's and Dementia, 2021, 17, .	0.8	0
12	Amyloid pathology, smallâ€vessel disease, atrophy, and cognition in normal adults with type 2 diabetes. Alzheimer's and Dementia, 2021, 17, .	0.8	0
13	Lower motor function prior to the Covidâ€19â€imposed lockdown predicts emotional distress in older adults with T2D. Alzheimer's and Dementia, 2021, 17, .	0.8	Ο
14	Consumption of ultraâ€processed food and cognitive decline among older adults with typeâ€2 diabetes. Alzheimer's and Dementia, 2021, 17, .	0.8	1
15	APOE Æ4 genotype is associated with thicker retinal layers in asymptomatic middleâ€aged adults at high Alzheimer's disease risk. Alzheimer's and Dementia, 2021, 17, .	0.8	0
16	Regional abdominal adiposity and related factors are associated with brain volumes and cognitive functioning in middleâ€aged adults at high ADâ€risk. Alzheimer's and Dementia, 2021, 17, .	0.8	0
17	No evidence for association of Alzheimer's disease risk loci with the rate of cognitive decline among elderly with type 2 diabetes. Alzheimer's and Dementia, 2021, 17, e050081.	0.8	0
18	Gait speed is associated with changes in number of depression symptoms in older adults with type 2 diabetes. Alzheimer's and Dementia, 2021, 17, .	0.8	0

#	Article	IF	CITATIONS
19	Current and longâ€term trajectories of high BMI are associated with poorer cognitive functioning in middleâ€aged adults at high Alzheimer's risk due to a maternal family history. Alzheimer's and Dementia, 2021, 17, e051464.	0.8	0
20	Computerized Cognitive Training for Older Adults at Higher Dementia Risk due to Diabetes: Findings From a Randomized Controlled Trial. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2020, 75, 747-754.	3.6	20
21	C-reactive protein in midlife is associated with depressive symptoms two decades later among men with coronary heart disease. Nordic Journal of Psychiatry, 2020, 74, 226-233.	1.3	3
22	The Israel Registry for Alzheimer's Prevention (IRAP) Study: Design and Baseline Characteristics. Journal of Alzheimer's Disease, 2020, 78, 777-788.	2.6	11
23	Higher BMI is associated with smaller regional brain volume in older adults with type 2 diabetes. Diabetologia, 2020, 63, 2446-2451.	6.3	12
24	Long-term trajectories of BMI predict carotid stiffness and plaque volume in type 2 diabetes older adults: a cohort study. Cardiovascular Diabetology, 2020, 19, 138.	6.8	12
25	Longâ€ŧerm adiposity is associated with poorer cognitive performance in middleâ€aged adults at high risk of Alzheimer's disease. Alzheimer's and Dementia, 2020, 16, e041829.	0.8	0
26	HAPPCAPâ€AD (Humanâ€Application Combined Approach for Prevention of Alzheimer's Disease): A novel feasibility study of a personalized midlife intervention to prevent AD. Alzheimer's and Dementia, 2020, 16, e044234.	0.8	0
27	The Israel Registry for Alzheimer's Prevention (IRAP) study of middleâ€aged offspring of Alzheimer's disease patients. Alzheimer's and Dementia, 2020, 16, e045222.	0.8	0
28	The COMT rs4680 polymorphism is associated with rate of cognitive decline in older adults with type 2 diabetes. Alzheimer's and Dementia, 2020, 16, e045227.	0.8	0
29	Trajectories of depression symptoms over time differ by APOE4 carriership in older adults with type 2 diabetes. Alzheimer's and Dementia, 2020, 16, e045304.	0.8	1
30	Greater apathy is associated with faster rate of cognitive decline in older adults with type 2 diabetes. Alzheimer's and Dementia, 2020, 16, e045641.	0.8	0
31	Vitamin E Intake Is Associated with Lower Brain Volume in Haptoglobin 1-1 Elderly with Type 2 Diabetes. Journal of Alzheimer's Disease, 2020, 74, 649-658.	2.6	2
32	Deterioration in Motor Function Over Time in Older Adults With Type 2 Diabetes is Associated with Accelerated Cognitive Decline. Endocrine Practice, 2020, 26, 1143-1152.	2.1	11
33	Age Modulates the Association of Caffeine Intake With Cognition and With Gray Matter in Elderly Diabetics. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2019, 74, 683-688.	3.6	12
34	TCF7L2 polymorphisms are associated with amygdalar volume in elderly individuals with Type 2 Diabetes. Scientific Reports, 2019, 9, 15818.	3.3	5
35	The associations between objective and subjective health among older adults with type 2 diabetes: The moderating role of personality. Journal of Psychosomatic Research, 2019, 117, 41-47.	2.6	17
36	The Association of Depressive Symptoms With Brain Volume Is Stronger Among Diabetic Elderly Carriers of the Haptoglobin 1-1 Genotype Compared to Non-carriers. Frontiers in Endocrinology, 2019, 10, 68.	3.5	1

#	Article	IF	CITATIONS
37	P4â€592: RETINAL THICKNESS CHANGES IN ASYMPTOMATIC MIDDLEâ€AGED INDIVIDUALS AT HIGH RISK FOR ALZHEIMERS DISEASE. Alzheimer's and Dementia, 2019, 15, P1550.	0.8	0
38	Differences in Semantic Memory Encoding Strategies in Young, Healthy Old and MCI Patients. Frontiers in Aging Neuroscience, 2019, 11, 306.	3.4	6
39	Personality traits and cognitive function in old-adults with type-2 diabetes. Aging and Mental Health, 2019, 23, 1317-1325.	2.8	5
40	Depression is more strongly associated with cognition in elderly women than men with type 2 diabetes. International Psychogeriatrics, 2019, 31, 591-595.	1.0	0
41	The <i>CADM2 </i> gene is associated with processing speed performance – evidence among elderly with type 2 diabetes. World Journal of Biological Psychiatry, 2019, 20, 577-583.	2.6	15
42	O1â€14â€02: THE HAPTOGLOBIN 1â€1 GENOTYPE IS ASSOCIATED WITH COGNITIVE FUNCTION AND DECLINE IN AFRICAN AMERICAN PERSONS WITH TYPE 2 DIABETES: THE ACCORDâ€MIND STUDY. Alzheimer's and Dementia, 2018, 14, P257.	0.8	0
43	O3â€07â€02: THE EFFECTS OF COMPUTERISED COGNITIVE TRAINING ON COGNITIVE OUTCOMES AND SELFâ€MANAGEMENT IN DIABETIC OLDER ADULTS AT RISK OF DEMENTIA: RESULTS FROM A DOUBLEâ€BLIND RC Alzheimer's and Dementia, 2018, 14, P1029.	To.8	0
44	P2â€⊋50: ASSOCIATION OF STRUCTURAL RETINAL MARKERS WITH COGNITIVE FUNCTION IN ASYMPTOMATIC INDIVIDUALS AT HIGH RISK FOR ALZHEIMER'S DISEASE. Alzheimer's and Dementia, 2018, 14, P769.	0.8	0
45	Association of the Haptoglobin Gene Polymorphism With Cognitive Function and Decline in Elderly African American Adults With Type 2 Diabetes. JAMA Network Open, 2018, 1, e184458.	5.9	17
46	P3â€627: PERSONALITY TRAITS AND COGNITIVE FUNCTION IN OLDâ€ADULTS WITH TYPE 2 DIABETES. Alzheimer': and Dementia, 2018, 14, P1372.	^{\$} 0.8	0
47	Depressive Symptoms Are Associated with Cognitive Function in the Elderly with Type 2 Diabetes. Journal of Alzheimer's Disease, 2018, 65, 683-692.	2.6	12
48	Hemoglobin A1c Variability Predicts Symptoms of Depression in Elderly Individuals With Type 2 Diabetes. Diabetes Care, 2017, 40, 1187-1193.	8.6	27
49	Computerized cognitive training for older diabetic adults at risk of dementia: Study protocol for a randomized controlled trial. Alzheimer's and Dementia: Translational Research and Clinical Interventions, 2017, 3, 636-650.	3.7	6
50	[P1–137]: TYPE 2 DIABETES GENETIC RISK VARIANTS WITHIN <i>TCF7L2</i> ARE ASSOCIATED WITH SMALLER AMYGDALAR VOLUME AMONG DIABETIC ELDERLY PATIENTS. Alzheimer's and Dementia, 2017, 13, P294.	0.8	0
51	[P4–366]: DIETARY REDUCTION OF ADVANCED GLYCATION END PRODUCTS TO PREVENT COGNITIVE DECLINE IN HIGHâ€RISK DIABETICS: A PILOT RANDOMIZED TRIAL. Alzheimer's and Dementia, 2017, 13, P1432.	0.8	1
52	[P4–340]: THE ISRAEL REGISTRY FOR ALZHEIMER'S PREVENTION (IRAP) STUDY OF MIDDLEâ€AGED OFFSPRING ALZHEIMER'S DISEASE PATIENTS WITH 20 YEARS OF EXQUISITE HEALTH CHARACTERIZATION. Alzheimer's and Dementia, 2017, 13, P1419.	OF 0.8	0
53	Haptoglobin 1-1 Genotype Modulates the Association of Glycemic Control With Hippocampal Volume in Elderly Individuals With Type 2 Diabetes. Diabetes, 2017, 66, 2927-2932.	0.6	13
54	Ethnicity/culture modulates the relationships of the haptoglobin (Hp) 1â€1 phenotype with cognitive function in older individuals with type 2 diabetes. International Journal of Geriatric Psychiatry, 2016, 31, 494-501.	2.7	4

#	Article	IF	CITATIONS
55	P1â€425: The Effects of Computerized Cognitive Training on Cognitive and Nonâ€Cognitive Outcomes Among Older Adults with Diabetes: Protocol for a Doubleâ€Blind Randomized Controlled Trial. Alzheimer's and Dementia, 2016, 12, P598.	0.8	0
56	P2â€252: Structural Brain Abnormalities Associated with Depression in Elderly with Type 2 Diabetes Differ by Haptoglobin Genotype. Alzheimer's and Dementia, 2016, 12, P722.	0.8	1
57	P3â€375: Neuropsychiatric Symptoms are Associated with Cognitive Function in Elderly with Type 2 Diabetes. Alzheimer's and Dementia, 2016, 12, P993.	0.8	0
58	P4-155: The Relationship of Depressive Symptoms with Cognitive Function is Different for Men and Women with Type 2 Diabetes. , 2016, 12, P1074-P1074.		0
59	O4â€12â€06: Adherence to Type 2 Diabetes Medications is Associated with Better Cognitive Function. Alzheimer's and Dementia, 2016, 12, P365.	0.8	0
60	Long-term Variability in Glycemic Control Is Associated With White Matter Hyperintensities in APOE4 Genotype Carriers With Type 2 Diabetes. Diabetes Care, 2016, 39, 1056-1059.	8.6	24
61	Waist circumference is correlated with poorer cognition in elderly type 2 diabetes women. Alzheimer's and Dementia, 2016, 12, 925-929.	0.8	22
62	A Systematic Review and Meta-Analysis of the Association between Helicobacter pylori Infection and Dementia. Journal of Alzheimer's Disease, 2016, 52, 1431-1442.	2.6	29
63	Potential contribution of the Alzheimer׳s disease risk locus BIN1 to episodic memory performance in cognitively normal Type 2 diabetes elderly. European Neuropsychopharmacology, 2016, 26, 787-795.	0.7	14
64	P4-011: The Alzheimer's disease risk loci bin1 is associated with poorer episodic memory performance in cognitively normal type 2 diabetes elderly. , 2015, 11, P769-P770.		0
65	Statin Use is Associated with Better Cognitive Function in Elderly with Type 2 Diabetes. Journal of Alzheimer's Disease, 2015, 47, 55-59.	2.6	9
66	P1-266: Trajectories of glycemic control are associated with depressive symptoms beyond cognitive function in nondemented very elderly with type 2 diabetes. , 2015, 11, P457-P457.		0
67	P1-202: Haptoglobin genotype modulates the relationships of glycemic control with white matter hyperintensities in elderly with type 2 diabetes. , 2015, 11, P427-P427.		Ο
68	Impaired Cerebral Hemodynamics and Cognitive Performance in Patients with Atherothrombotic Disease. Journal of Alzheimer's Disease, 2015, 46, 137-144.	2.6	22
69	Shorter Adult Height is Associated with Poorer Cognitive Performance in Elderly Men with Type II Diabetes. Journal of Alzheimer's Disease, 2015, 44, 927-935.	2.6	16
70	P3-250: Haptoglobin genotype modulates the relationships of glycaemic control with cognitive function in elderly individuals with type 2 diabetes. , 2015, 11, P726-P726.		0
71	P3-251: Ethnicity/culture modulates the relationships of the haptoblobin (Hp) 1-1 phenotype with cognitive function in elderly with type 2 diabetes. , 2015, 11, P727-P727.		0
72	Haptoglobin genotype modulates the relationships of glycaemic control with cognitive function in elderly individuals with type 2 diabetes. Diabetologia, 2015, 58, 736-744.	6.3	14

#	Article	IF	CITATIONS
73	Glycemic control, inflammation, and cognitive function in older patients with type 2 diabetes. International Journal of Geriatric Psychiatry, 2015, 30, 1093-1100.	2.7	15
74	Arterial Wall Function is Associated with Cognitive Performance Primarily in Elderly with Type 2 Diabetes. Journal of Alzheimer's Disease, 2015, 44, 687-693.	2.6	1
75	Considerations in psychotropic treatments in dementia–Âcan polypharmacy be avoided?. International Journal of Neuropsychopharmacology, 2014, 17, 1107-1117.	2.1	12
76	A spectrum of contributions of type 2 diabetes and related metabolic characteristics to dementia. European Neuropsychopharmacology, 2014, 24, 1945-1946.	0.7	1
77	The Association of Duration of Type 2 Diabetes with Cognitive Performance is Modulated by Long-Term Glycemic Control. American Journal of Geriatric Psychiatry, 2014, 22, 1055-1059.	1.2	54
78	The ApoE4 genotype modifies the relationship of long-term glycemic control with cognitive functioning in elderly with type 2 diabetes. European Neuropsychopharmacology, 2014, 24, 1303-1308.	0.7	15
79	The Israel Diabetes and Cognitive Decline (IDCD) study: Design and baseline characteristics. Alzheimer's and Dementia, 2014, 10, 769-778.	0.8	52
80	Decreased Motor Function Is Associated with Poorer Cognitive Function in Elderly with Type 2 Diabetes. Dementia and Geriatric Cognitive Disorders Extra, 2014, 4, 103-112.	1.3	18
81	P4-239: THE ASSOCIATION OF THE TCF7L2 DIABETES SUSCEPTIBILITY GENE WITH BRAIN FUNCTION AMONG ELDERLY SUBJECTS WITH TYPE 2 DIABETES. , 2014, 10, P874-P874.		0
82	F3-01-04: SOCIOECONOMIC STATUS AND CHOLESTEROL LEVELS IN MIDLIFE AND RISK OF DEMENTIA 35 YEARS LATER. , 2014, 10, P203-P203.		1
83	O2-09-04: TRAJECTORIES IN GLYCEMIC CONTROL OVER TIME ARE ASSOCIATED WITH COGNITIVE PERFORMANCE IN ELDERLY SUBJECTS WITH TYPE 2 DIABETES. , 2014, 10, P184-P185.		1
84	P3-343: ARTERIAL WALL FUNCTION AND COGNITIVE PERFORMANCE IN DIABETIC AND NON-DIABETIC ELDERLY SUBJECTS. , 2014, 10, P756-P756.		0
85	O3-07-05: DIABETES AND ALZHEIMER'S-ASSOCIATED GENES AFFECT DIFFERENTLY THE ASSOCIATION OF GLYCEMIC CONTROL WITH COGNITIVE FUNCTION. , 2014, 10, P222-P223.		0
86	O3-07-04: MIDLIFE RISK FACTORS FOR MORTALITY AND DEMENTIA AT VERY OLD AGE: THE ISRAEL ISCHEMIC HEART DISEASE STUDY. , 2014, 10, P222-P222.		0
87	Trajectories in Glycemic Control over Time Are Associated with Cognitive Performance in Elderly Subjects with Type 2 Diabetes. PLoS ONE, 2014, 9, e97384.	2.5	53
88	Haptoglobin 1-1 Genotype Is Associated With Poorer Cognitive Functioning in the Elderly With Type 2 Diabetes. Diabetes Care, 2013, 36, 3139-3145.	8.6	26
89	Satisfaction with Current Status at Work and Lack of Motivation to Improve It During Midlife is Associated with Increased Risk for Dementia in Subjects who Survived Thirty-Seven Years Later. Journal of Alzheimer's Disease, 2013, 36, 769-780.	2.6	3
90	Younger Age at Crisis Following Parental Death in Male Children and Adolescents Is Associated With Higher Risk for Dementia at Old Age. Alzheimer Disease and Associated Disorders, 2012, 26, 68-73.	1.3	27

#	Article	IF	CITATIONS
91	Changes in Glycemic Control are Associated with Changes in Cognition in Non-Diabetic Elderly. Journal of Alzheimer's Disease, 2012, 30, 299-309.	2.6	65
92	Exposure to the Holocaust and World War II Concentration Camps during Late Adolescence and Adulthood is not Associated with Increased Risk for Dementia at Old Age. Journal of Alzheimer's Disease, 2011, 23, 709-716.	2.6	15
93	The Association of Age With Rate of Cognitive Decline in Elderly Individuals Residing in Supporting Care Facilities. Alzheimer Disease and Associated Disorders, 2011, 25, 312-316.	1.3	11
94	The association of diabetes and dementia and possible implications for nondiabetic populations. Expert Review of Neurotherapeutics, 2011, 11, 1609-1617.	2.8	40
95	Diabetes Is Associated with Increased Rate of Cognitive Decline in Questionably Demented Elderly. Dementia and Geriatric Cognitive Disorders, 2010, 29, 68-74.	1.5	55
96	Repetitive Thinking as a Psychological Cognitive Style in Midlife Is Associated with Lower Risk for Dementia Three Decades Later. Dementia and Geriatric Cognitive Disorders, 2009, 28, 513-520.	1.5	9
97	The effects of cardiovascular risk factors on cognitive compromise. Dialogues in Clinical Neuroscience, 2009, 11, 201-212.	3.7	89
98	The Role of Cardiovascular Risk Factors in Alzheimer's Disease. CNS Spectrums, 2003, 8, 824-831.	1.2	18
99	Is the distinction between Alzheimer's disease and vascular dementia possible and relevant?. Dialogues in Clinical Neuroscience, 2003, 5, 7-15.	3.7	14