

Gangyi Yang

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5486658/gangyi-yang-publications-by-citations.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

120
papers

1,853
citations

23
h-index

35
g-index

128
ext. papers

2,390
ext. citations

5.4
avg, IF

4.61
L-index

#	Paper	IF	Citations
120	Nesfatin-1 action in the brain increases insulin sensitivity through Akt/AMPK/TORC2 pathway in diet-induced insulin resistance. <i>Diabetes</i> , 2012 , 61, 1959-68	0.9	91
119	Circulating Sfrp5 is a signature of obesity-related metabolic disorders and is regulated by glucose and liraglutide in humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 290-8	5.6	86
118	Elevated circulating levels of irisin and the effect of metformin treatment in women with polycystic ovary syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 1485-93	5.6	69
117	Zinc- α -glycoprotein is associated with insulin resistance in humans and is regulated by hyperglycemia, hyperinsulinemia, or liraglutide administration: cross-sectional and interventional studies in normal subjects, insulin-resistant subjects, and subjects with newly diagnosed diabetes. <i>Diabetes Care</i> , 2013 , 36, 1074-82	14.6	67
116	The adipose triglyceride lipase, adiponectin and visfatin are downregulated by tumor necrosis factor-alpha (TNF-alpha) in vivo. <i>Cytokine</i> , 2009 , 45, 12-9	4	52
115	Overexpression of visfatin/PBEF/Nampt alters whole-body insulin sensitivity and lipid profile in rats. <i>Annals of Medicine</i> , 2009 , 41, 311-20	1.5	49
114	Serum C1q/TNF-related protein-3 (CTRP3) levels are decreased in obesity and hypertension and are negatively correlated with parameters of insulin resistance. <i>Diabetology and Metabolic Syndrome</i> , 2015 , 7, 33	5.6	43
113	Myonectin Predicts the Development of Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 139-147	5.6	42
112	Hypothalamic nesfatin-1/NUCB2 knockdown augments hepatic gluconeogenesis that is correlated with inhibition of mTOR-STAT3 signaling pathway in rats. <i>Diabetes</i> , 2014 , 63, 1234-47	0.9	42
111	Liraglutide prevents hypoadiponectinemia-induced insulin resistance and alterations of gene expression involved in glucose and lipid metabolism. <i>Molecular Medicine</i> , 2011 , 17, 1168-78	6.2	42
110	The effects of fibroblast growth factor-21 knockdown and over-expression on its signaling pathway and glucose-lipid metabolism in vitro. <i>Molecular and Cellular Endocrinology</i> , 2012 , 348, 21-6	4.4	41
109	DPP-4 Inhibitors Improve Diabetic Wound Healing via Direct and Indirect Promotion of Epithelial-Mesenchymal Transition and Reduction of Scarring. <i>Diabetes</i> , 2018 , 67, 518-531	0.9	38
108	The role of JAZF1 on lipid metabolism and related genes in vitro. <i>Metabolism: Clinical and Experimental</i> , 2011 , 60, 523-30	12.7	38
107	Circulating preptin levels in normal, impaired glucose tolerance, and type 2 diabetic subjects. <i>Annals of Medicine</i> , 2009 , 41, 52-6	1.5	38
106	Liraglutide increases FGF-21 activity and insulin sensitivity in high fat diet and adiponectin knockdown induced insulin resistance. <i>PLoS ONE</i> , 2012 , 7, e48392	3.7	36
105	Alteration of gut microbiota induced by DPP-4i treatment improves glucose homeostasis. <i>EBioMedicine</i> , 2019 , 44, 665-674	8.8	34
104	Kr β pel-like factor 14 increases insulin sensitivity through activation of PI3K/Akt signal pathway. <i>Cellular Signalling</i> , 2015 , 27, 2201-8	4.9	33

103	Sodium-Glucose Cotransporter 2 (SGLT2) Inhibitor Increases Circulating Zinc- α -Glycoprotein Levels in Patients with Type 2 Diabetes. <i>Scientific Reports</i> , 2016 , 6, 32887	4.9	30
102	LAMP3 regulates hepatic lipid metabolism through activating PI3K/Akt pathway. <i>Molecular and Cellular Endocrinology</i> , 2018 , 470, 160-167	4.4	26
101	Decreased circulating BMP-9 levels in patients with Type 2 diabetes is a signature of insulin resistance. <i>Clinical Science</i> , 2017 , 131, 239-246	6.5	25
100	Inhibition of Mitochondrial Calcium Overload by SIRT3 Prevents Obesity- or Age-Related Whitening of Brown Adipose Tissue. <i>Diabetes</i> , 2020 , 69, 165-180	0.9	25
99	Silencing of FGF-21 expression promotes hepatic gluconeogenesis and glycogenolysis by regulation of the STAT3-SOCS3 signal. <i>FEBS Journal</i> , 2014 , 281, 2136-47	5.7	25
98	Efficacy and safety of aldose reductase inhibitor for the treatment of diabetic cardiovascular autonomic neuropathy: systematic review and meta-analysis. <i>PLoS ONE</i> , 2014 , 9, e87096	3.7	25
97	Plasma Sfrp5 levels correlate with determinants of the metabolic syndrome in Chinese adults. <i>Diabetes/Metabolism Research and Reviews</i> , 2017 , 33, e2896	7.5	23
96	Osteoprotegerin Promotes Liver Steatosis by Targeting the ERK-PPAR- γ -CD36 Pathway. <i>Diabetes</i> , 2019 , 68, 1902-1914	0.9	23
95	Role of bone morphogenetic protein-9 in the regulation of glucose and lipid metabolism. <i>FASEB Journal</i> , 2019 , 33, 10077-10088	0.9	22
94	JAZF1 ameliorates age and diet-associated hepatic steatosis through SREBP-1c-dependent mechanism. <i>Cell Death and Disease</i> , 2018 , 9, 859	9.8	22
93	Duodenal GLP-1 signaling regulates hepatic glucose production through a PKC-dependent neurocircuitry. <i>Cell Death and Disease</i> , 2017 , 8, e2609	9.8	21
92	Activation of TRPV1 channel antagonizes diabetic nephropathy through inhibiting endoplasmic reticulum-mitochondria contact in podocytes. <i>Metabolism: Clinical and Experimental</i> , 2020 , 105, 154182	12.7	21
91	Overexpression of juxtaposed with another zinc finger gene 1 reduces proinflammatory cytokine release via inhibition of stress-activated protein kinases and nuclear factor- κ B. <i>FEBS Journal</i> , 2014 , 281, 3193-205	5.7	21
90	Circulating zinc- α -glycoprotein levels are low in newly diagnosed patients with metabolic syndrome and correlate with adiponectin. <i>Nutrition and Metabolism</i> , 2017 , 14, 53	4.6	20
89	Overexpression of JAZF1 protected ApoE-deficient mice from atherosclerosis by inhibiting hepatic cholesterol synthesis via CREB-dependent mechanisms. <i>International Journal of Cardiology</i> , 2014 , 177, 100-10	3.2	20
88	The natural logarithm of zinc- α -glycoprotein/HOMA-IR is a better predictor of insulin sensitivity than the product of triglycerides and glucose and the other lipid ratios. <i>Cytokine</i> , 2016 , 79, 96-102	4	20
87	Effects of sitagliptin on circulating zinc- α -glycoprotein levels in newly diagnosed type 2 diabetes patients: a randomized trial. <i>European Journal of Endocrinology</i> , 2016 , 174, 147-55	6.5	19
86	Glypican-4 is increased in human subjects with impaired glucose tolerance and decreased in patients with newly diagnosed type 2 diabetes. <i>Acta Diabetologica</i> , 2014 , 51, 981-90	3.9	19

85	Circulating Zinc- α -glycoprotein levels and Insulin Resistance in Polycystic Ovary Syndrome. <i>Scientific Reports</i> , 2016 , 6, 25934	4.9	19
84	Rictor positively regulates B cell receptor signaling by modulating actin reorganization via ezrin. <i>PLoS Biology</i> , 2017 , 15, e2001750	9.7	18
83	A novel role for the Krüppel-like factor 14 on macrophage inflammatory response and atherosclerosis development. <i>Cardiovascular Pathology</i> , 2017 , 27, 1-8	3.8	17
82	Efficacy of low-level light therapy for treatment of diabetic foot ulcer: A systematic review and meta-analysis of randomized controlled trials. <i>Diabetes Research and Clinical Practice</i> , 2018 , 143, 215-224	7.4	17
81	Transcutaneous oxygen pressure (TcPO ₂) a novel diagnostic tool for peripheral neuropathy in type 2 diabetes patients. <i>Diabetes Research and Clinical Practice</i> , 2014 , 105, 336-43	7.4	17
80	Short-term pioglitazone treatment prevents free fatty acid-induced hepatic insulin resistance in normal rats: possible role of the resistin and adiponectin. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 339, 1190-6	3.4	17
79	Efficacy and safety of linagliptin in Asian patients with type 2 diabetes mellitus inadequately controlled by metformin: A multinational 24-week, randomized clinical trial. <i>Journal of Diabetes</i> , 2016 , 8, 229-37	3.8	17
78	Deficiency of PKD2L1 (TRPP3) Exacerbates Pathological Cardiac Hypertrophy by Augmenting NCX1-Mediated Mitochondrial Calcium Overload. <i>Cell Reports</i> , 2018 , 24, 1639-1652	10.6	16
77	Effective gene delivery of shBMP-9 using polyethyleneimine-based core-shell nanoparticles in an animal model of insulin resistance. <i>Nanoscale</i> , 2019 , 11, 2008-2016	7.7	15
76	Deficiency of Mitochondrial Glycerol 3-Phosphate Dehydrogenase Contributes to Hepatic Steatosis. <i>Hepatology</i> , 2019 , 70, 84-97	11.2	15
75	Polyethylene glycol loxenatide injections added to metformin effectively improve glycemic control and exhibit favorable safety in type 2 diabetic patients. <i>Journal of Diabetes</i> , 2017 , 9, 158-167	3.8	14
74	Changes in whole metabolites after exenatide treatment in overweight/obese polycystic ovary syndrome patients. <i>Clinical Endocrinology</i> , 2019 , 91, 508-516	3.4	14
73	Circulating bone morphogenetic protein-9 in relation to metabolic syndrome and insulin resistance. <i>Scientific Reports</i> , 2017 , 7, 17529	4.9	14
72	Circulating C1q/TNF-related protein isoform 15 is a marker for the presence of metabolic syndrome. <i>Diabetes/Metabolism Research and Reviews</i> , 2019 , 35, e3085	7.5	14
71	Mitochondrial glycerol 3-phosphate dehydrogenase promotes skeletal muscle regeneration. <i>EMBO Molecular Medicine</i> , 2018 , 10,	12	14
70	JAZF1 Inhibits Adipose Tissue Macrophages and Adipose Tissue Inflammation in Diet-Induced Diabetic Mice. <i>BioMed Research International</i> , 2018 , 2018, 4507659	3	14
69	C1q/TNF-related protein-6 is associated with insulin resistance and the development of diabetes in Chinese population. <i>Acta Diabetologica</i> , 2018 , 55, 1221-1229	3.9	13
68	Effects of free fatty acids on plasma resistin and insulin resistance in awake rats. <i>Metabolism: Clinical and Experimental</i> , 2005 , 54, 1142-6	12.7	13

67	Circulating betatrophin is associated with insulin resistance in humans: cross-sectional and interventional studies and. <i>Oncotarget</i> , 2017 , 8, 96604-96614	3.3	13
66	NAMPT knockdown attenuates atherosclerosis and promotes reverse cholesterol transport in ApoE KO mice with high-fat-induced insulin resistance. <i>Scientific Reports</i> , 2016 , 6, 26746	4.9	13
65	Effects of glucagon-like peptide-1 agents on left ventricular function: systematic review and meta-analysis. <i>Annals of Medicine</i> , 2014 , 46, 664-71	1.5	11
64	Follistatin-like 1 as a Novel Adipomyokine Related to Insulin Resistance and Physical Activity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	11
63	Gut ghrelin regulates hepatic glucose production and insulin signaling via a gut-brain-liver pathway. <i>Cell Communication and Signaling</i> , 2019 , 17, 8	7.5	10
62	CILP-2 is a novel secreted protein and associated with insulin resistance. <i>Journal of Molecular Cell Biology</i> , 2019 , 11, 1083-1094	6.3	10
61	Impairment of Bitter Taste Sensor Transient Receptor Potential Channel M5-Mediated Aversion Aggravates High-Salt Intake and Hypertension. <i>Hypertension</i> , 2019 , 74, 1021-1032	8.5	10
60	Serum retinol-binding protein 4 levels are elevated but do not contribute to insulin resistance in newly diagnosed Chinese hypertensive patients. <i>Diabetology and Metabolic Syndrome</i> , 2014 , 6, 72	5.6	10
59	DOCK5 regulates energy balance and hepatic insulin sensitivity by targeting mTORC1 signaling. <i>EMBO Reports</i> , 2020 , 21, e49473	6.5	9
58	Serum Fetuin-A levels are increased and associated with insulin resistance in women with polycystic ovary syndrome. <i>BMC Endocrine Disorders</i> , 2020 , 20, 67	3.3	8
57	Comparison of Allogeneic Platelet-rich Plasma With Autologous Platelet-rich Plasma for the Treatment of Diabetic Lower Extremity Ulcers. <i>Cell Transplantation</i> , 2020 , 29, 963689720931428	4	8
56	Elevated circulating vaspin levels were decreased by rosiglitazone therapy in T2DM patients with poor glycemic control on metformin alone. <i>Cytokine</i> , 2011 , 56, 399-402	4	8
55	Central Sfrp5 regulates hepatic glucose flux and VLDL-triglyceride secretion. <i>Metabolism: Clinical and Experimental</i> , 2020 , 103, 154029	12.7	8
54	FGF21 facilitates autophagy in prostate cancer cells by inhibiting the PI3K-Akt-mTOR signaling pathway. <i>Cell Death and Disease</i> , 2021 , 12, 303	9.8	8
53	The potential effects of clinical antidiabetic agents on SARS-CoV-2. <i>Journal of Diabetes</i> , 2021 , 13, 243-253	3.8	8
52	Circulating bone morphogenetic protein-9 levels are associated with hypertension and insulin resistance in humans. <i>Journal of the American Society of Hypertension</i> , 2018 , 12, 372-380		7
51	Circulating ANGPTL8 Is Associated with the Presence of Metabolic Syndrome and Insulin Resistance in Polycystic Ovary Syndrome Young Women. <i>Mediators of Inflammation</i> , 2019 , 2019, 6321427	4.3	7
50	Global and Regional Effects of Bladder Cancer Risk Associated with Pioglitazone Therapy in Patients with Diabetes. <i>Scientific Reports</i> , 2017 , 7, 15804	4.9	7

49	Association between Serum Cystatin C and Diabetic Foot Ulceration in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Journal of Diabetes Research</i> , 2016 , 2016, 8029340	3.9	7
48	Increased plasma osteopontin levels are associated with nonalcoholic fatty liver disease in patients with type 2 diabetes mellitus. <i>Cytokine</i> , 2020 , 125, 154837	4	7
47	High Circulating Alarin Levels Are Associated with Presence of Metabolic Syndrome. <i>Cellular Physiology and Biochemistry</i> , 2018 , 51, 2041-2051	3.9	7
46	C1q/TNF-Related Protein5 (CTRP5) as a Biomarker to Predict Metabolic Syndrome and Each of Its Components. <i>International Journal of Endocrinology</i> , 2018 , 2018, 7201473	2.7	7
45	Effect of central JAZF1 on glucose production is regulated by the PI3K-Akt-AMPK pathway. <i>FASEB Journal</i> , 2020 , 34, 7058-7074	0.9	6
44	Adrenal artery ablation for primary aldosteronism without apparent aldosteronoma: An efficacy and safety, proof-of-principle trial. <i>Journal of Clinical Hypertension</i> , 2020 , 22, 1618-1626	2.3	6
43	Efficacy and Safety of Mulberry Twig Alkaloids Tablet for the Treatment of Type 2 Diabetes: A Multicenter, Randomized, Double-Blind, Double-Dummy, and Parallel Controlled Clinical Trial. <i>Diabetes Care</i> , 2021 , 44, 1324-1333	14.6	6
42	Circulating complement-1q tumor necrosis factor- β -related protein isoform β levels are low in type β diabetes patients and reduced by dapagliflozin. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 88-95	3.9	6
41	Non-insulin determinant pathways maintain glucose homeostasis upon metabolic surgery. <i>Cell Discovery</i> , 2018 , 4, 58	22.3	6
40	Circulating CTRP6 Levels are Increased in Overweight or Obese Chinese Individuals and Associated with Insulin Resistance Parameters: A Pilot Study. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021 , 129, 535-541	2.3	5
39	Pioglitazone Improved Insulin Sensitivity and First Phase Insulin Secretion Among Obese and Lean People with Diabetes: A Multicenter Clamp Study. <i>Diabetes Therapy</i> , 2018 , 9, 815-826	3.6	5
38	Reducing NADPH Synthesis Counteracts Diabetic Nephropathy through Restoration of AMPK Activity in Type 1 Diabetic Rats. <i>Cell Reports</i> , 2020 , 32, 108207	10.6	5
37	Elevated Circulating Fetuin-B Levels Are Associated with Insulin Resistance and Reduced by GLP-1RA in Newly Diagnosed PCOS Women. <i>Mediators of Inflammation</i> , 2020 , 2020, 2483435	4.3	5
36	Circulating alarin concentrations are high in patients with type 2 diabetes and increased by glucagon-like peptide-1 receptor agonist treatment: An Consort-compliant study. <i>Medicine (United States)</i> , 2019 , 98, e16428	1.8	5
35	Efficacy and safety of polyethylene glycol loxenatide monotherapy in type 2 diabetes patients: A multicentre, randomized, double-blind, placebo-controlled phase 3a clinical trial. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 116-124	6.7	5
34	TERT and Akt Are Involved in the Par-4-Dependent Apoptosis of Islet Cells in Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2018 , 2018, 7653904	3.9	5
33	Association of circulating BMP9 with coronary heart disease and hypertension in Chinese populations. <i>BMC Cardiovascular Disorders</i> , 2019 , 19, 131	2.3	4
32	Sfrp5 interacts with Slurp1 to regulate the accumulation of triglycerides in hepatocyte steatosis model. <i>Biochemical and Biophysical Research Communications</i> , 2019 , 512, 256-262	3.4	4

31	Adipose Insulin Resistance and Circulating Betatrophin Levels in Women with PCOS. <i>BioMed Research International</i> , 2020 , 2020, 1253164	3	4
30	Adiponectin/(FBG [FIns) as a predictor of insulin sensitivity and metabolic syndrome in patients with polycystic ovary syndrome. <i>Medicine (United States)</i> , 2016 , 95, e5524	1.8	4
29	Human C-reactive protein impedes entry of leptin into the CNS and attenuates its physiological actions in the CNS. <i>Biochemical Journal</i> , 2016 , 473, 1215-24	3.8	4
28	Exenatide once-weekly injection for the treatment of type 2 diabetes in Chinese patients: current perspectives. <i>Therapeutics and Clinical Risk Management</i> , 2015 , 11, 1153-62	2.9	4
27	LASS2 inhibits proliferation and induces apoptosis in HepG2 cells by affecting mitochondrial dynamics, the cell cycle and the nuclear factor- κ B pathways. <i>Oncology Reports</i> , 2019 , 41, 3005-3014	3.5	4
26	Association between circulating follistatin-like-1 and metabolic syndrome in middle-aged and old population: A cross-sectional study. <i>Diabetes/Metabolism Research and Reviews</i> , 2021 , 37, e3373	7.5	4
25	Genetic ablation of C-reactive protein gene confers resistance to obesity and insulin resistance in rats. <i>Diabetologia</i> , 2021 , 64, 1169-1183	10.3	4
24	In vivo effects of phosphodiesterase III inhibitors on glucose metabolism and insulin sensitivity. <i>Journal of the Chinese Medical Association</i> , 2003 , 66, 210-6	2.8	4
23	Several Circulating Biomarkers for PCOS Diagnosis. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021 , 129, 705-712	2.3	3
22	Salt-Induced Hepatic Inflammatory Memory Contributes to Cardiovascular Damage Through Epigenetic Modulation of SIRT3.. <i>Circulation</i> , 2022 , 145, 375-391	16.7	3
21	Efficacy and safety of polyethylene glycol loxenatide as add-on to metformin in patients with type 2 diabetes: A multicentre, randomized, double-blind, placebo-controlled, phase 3b trial. <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 2375-2383	6.7	3
20	Dock5 controls the peripheral B cell differentiation via regulating BCR signaling and actin reorganization. <i>Cellular Immunology</i> , 2019 , 337, 15-21	4.4	2
19	Effective and safe delivery of GLP-1AR and FGF-21 plasmids using amino-functionalized dual-mesoporous silica nanoparticles in vitro and in vivo. <i>Biomaterials</i> , 2021 , 271, 120763	15.6	2
18	Dedicator of Cytokinesis 5 Regulates Keratinocyte Function and Promotes Diabetic Wound Healing. <i>Diabetes</i> , 2021 , 70, 1170-1184	0.9	2
17	Hepatic lipid accumulation induced by a high-fat diet is regulated by Nrf2 through multiple pathways.. <i>FASEB Journal</i> , 2022 , 36, e22280	0.9	2
16	Circulating CTRP7 Is a Potential Predictor for Metabolic Syndrome. <i>Frontiers in Endocrinology</i> , 2021 , 12, 774309	5.7	1
15	Serum Fetuin-B Levels Are Elevated in Women with Metabolic Syndrome and Associated with Increased Oxidative Stress. <i>Oxidative Medicine and Cellular Longevity</i> , 2021 , 2021, 6657658	6.7	1
14	Circulating Levels of CILP2 Are Elevated in Coronary Heart Disease and Associated with Atherosclerosis. <i>Oxidative Medicine and Cellular Longevity</i> , 2020 , 2020, 1871984	6.7	1

13	Association between Visceral Fat and Bone Mineral Density in Both Male and Female Patients with Adult Growth Hormone Deficiency. <i>Biochemistry Research International</i> , 2020 , 2020, 5079625	2.4	1
12	LASS2 regulates hepatocyte steatosis by interacting with NDUFS2/OXPHOS related proteins. <i>Biochemical and Biophysical Research Communications</i> , 2020 , 526, 871-879	3.4	1
11	A reusable colorimetric assay based on mixed valence state Ce-MOF@Pt nanoparticles for highly sensitive detection of visfatin. <i>Analytica Chimica Acta</i> , 2021 , 1146, 24-32	6.6	1
10	Response to Letter to the Editor: "Myonectin Predicts the Development of Type 2 Diabetes". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 4040-4041	5.6	1
9	Catheter-Based Adrenal Ablation Remits Primary Aldosteronism: A Randomized Medication-Controlled Trial. <i>Circulation</i> , 2021 , 144, 580-582	16.7	1
8	Association of serum fetuin-B with insulin resistance and pre-diabetes in young Chinese women: evidence from a cross-sectional study and effect of liraglutide. <i>PeerJ</i> , 2021 , 9, e11869	3.1	1
7	Association of metabolic syndrome components with circulating levels of cytokine clusters in young women. <i>Endocrine Connections</i> , 2021 , 10, 66-75	3.5	1
6	A Novel Immune and Stroma Related Prognostic Marker for Invasive Breast Cancer in Tumor Microenvironment: A TCGA Based Study. <i>Frontiers in Endocrinology</i> , 2021 , 12, 774244	5.7	0
5	Component of oligomeric Golgi complex 1 deficiency leads to hypoglycemia: a case report and literature review. <i>BMC Pediatrics</i> , 2021 , 21, 442	2.6	0
4	Hypothalamic BMP9 suppresses glucose production by central PI3K/Akt/mTOR pathway. <i>Journal of Endocrinology</i> , 2021 , 248, 221-235	4.7	0
3	Construction and Validation of a Prediction Model for Identifying Clinical Risk Factors of Lateral Lymph Node Metastasis in Medullary Thyroid Carcinoma.. <i>International Journal of General Medicine</i> , 2022 , 15, 2301-2309	2.3	0
2	GNG2 acts as a tumor suppressor in breast cancer through stimulating MRAS signaling.. <i>Cell Death and Disease</i> , 2022 , 13, 260	9.8	0
1	CTRP7 Is a Biomarker Related to Insulin Resistance and Oxidative Stress: Cross-Sectional and Intervention Studies In and In .. <i>Oxidative Medicine and Cellular Longevity</i> , 2022 , 2022, 6877609	6.7	0