

Gangyi Yang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5486658/publications.pdf>

Version: 2024-02-01

123
papers

2,920
citations

185998

28
h-index

253896

43
g-index

128
all docs

128
docs citations

128
times ranked

3925
citing authors

#	ARTICLE	IF	CITATIONS
1	Nesfatin-1 Action in the Brain Increases Insulin Sensitivity Through Akt/AMPK/TORC2 Pathway in Diet-Induced Insulin Resistance. <i>Diabetes</i> , 2012, 61, 1959-1968.	0.3	112
2	Circulating Sfrp5 Is a Signature of Obesity-Related Metabolic Disorders and Is Regulated by Glucose and Liraglutide in Humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 290-298.	1.8	103
3	Zinc- α 2-Glycoprotein Is Associated With Insulin Resistance in Humans and Is Regulated by Hyperglycemia, Hyperinsulinemia, or Liraglutide Administration. <i>Diabetes Care</i> , 2013, 36, 1074-1082.	4.3	83
4	Elevated Circulating Levels of Irisin and the Effect of Metformin Treatment in Women With Polycystic Ovary Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 1485-1493.	1.8	78
5	Inhibition of Mitochondrial Calcium Overload by SIRT3 Prevents Obesity- or Age-Related Whitening of Brown Adipose Tissue. <i>Diabetes</i> , 2020, 69, 165-180.	0.3	77
6	Alteration of gut microbiota induced by DPP-4i treatment improves glucose homeostasis. <i>EBioMedicine</i> , 2019, 44, 665-674.	2.7	66
7	Serum C1q/TNF-related protein-3 (CTRP3) levels are decreased in obesity and hypertension and are negatively correlated with parameters of insulin resistance. <i>Diabetology and Metabolic Syndrome</i> , 2015, 7, 33.	1.2	60
8	The adipose triglyceride lipase, adiponectin and visfatin are downregulated by tumor necrosis factor- α (TNF- α) in vivo. <i>Cytokine</i> , 2009, 45, 12-19.	1.4	58
9	Overexpression of visfatin/PBEF/Nampt alters whole-body insulin sensitivity and lipid profile in rats. <i>Annals of Medicine</i> , 2009, 41, 311-320.	1.5	56
10	DPP-4 Inhibitors Improve Diabetic Wound Healing via Direct and Indirect Promotion of Epithelial-Mesenchymal Transition and Reduction of Scarring. <i>Diabetes</i> , 2018, 67, 518-531.	0.3	56
11	Osteoprotegerin Promotes Liver Steatosis by Targeting the ERK-PPAR- α -CD36 Pathway. <i>Diabetes</i> , 2019, 68, 1902-1914.	0.3	56
12	Circulating preptin levels in normal, impaired glucose tolerance, and type 2 diabetic subjects. <i>Annals of Medicine</i> , 2009, 41, 52-56.	1.5	55
13	Activation of TRPV1 channel antagonizes diabetic nephropathy through inhibiting endoplasmic reticulum-mitochondria contact in podocytes. <i>Metabolism: Clinical and Experimental</i> , 2020, 105, 154182.	1.5	53
14	Myonectin Predicts the Development of Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 139-147.	1.8	52
15	Hypothalamic Nesfatin-1/NUCB2 Knockdown Augments Hepatic Gluconeogenesis That Is Correlated With Inhibition of mTOR-STAT3 Signaling Pathway in Rats. <i>Diabetes</i> , 2014, 63, 1234-1247.	0.3	50
16	Liraglutide Prevents Hypoadiponectinemia-Induced Insulin Resistance and Alterations of Gene Expression Involved in Glucose and Lipid Metabolism. <i>Molecular Medicine</i> , 2011, 17, 1168-1178.	1.9	49
17	The role of JAZF1 on lipid metabolism and related genes in vitro. <i>Metabolism: Clinical and Experimental</i> , 2011, 60, 523-530.	1.5	48
18	Liraglutide Increases FGF-21 Activity and Insulin Sensitivity in High Fat Diet and Adiponectin Knockdown Induced Insulin Resistance. <i>PLoS ONE</i> , 2012, 7, e48392.	1.1	48

#	ARTICLE	IF	CITATIONS
19	Kräppl-like factor 14 increases insulin sensitivity through activation of PI3K/Akt signal pathway. <i>Cellular Signalling</i> , 2015, 27, 2201-2208.	1.7	48
20	Sodium-Glucose Cotransporter 2 (SGLT2) Inhibitor Increases Circulating Zinc-Î2-Glycoprotein Levels in Patients with Type 2 Diabetes. <i>Scientific Reports</i> , 2016, 6, 32887.	1.6	47
21	The effects of fibroblast growth factor-21 knockdown and over-expression on its signaling pathway and glucoseâ€“lipid metabolism in vitro. <i>Molecular and Cellular Endocrinology</i> , 2012, 348, 21-26.	1.6	45
22	LAMP3 regulates hepatic lipid metabolism through activating PI3K/Akt pathway. <i>Molecular and Cellular Endocrinology</i> , 2018, 470, 160-167.	1.6	44
23	Salt-Induced Hepatic Inflammatory Memory Contributes to Cardiovascular Damage Through Epigenetic Modulation of SIRT3. <i>Circulation</i> , 2022, 145, 375-391.	1.6	38
24	Decreased circulating BMP-9 levels in patients with Type 2 diabetes is a signature of insulin resistance. <i>Clinical Science</i> , 2017, 131, 239-246.	1.8	37
25	JAZF1 ameliorates age and diet-associated hepatic steatosis through SREBP-1c -dependent mechanism. <i>Cell Death and Disease</i> , 2018, 9, 859.	2.7	36
26	Role of bone morphogenetic proteinâ€“9 in the regulation of glucose and lipid metabolism. <i>FASEB Journal</i> , 2019, 33, 10077-10088.	0.2	35
27	Efficacy and Safety of Aldose Reductase Inhibitor for the Treatment of Diabetic Cardiovascular Autonomic Neuropathy: Systematic Review and Meta-Analysis. <i>PLoS ONE</i> , 2014, 9, e87096.	1.1	35
28	The natural logarithm of zinc-Î2-glycoprotein/HOMA-IR is a better predictor of insulin sensitivity than the product of triglycerides and glucose and the other lipid ratios. <i>Cytokine</i> , 2016, 79, 96-102.	1.4	32
29	FGF21 facilitates autophagy in prostate cancer cells by inhibiting the PI3Kâ€“Aktâ€“mTOR signaling pathway. <i>Cell Death and Disease</i> , 2021, 12, 303.	2.7	31
30	Efficacy of low-level light therapy for treatment of diabetic foot ulcer: A systematic review and meta-analysis of randomized controlled trials. <i>Diabetes Research and Clinical Practice</i> , 2018, 143, 215-224.	1.1	30
31	Deficiency of Mitochondrial Glycerol 3â€“Phosphate Dehydrogenase Contributes to Hepatic Steatosis. <i>Hepatology</i> , 2019, 70, 84-97.	3.6	30
32	Glypican-4 is increased in human subjects with impaired glucose tolerance and decreased in patients with newly diagnosed type 2 diabetes. <i>Acta Diabetologica</i> , 2014, 51, 981-990.	1.2	29
33	Duodenal GLP-1 signaling regulates hepatic glucose production through a PKC-Î-dependent neurocircuitry. <i>Cell Death and Disease</i> , 2017, 8, e2609-e2609.	2.7	29
34	Plasma Sfrp5 levels correlate with determinants of the metabolic syndrome in Chinese adults. <i>Diabetes/Metabolism Research and Reviews</i> , 2017, 33, e2896.	1.7	29
35	Silencing of <sc>FGF</sc>â€“21 expression promotes hepatic gluconeogenesis and glycogenolysis by regulation of the <sc>STAT</sc>â€“3â€“<sc>SOCS</sc>3 signal. <i>FEBS Journal</i> , 2014, 281, 2136-2147.	2.2	28
36	Circulating Zinc-Î2-glycoprotein levels and Insulin Resistance in Polycystic Ovary Syndrome. <i>Scientific Reports</i> , 2016, 6, 25934.	1.6	28

#	ARTICLE	IF	CITATIONS
37	Circulating bone morphogenetic protein-9 in relation to metabolic syndrome and insulin resistance. <i>Scientific Reports</i> , 2017, 7, 17529.	1.6	28
38	Circulating zinc- α 2-glycoprotein levels are low in newly diagnosed patients with metabolic syndrome and correlate with adiponectin. <i>Nutrition and Metabolism</i> , 2017, 14, 53.	1.3	27
39	Deficiency of PKD2L1 (TRPP3) Exacerbates Pathological Cardiac Hypertrophy by Augmenting NCX1-Mediated Mitochondrial Calcium Overload. <i>Cell Reports</i> , 2018, 24, 1639-1652.	2.9	27
40	Overexpression of JAZF1 protected ApoE-deficient mice from atherosclerosis by inhibiting hepatic cholesterol synthesis via CREB-dependent mechanisms. <i>International Journal of Cardiology</i> , 2014, 177, 100-110.	0.8	26
41	Effects of sitagliptin on circulating zinc- α 2-glycoprotein levels in newly diagnosed type 2 diabetes patients: a randomized trial. <i>European Journal of Endocrinology</i> , 2016, 174, 147-155.	1.9	25
42	Changes in whole metabolites after exenatide treatment in overweight/obese polycystic ovary syndrome patients. <i>Clinical Endocrinology</i> , 2019, 91, 508-516.	1.2	25
43	Circulating C1q/TNF-related protein isoform 15 is a marker for the presence of metabolic syndrome. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3085.	1.7	25
44	Follistatin-like 1 as a Novel Adipomyokine Related to Insulin Resistance and Physical Activity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e4499-e4509.	1.8	25
45	Transcutaneous oxygen pressure (TcPO ₂): A novel diagnostic tool for peripheral neuropathy in type 2 diabetes patients. <i>Diabetes Research and Clinical Practice</i> , 2014, 105, 336-343.	1.1	24
46	Polyethylene glycol loxenatide injections added to metformin effectively improve glycemic control and exhibit favorable safety in type 2 diabetic patients. <i>Journal of Diabetes</i> , 2017, 9, 158-167.	0.8	24
47	Rictor positively regulates B cell receptor signaling by modulating actin reorganization via ezrin. <i>PLoS Biology</i> , 2017, 15, e2001750.	2.6	24
48	Mitochondrial glycerol 3-phosphate dehydrogenase promotes skeletal muscle regeneration. <i>EMBO Molecular Medicine</i> , 2018, 10, .	3.3	24
49	Efficacy and Safety of Mulberry Twig Alkaloids Tablet for the Treatment of Type 2 Diabetes: A Multicenter, Randomized, Double-Blind, Double-Dummy, and Parallel Controlled Clinical Trial. <i>Diabetes Care</i> , 2021, 44, 1324-1333.	4.3	24
50	Overexpression of juxtaposed with another zinc finger gene Δ 1 reduces proinflammatory cytokine release via inhibition of stress-activated protein kinases and nuclear factor- κ B. <i>FEBS Journal</i> , 2014, 281, 3193-3205.	2.2	23
51	A novel role for the Kr α ppel-like factor 14 on macrophage inflammatory response and atherosclerosis development. <i>Cardiovascular Pathology</i> , 2017, 27, 1-8.	0.7	23
52	Comparison of Allogeneic Platelet-rich Plasma With Autologous Platelet-rich Plasma for the Treatment of Diabetic Lower Extremity Ulcers. <i>Cell Transplantation</i> , 2020, 29, 096368972093142.	1.2	22
53	Efficacy and safety of linagliptin in Asian patients with type 2 diabetes mellitus inadequately controlled by metformin: A multinational 24-week, randomized clinical trial. <i>Journal of Diabetes</i> , 2016, 8, 229-237.	0.8	20
54	JAZF1 Inhibits Adipose Tissue Macrophages and Adipose Tissue Inflammation in Diet-Induced Diabetic Mice. <i>BioMed Research International</i> , 2018, 2018, 1-10.	0.9	20

#	ARTICLE	IF	CITATIONS
55	CILP-2 is a novel secreted protein and associated with insulin resistance. <i>Journal of Molecular Cell Biology</i> , 2019, 11, 1083-1094.	1.5	19
56	Serum Fetuin-A levels are increased and associated with insulin resistance in women with polycystic ovary syndrome. <i>BMC Endocrine Disorders</i> , 2020, 20, 67.	0.9	19
57	Circulating betatrophin is associated with insulin resistance in humans: cross-sectional and interventional studies <i>in vivo</i> and <i>in vitro</i> . <i>Oncotarget</i> , 2017, 8, 96604-96614.	0.8	19
58	C1q/TNF-related protein-6 is associated with insulin resistance and the development of diabetes in Chinese population. <i>Acta Diabetologica</i> , 2018, 55, 1221-1229.	1.2	18
59	Effective gene delivery of shBMP-9 using polyethyleneimine-based core-shell nanoparticles in an animal model of insulin resistance. <i>Nanoscale</i> , 2019, 11, 2008-2016.	2.8	18
60	Adrenal artery ablation for primary aldosteronism without apparent aldosteronoma: An efficacy and safety, proof-of-principle trial. <i>Journal of Clinical Hypertension</i> , 2020, 22, 1618-1626.	1.0	18
61	Short-term pioglitazone treatment prevents free fatty acid-induced hepatic insulin resistance in normal rats: Possible role of the resistin and adiponectin. <i>Biochemical and Biophysical Research Communications</i> , 2006, 339, 1190-1196.	1.0	17
62	Central Sfrp5 regulates hepatic glucose flux and VLDL-triglyceride secretion. <i>Metabolism: Clinical and Experimental</i> , 2020, 103, 154029.	1.5	17
63	Efficacy and safety of polyethylene glycol loxenatide monotherapy in type 2 diabetes patients: A multicentre, randomized, double-blind, placebo-controlled phase 3 clinical trial. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 116-124.	2.2	17
64	Genetic ablation of C-reactive protein gene confers resistance to obesity and insulin resistance in rats. <i>Diabetologia</i> , 2021, 64, 1169-1183.	2.9	17
65	Effects of free fatty acids on plasma resistin and insulin resistance in awake rats. <i>Metabolism: Clinical and Experimental</i> , 2005, 54, 1142-1146.	1.5	16
66	NAMPT knockdown attenuates atherosclerosis and promotes reverse cholesterol transport in ApoE KO mice with high-fat-induced insulin resistance. <i>Scientific Reports</i> , 2016, 6, 26746.	1.6	16
67	Gut ghrelin regulates hepatic glucose production and insulin signaling via a gut-brain-liver pathway. <i>Cell Communication and Signaling</i> , 2019, 17, 8.	2.7	16
68	DOCK5 regulates energy balance and hepatic insulin sensitivity by targeting mTORC1 signaling. <i>EMBO Reports</i> , 2020, 21, e49473.	2.0	16
69	Effect of central JAZF1 on glucose production is regulated by the PI3K-Akt-AMPK pathway. <i>FASEB Journal</i> , 2020, 34, 7058-7074.	0.2	16
70	Catheter-Based Adrenal Ablation Remits Primary Aldosteronism: A Randomized Medication-Controlled Trial. <i>Circulation</i> , 2021, 144, 580-582.	1.6	16
71	Serum retinol-binding protein 4 levels are elevated but do not contribute to insulin resistance in newly diagnosed Chinese hypertensive patients. <i>Diabetology and Metabolic Syndrome</i> , 2014, 6, 72.	1.2	15
72	Elevated Circulating Fetuin-B Levels Are Associated with Insulin Resistance and Reduced by GLP-1RA in Newly Diagnosed PCOS Women. <i>Mediators of Inflammation</i> , 2020, 2020, 1-12.	1.4	15

#	ARTICLE	IF	CITATIONS
73	The potential effects of clinical antidiabetic agents on <sc>SARSâ€CoV</sc> â€2. Journal of Diabetes, 2021, 13, 243-252.	0.8	15
74	Effective and safe delivery of GLP-1AR and FGF-21 plasmids using amino-functionalized dual-mesoporous silica nanoparticles in vitro and in vivo. Biomaterials, 2021, 271, 120763.	5.7	15
75	Impairment of Bitter Taste Sensor Transient Receptor Potential Channel M5-Mediated Aversion Aggravates High-Salt Intake and Hypertension. Hypertension, 2019, 74, 1021-1032.	1.3	14
76	Efficacy and safety of polyethylene glycol loxenatide as addâ€on to metformin in patients with type 2 diabetes: A multicentre, randomized, doubleâ€blind, placeboâ€controlled, phase 3b trial. Diabetes, Obesity and Metabolism, 2020, 22, 2375-2383.	2.2	14
77	Hepatic lipid accumulation induced by a highâ€fat diet is regulated by Nrf2 through multiple pathways. FASEB Journal, 2022, 36, e22280.	0.2	14
78	Global and Regional Effects of Bladder Cancer Risk Associated with Pioglitazone Therapy in Patients with Diabetes. Scientific Reports, 2017, 7, 15804.	1.6	13
79	Effects of glucagon-like peptide-1 agents on left ventricular function: Systematic review and meta-analysis. Annals of Medicine, 2014, 46, 664-671.	1.5	12
80	Circulating bone morphogenetic protein-9 levels are associated with hypertension and insulin resistance in humans. Journal of the American Society of Hypertension, 2018, 12, 372-380.	2.3	12
81	Reducing NADPH Synthesis Counteracts Diabetic Nephropathy through Restoration of AMPK Activity in Type 1 Diabetic Rats. Cell Reports, 2020, 32, 108207.	2.9	12
82	Association between Serum Cystatin C and Diabetic Foot Ulceration in Patients with Type 2 Diabetes: A Cross-Sectional Study. Journal of Diabetes Research, 2016, 2016, 1-7.	1.0	11
83	Pioglitazone Improved Insulin Sensitivity and First Phase Insulin Secretion Among Obese and Lean People with Diabetes: A Multicenter Clamp Study. Diabetes Therapy, 2018, 9, 815-826.	1.2	11
84	Circulating CTRP6 Levels are Increased in Overweight or Obese Chinese Individuals and Associated with Insulin Resistance Parameters: A Pilot Study. Experimental and Clinical Endocrinology and Diabetes, 2021, 129, 535-541.	0.6	11
85	Increased plasma osteopontin levels are associated with nonalcoholic fatty liver disease in patients with type 2 diabetes mellitus. Cytokine, 2020, 125, 154837.	1.4	11
86	Adipose Insulin Resistance and Circulating Betatrophin Levels in Women with PCOS. BioMed Research International, 2020, 2020, 1-9.	0.9	11
87	Dedicator of Cytokinesis 5 Regulates Keratinocyte Function and Promotes Diabetic Wound Healing. Diabetes, 2021, 70, 1170-1184.	0.3	11
88	Circulating ANGPTL8 Is Associated with the Presence of Metabolic Syndrome and Insulin Resistance in Polycystic Ovary Syndrome Young Women. Mediators of Inflammation, 2019, 2019, 1-10.	1.4	10
89	Elevated circulating vaspin levels were decreased by rosiglitazone therapy in T2DM patients with poor glycemic control on metformin alone. Cytokine, 2011, 56, 399-402.	1.4	9
90	High Circulating Alarin Levels Are Associated with Presence of Metabolic Syndrome. Cellular Physiology and Biochemistry, 2018, 51, 2041-2051.	1.1	9

#	ARTICLE	IF	CITATIONS
91	C1q/TNF-Related Protein5 (CTRP5) as a Biomarker to Predict Metabolic Syndrome and Each of Its Components. <i>International Journal of Endocrinology</i> , 2018, 2018, 1-8.	0.6	9
92	Association between circulating follistatin-like-1 and metabolic syndrome in middle-aged and old population: A cross-sectional study. <i>Diabetes/Metabolism Research and Reviews</i> , 2021, 37, e3373.	1.7	9
93	C-reactive protein perturbs alveolar bone homeostasis: An experimental study of periodontitis and diabetes in the rat. <i>Journal of Clinical Periodontology</i> , 2022, 49, 1052-1066.	2.3	9
94	Non-insulin determinant pathways maintain glucose homeostasis upon metabolic surgery. <i>Cell Discovery</i> , 2018, 4, 58.	3.1	8
95	Several Circulating Biomarkers for PCOS Diagnosis. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021, 129, 705-712.	0.6	8
96	Sfrp5 interacts with Slurp1 to regulate the accumulation of triglycerides in hepatocyte steatosis model. <i>Biochemical and Biophysical Research Communications</i> , 2019, 512, 256-262.	1.0	8
97	Circulating alarin concentrations are high in patients with type 2 diabetes and increased by glucagon-like peptide-1 receptor agonist treatment. <i>Medicine (United States)</i> , 2019, 98, e16428.	0.4	8
98	TERT and Akt Are Involved in the Par-4-Dependent Apoptosis of Islet β Cells in Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2018, 2018, 1-13.	1.0	7
99	Association of circulating BMP9 with coronary heart disease and hypertension in Chinese populations. <i>BMC Cardiovascular Disorders</i> , 2019, 19, 131.	0.7	7
100	Circulating Levels of CILP2 Are Elevated in Coronary Heart Disease and Associated with Atherosclerosis. <i>Oxidative Medicine and Cellular Longevity</i> , 2020, 2020, 1-11.	1.9	7
101	A reusable colorimetric assay based on mixed valence state Ce-MOF@Pt nanoparticles for highly sensitive detection of visfatin. <i>Analytica Chimica Acta</i> , 2021, 1146, 24-32.	2.6	7
102	Circulating CTRP7 Is a Potential Predictor for Metabolic Syndrome. <i>Frontiers in Endocrinology</i> , 2021, 12, 774309.	1.5	7
103	Adiponectin/(FBG—HbA1c) as a predictor of insulin sensitivity and metabolic syndrome in patients with polycystic ovary syndrome. <i>Medicine (United States)</i> , 2016, 95, e5524.	0.4	6
104	LASS2 inhibits proliferation and induces apoptosis in HepG2 cells by affecting mitochondrial dynamics, the cell cycle and the nuclear factor- κ B pathways. <i>Oncology Reports</i> , 2019, 41, 3005-3014.	1.2	6
105	Circulating complement C1q tumor necrosis factor-related protein isoform 5 levels are low in type 2 diabetes patients and reduced by dapagliflozin. <i>Journal of Diabetes Investigation</i> , 2020, 11, 88-95.	1.1	6
106	Serum Fetuin-B Levels Are Elevated in Women with Metabolic Syndrome and Associated with Increased Oxidative Stress. <i>Oxidative Medicine and Cellular Longevity</i> , 2021, 2021, 1-17.	1.9	6
107	Human C-reactive protein impedes entry of leptin into the CNS and attenuates its physiological actions in the CNS. <i>Biochemical Journal</i> , 2016, 473, 1215-1224.	1.7	5
108	LASS2 regulates hepatocyte steatosis by interacting with NDUFS2/OXPHOS related proteins. <i>Biochemical and Biophysical Research Communications</i> , 2020, 526, 871-879.	1.0	5

#	ARTICLE	IF	CITATIONS
109	A Novel Immune and Stroma Related Prognostic Marker for Invasive Breast Cancer in Tumor Microenvironment: A TCGA Based Study. <i>Frontiers in Endocrinology</i> , 2021, 12, 774244.	1.5	5
110	CTRP7 Is a Biomarker Related to Insulin Resistance and Oxidative Stress: Cross-Sectional and Intervention Studies In Vivo and In Vitro. <i>Oxidative Medicine and Cellular Longevity</i> , 2022, 2022, 1-17.	1.9	5
111	Exenatide once-weekly injection for the treatment of type 2 diabetes in Chinese patients: current perspectives. <i>Therapeutics and Clinical Risk Management</i> , 2015, 11, 1153.	0.9	4
112	Hypothalamic BMP9 suppresses glucose production by central PI3K/Akt/mTOR pathway. <i>Journal of Endocrinology</i> , 2021, 248, 221-235.	1.2	4
113	Association of metabolic syndrome components with circulating levels of cytokine clusters in young women. <i>Endocrine Connections</i> , 2021, 10, 66-75.	0.8	4
114	In vivo effects of phosphodiesterase III inhibitors on glucose metabolism and insulin sensitivity. <i>Journal of the Chinese Medical Association</i> , 2003, 66, 210-6.	0.6	4
115	GNG2 acts as a tumor suppressor in breast cancer through stimulating MRAS signaling. <i>Cell Death and Disease</i> , 2022, 13, 260.	2.7	4
116	Dock5 controls the peripheral B cell differentiation via regulating BCR signaling and actin reorganization. <i>Cellular Immunology</i> , 2019, 337, 15-21.	1.4	3
117	Association of serum fetuin-B with insulin resistance and pre-diabetes in young Chinese women: evidence from a cross-sectional study and effect of liraglutide. <i>PeerJ</i> , 2021, 9, e11869.	0.9	3
118	Response to Letter to the Editor: "Myonectin Predicts the Development of Type 2 Diabetes". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 4040-4041.	1.8	2
119	GPHB5 Is a Biomarker in Women With Metabolic Syndrome: Results From Cross-Sectional and Intervention Studies. <i>Frontiers in Endocrinology</i> , 0, 13, .	1.5	2
120	Association between Visceral Fat and Bone Mineral Density in Both Male and Female Patients with Adult Growth Hormone Deficiency. <i>Biochemistry Research International</i> , 2020, 2020, 1-6.	1.5	1
121	Component of oligomeric Golgi complex 1 deficiency leads to hypoglycemia: a case report and literature review. <i>BMC Pediatrics</i> , 2021, 21, 442.	0.7	1
122	Construction and Validation of a Prediction Model for Identifying Clinical Risk Factors of Lateral Lymph Node Metastasis in Medullary Thyroid Carcinoma. <i>International Journal of General Medicine</i> , 2022, Volume 15, 2301-2309.	0.8	1
123	TRPC5 deletion in the central amygdala antagonizes high-fat diet-induced obesity by increasing sympathetic innervation. <i>International Journal of Obesity</i> , 2022, 46, 1544-1555.	1.6	1