

Gangyi Yang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5486658/publications.pdf>

Version: 2024-02-01

123
papers

2,920
citations

185998

28
h-index

253896

43
g-index

128
all docs

128
docs citations

128
times ranked

3925
citing authors

#	ARTICLE	IF	CITATIONS
1	Salt-Induced Hepatic Inflammatory Memory Contributes to Cardiovascular Damage Through Epigenetic Modulation of SIRT3. <i>Circulation</i> , 2022, 145, 375-391.	1.6	38
2	Construction and Validation of a Prediction Model for Identifying Clinical Risk Factors of Lateral Lymph Node Metastasis in Medullary Thyroid Carcinoma. <i>International Journal of General Medicine</i> , 2022, Volume 15, 2301-2309.	0.8	1
3	GNG2 acts as a tumor suppressor in breast cancer through stimulating MRAS signaling. <i>Cell Death and Disease</i> , 2022, 13, 260.	2.7	4
4	CTRP7 Is a Biomarker Related to Insulin Resistance and Oxidative Stress: Cross-Sectional and Intervention Studies In Vivo and In Vitro. <i>Oxidative Medicine and Cellular Longevity</i> , 2022, 2022, 1-17.	1.9	5
5	Hepatic lipid accumulation induced by a high-fat diet is regulated by Nrf2 through multiple pathways. <i>FASEB Journal</i> , 2022, 36, e22280.	0.2	14
6	TRPC5 deletion in the central amygdala antagonizes high-fat diet-induced obesity by increasing sympathetic innervation. <i>International Journal of Obesity</i> , 2022, 46, 1544-1555.	1.6	1
7	C-reactive protein perturbs alveolar bone homeostasis: An experimental study of periodontitis and diabetes in the rat. <i>Journal of Clinical Periodontology</i> , 2022, 49, 1052-1066.	2.3	9
8	Several Circulating Biomarkers for PCOS Diagnosis. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021, 129, 705-712.	0.6	8
9	Circulating CTRP6 Levels are Increased in Overweight or Obese Chinese Individuals and Associated with Insulin Resistance Parameters: A Pilot Study. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021, 129, 535-541.	0.6	11
10	Association between circulating follistatin-like-1 and metabolic syndrome in middle-aged and old population: A cross-sectional study. <i>Diabetes/Metabolism Research and Reviews</i> , 2021, 37, e3373.	1.7	9
11	The potential effects of clinical antidiabetic agents on SARS-CoV-2. <i>Journal of Diabetes</i> , 2021, 13, 243-252.	0.8	15
12	Efficacy and safety of polyethylene glycol loxenatide monotherapy in type 2 diabetes patients: A multicentre, randomized, double-blind, placebo-controlled phase 3a clinical trial. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 116-124.	2.2	17
13	Genetic ablation of C-reactive protein gene confers resistance to obesity and insulin resistance in rats. <i>Diabetologia</i> , 2021, 64, 1169-1183.	2.9	17
14	Hypothalamic BMP9 suppresses glucose production by central PI3K/Akt/mTOR pathway. <i>Journal of Endocrinology</i> , 2021, 248, 221-235.	1.2	4
15	Dedicator of Cytokinesis 5 Regulates Keratinocyte Function and Promotes Diabetic Wound Healing. <i>Diabetes</i> , 2021, 70, 1170-1184.	0.3	11
16	A reusable colorimetric assay based on mixed valence state Ce-MOF@Pt nanoparticles for highly sensitive detection of visfatin. <i>Analytica Chimica Acta</i> , 2021, 1146, 24-32.	2.6	7
17	FGF21 facilitates autophagy in prostate cancer cells by inhibiting the PI3K-Akt-mTOR signaling pathway. <i>Cell Death and Disease</i> , 2021, 12, 303.	2.7	31
18	Effective and safe delivery of GLP-1AR and FGF-21 plasmids using amino-functionalized dual-mesoporous silica nanoparticles in vitro and in vivo. <i>Biomaterials</i> , 2021, 271, 120763.	5.7	15

#	ARTICLE	IF	CITATIONS
19	Efficacy and Safety of Mulberry Twig Alkaloids Tablet for the Treatment of Type 2 Diabetes: A Multicenter, Randomized, Double-Blind, Double-Dummy, and Parallel Controlled Clinical Trial. <i>Diabetes Care</i> , 2021, 44, 1324-1333.	4.3	24
20	Catheter-Based Adrenal Ablation Remits Primary Aldosteronism: A Randomized Medication-Controlled Trial. <i>Circulation</i> , 2021, 144, 580-582.	1.6	16
21	Association of serum fetuin-B with insulin resistance and pre-diabetes in young Chinese women: evidence from a cross-sectional study and effect of liraglutide. <i>PeerJ</i> , 2021, 9, e11869.	0.9	3
22	Association of metabolic syndrome components with circulating levels of cytokine clusters in young women. <i>Endocrine Connections</i> , 2021, 10, 66-75.	0.8	4
23	Component of oligomeric Golgi complex 1 deficiency leads to hypoglycemia: a case report and literature review. <i>BMC Pediatrics</i> , 2021, 21, 442.	0.7	1
24	Serum Fetuin-B Levels Are Elevated in Women with Metabolic Syndrome and Associated with Increased Oxidative Stress. <i>Oxidative Medicine and Cellular Longevity</i> , 2021, 2021, 1-17.	1.9	6
25	Circulating CTRP7 Is a Potential Predictor for Metabolic Syndrome. <i>Frontiers in Endocrinology</i> , 2021, 12, 774309.	1.5	7
26	A Novel Immune and Stroma Related Prognostic Marker for Invasive Breast Cancer in Tumor Microenvironment: A TCGA Based Study. <i>Frontiers in Endocrinology</i> , 2021, 12, 774244.	1.5	5
27	Circulating complement C1q tumor necrosis factor-related protein isoform 5 levels are low in type 2 diabetes patients and reduced by dapagliflozin. <i>Journal of Diabetes Investigation</i> , 2020, 11, 88-95.	1.1	6
28	Increased plasma osteopontin levels are associated with nonalcoholic fatty liver disease in patients with type 2 diabetes mellitus. <i>Cytokine</i> , 2020, 125, 154837.	1.4	11
29	Central Sfrp5 regulates hepatic glucose flux and VLDL-triglyceride secretion. <i>Metabolism: Clinical and Experimental</i> , 2020, 103, 154029.	1.5	17
30	Reducing NADPH Synthesis Counteracts Diabetic Nephropathy through Restoration of AMPK Activity in Type 1 Diabetic Rats. <i>Cell Reports</i> , 2020, 32, 108207.	2.9	12
31	Elevated Circulating Fetuin-B Levels Are Associated with Insulin Resistance and Reduced by GLP-1RA in Newly Diagnosed PCOS Women. <i>Mediators of Inflammation</i> , 2020, 2020, 1-12.	1.4	15
32	Circulating Levels of CILP2 Are Elevated in Coronary Heart Disease and Associated with Atherosclerosis. <i>Oxidative Medicine and Cellular Longevity</i> , 2020, 2020, 1-11.	1.9	7
33	Association between Visceral Fat and Bone Mineral Density in Both Male and Female Patients with Adult Growth Hormone Deficiency. <i>Biochemistry Research International</i> , 2020, 2020, 1-6.	1.5	1
34	Efficacy and safety of polyethylene glycol loxenatide as add-on to metformin in patients with type 2 diabetes: A multicentre, randomized, double-blind, placebo-controlled, phase 3b trial. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 2375-2383.	2.2	14
35	Adrenal artery ablation for primary aldosteronism without apparent aldosteronoma: An efficacy and safety, proof-of-principle trial. <i>Journal of Clinical Hypertension</i> , 2020, 22, 1618-1626.	1.0	18
36	Follistatin-like 1 as a Novel Adipomyokine Related to Insulin Resistance and Physical Activity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e4499-e4509.	1.8	25

#	ARTICLE	IF	CITATIONS
37	Serum Fetuin-A levels are increased and associated with insulin resistance in women with polycystic ovary syndrome. <i>BMC Endocrine Disorders</i> , 2020, 20, 67.	0.9	19
38	Comparison of Allogeneic Platelet-rich Plasma With Autologous Platelet-rich Plasma for the Treatment of Diabetic Lower Extremity Ulcers. <i>Cell Transplantation</i> , 2020, 29, 096368972093142.	1.2	22
39	DOCK5 regulates energy balance and hepatic insulin sensitivity by targeting mTORC1 signaling. <i>EMBO Reports</i> , 2020, 21, e49473.	2.0	16
40	Adipose Insulin Resistance and Circulating Betatrophin Levels in Women with PCOS. <i>BioMed Research International</i> , 2020, 2020, 1-9.	0.9	11
41	Activation of TRPV1 channel antagonizes diabetic nephropathy through inhibiting endoplasmic reticulum-mitochondria contact in podocytes. <i>Metabolism: Clinical and Experimental</i> , 2020, 105, 154182.	1.5	53
42	Inhibition of Mitochondrial Calcium Overload by SIRT3 Prevents Obesity- or Age-Related Whitening of Brown Adipose Tissue. <i>Diabetes</i> , 2020, 69, 165-180.	0.3	77
43	Effect of central JAZF1 on glucose production is regulated by the PI3Kâ€Aktâ€AMPK pathway. <i>FASEB Journal</i> , 2020, 34, 7058-7074.	0.2	16
44	LASS2 regulates hepatocyte steatosis by interacting with NDUFS2/OXPHOS related proteins. <i>Biochemical and Biophysical Research Communications</i> , 2020, 526, 871-879.	1.0	5
45	Impairment of Bitter Taste Sensor Transient Receptor Potential Channel M5-Mediated Aversion Aggravates High-Salt Intake and Hypertension. <i>Hypertension</i> , 2019, 74, 1021-1032.	1.3	14
46	Osteoprotegerin Promotes Liver Steatosis by Targeting the ERKâ€PPAR-Î³â€CD36 Pathway. <i>Diabetes</i> , 2019, 68, 1902-1914.	0.3	56
47	Circulating ANGPTL8 Is Associated with the Presence of Metabolic Syndrome and Insulin Resistance in Polycystic Ovary Syndrome Young Women. <i>Mediators of Inflammation</i> , 2019, 2019, 1-10.	1.4	10
48	Role of bone morphogenetic proteinâ€9 in the regulation of glucose and lipid metabolism. <i>FASEB Journal</i> , 2019, 33, 10077-10088.	0.2	35
49	Changes in whole metabolites after exenatide treatment in overweight/obese polycystic ovary syndrome patients. <i>Clinical Endocrinology</i> , 2019, 91, 508-516.	1.2	25
50	Dock5 controls the peripheral B cell differentiation via regulating BCR signaling and actin reorganization. <i>Cellular Immunology</i> , 2019, 337, 15-21.	1.4	3
51	Effective gene delivery of shBMP-9 using polyethyleneimine-based coreâ€shell nanoparticles in an animal model of insulin resistance. <i>Nanoscale</i> , 2019, 11, 2008-2016.	2.8	18
52	Gut ghrelin regulates hepatic glucose production and insulin signaling via a gut-brain-liver pathway. <i>Cell Communication and Signaling</i> , 2019, 17, 8.	2.7	16
53	Association of circulating BMP9 with coronary heart disease and hypertension in Chinese populations. <i>BMC Cardiovascular Disorders</i> , 2019, 19, 131.	0.7	7
54	CILP-2 is a novel secreted protein and associated with insulin resistance. <i>Journal of Molecular Cell Biology</i> , 2019, 11, 1083-1094.	1.5	19

#	ARTICLE	IF	CITATIONS
55	Sfrp5 interacts with Slurp1 to regulate the accumulation of triglycerides in hepatocyte steatosis model. <i>Biochemical and Biophysical Research Communications</i> , 2019, 512, 256-262.	1.0	8
56	Alteration of gut microbiota induced by DPP-4i treatment improves glucose homeostasis. <i>EBioMedicine</i> , 2019, 44, 665-674.	2.7	66
57	Circulating alarin concentrations are high in patients with type 2 diabetes and increased by glucagon-like peptide-1 receptor agonist treatment. <i>Medicine (United States)</i> , 2019, 98, e16428.	0.4	8
58	LASS2 inhibits proliferation and induces apoptosis in HepG2 cells by affecting mitochondrial dynamics, the cell cycle and the nuclear factor- κ B pathways. <i>Oncology Reports</i> , 2019, 41, 3005-3014.	1.2	6
59	Deficiency of Mitochondrial Glycerol 3-Phosphate Dehydrogenase Contributes to Hepatic Steatosis. <i>Hepatology</i> , 2019, 70, 84-97.	3.6	30
60	Circulating C1q/TNF-related protein isoform 15 is a marker for the presence of metabolic syndrome. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3085.	1.7	25
61	DPP-4 Inhibitors Improve Diabetic Wound Healing via Direct and Indirect Promotion of Epithelial-Mesenchymal Transition and Reduction of Scarring. <i>Diabetes</i> , 2018, 67, 518-531.	0.3	56
62	Circulating bone morphogenetic protein-9 levels are associated with hypertension and insulin resistance in humans. <i>Journal of the American Society of Hypertension</i> , 2018, 12, 372-380.	2.3	12
63	Myonectin Predicts the Development of Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 139-147.	1.8	52
64	Pioglitazone Improved Insulin Sensitivity and First Phase Insulin Secretion Among Obese and Lean People with Diabetes: A Multicenter Clamp Study. <i>Diabetes Therapy</i> , 2018, 9, 815-826.	1.2	11
65	LAMP3 regulates hepatic lipid metabolism through activating PI3K/Akt pathway. <i>Molecular and Cellular Endocrinology</i> , 2018, 470, 160-167.	1.6	44
66	Mitochondrial glycerol 3-phosphate dehydrogenase promotes skeletal muscle regeneration. <i>EMBO Molecular Medicine</i> , 2018, 10, .	3.3	24
67	High Circulating Alarin Levels Are Associated with Presence of Metabolic Syndrome. <i>Cellular Physiology and Biochemistry</i> , 2018, 51, 2041-2051.	1.1	9
68	C1q/TNF-Related Protein5 (CTRP5) as a Biomarker to Predict Metabolic Syndrome and Each of Its Components. <i>International Journal of Endocrinology</i> , 2018, 2018, 1-8.	0.6	9
69	Non-insulin determinant pathways maintain glucose homeostasis upon metabolic surgery. <i>Cell Discovery</i> , 2018, 4, 58.	3.1	8
70	Response to Letter to the Editor: "Myonectin Predicts the Development of Type 2 Diabetes". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 4040-4041.	1.8	2
71	TERT and Akt Are Involved in the Par-4-Dependent Apoptosis of Islet β Cells in Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2018, 2018, 1-13.	1.0	7
72	JAZF1 ameliorates age and diet-associated hepatic steatosis through SREBP-1c -dependent mechanism. <i>Cell Death and Disease</i> , 2018, 9, 859.	2.7	36

#	ARTICLE	IF	CITATIONS
73	JAZF1 Inhibits Adipose Tissue Macrophages and Adipose Tissue Inflammation in Diet-Induced Diabetic Mice. <i>BioMed Research International</i> , 2018, 2018, 1-10.	0.9	20
74	Efficacy of low-level light therapy for treatment of diabetic foot ulcer: A systematic review and meta-analysis of randomized controlled trials. <i>Diabetes Research and Clinical Practice</i> , 2018, 143, 215-224.	1.1	30
75	C1q/TNF-related protein-6 is associated with insulin resistance and the development of diabetes in Chinese population. <i>Acta Diabetologica</i> , 2018, 55, 1221-1229.	1.2	18
76	Deficiency of PKD2L1 (TRPP3) Exacerbates Pathological Cardiac Hypertrophy by Augmenting NCX1-Mediated Mitochondrial Calcium Overload. <i>Cell Reports</i> , 2018, 24, 1639-1652.	2.9	27
77	Polyethylene glycol loxanatide injections added to metformin effectively improve glycemic control and exhibit favorable safety in type 2 diabetic patients. <i>Journal of Diabetes</i> , 2017, 9, 158-167.	0.8	24
78	Duodenal GLP-1 signaling regulates hepatic glucose production through a PKC- δ -dependent neurocircuitry. <i>Cell Death and Disease</i> , 2017, 8, e2609-e2609.	2.7	29
79	Plasma Sfrp5 levels correlate with determinants of the metabolic syndrome in Chinese adults. <i>Diabetes/Metabolism Research and Reviews</i> , 2017, 33, e2896.	1.7	29
80	A novel role for the Kr μ ppel-like factor 14 on macrophage inflammatory response and atherosclerosis development. <i>Cardiovascular Pathology</i> , 2017, 27, 1-8.	0.7	23
81	Decreased circulating BMP-9 levels in patients with Type 2 diabetes is a signature of insulin resistance. <i>Clinical Science</i> , 2017, 131, 239-246.	1.8	37
82	Circulating bone morphogenetic protein-9 in relation to metabolic syndrome and insulin resistance. <i>Scientific Reports</i> , 2017, 7, 17529.	1.6	28
83	Global and Regional Effects of Bladder Cancer Risk Associated with Pioglitazone Therapy in Patients with Diabetes. <i>Scientific Reports</i> , 2017, 7, 15804.	1.6	13
84	Rictor positively regulates B cell receptor signaling by modulating actin reorganization via ezrin. <i>PLoS Biology</i> , 2017, 15, e2001750.	2.6	24
85	Circulating zinc- β -glycoprotein levels are low in newly diagnosed patients with metabolic syndrome and correlate with adiponectin. <i>Nutrition and Metabolism</i> , 2017, 14, 53.	1.3	27
86	Circulating betatrophin is associated with insulin resistance in humans: cross-sectional and interventional studies <i>in vivo</i> and <i>in vitro</i> . <i>Oncotarget</i> , 2017, 8, 96604-96614.	0.8	19
87	Association between Serum Cystatin C and Diabetic Foot Ulceration in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-7.	1.0	11
88	NAMPT knockdown attenuates atherosclerosis and promotes reverse cholesterol transport in ApoE KO mice with high-fat-induced insulin resistance. <i>Scientific Reports</i> , 2016, 6, 26746.	1.6	16
89	Circulating Zinc- β -glycoprotein levels and Insulin Resistance in Polycystic Ovary Syndrome. <i>Scientific Reports</i> , 2016, 6, 25934.	1.6	28
90	The natural logarithm of zinc- β -glycoprotein/HOMA-IR is a better predictor of insulin sensitivity than the product of triglycerides and glucose and the other lipid ratios. <i>Cytokine</i> , 2016, 79, 96-102.	1.4	32

#	ARTICLE	IF	CITATIONS
91	Efficacy and safety of linagliptin in Asian patients with type 2 diabetes mellitus inadequately controlled by metformin: A multinational 24-week, randomized clinical trial. <i>Journal of Diabetes</i> , 2016, 8, 229-237.	0.8	20
92	Sodium-Glucose Cotransporter 2 (SGLT2) Inhibitor Increases Circulating Zinc-β2-Glycoprotein Levels in Patients with Type 2 Diabetes. <i>Scientific Reports</i> , 2016, 6, 32887.	1.6	47
93	Adiponectin/(FPG-β2-Glycoproteins) as a predictor of insulin sensitivity and metabolic syndrome in patients with polycystic ovary syndrome. <i>Medicine (United States)</i> , 2016, 95, e5524.	0.4	6
94	Human C-reactive protein impedes entry of leptin into the CNS and attenuates its physiological actions in the CNS. <i>Biochemical Journal</i> , 2016, 473, 1215-1224.	1.7	5
95	Effects of sitagliptin on circulating zinc-β2-glycoprotein levels in newly diagnosed type 2 diabetes patients: a randomized trial. <i>European Journal of Endocrinology</i> , 2016, 174, 147-155.	1.9	25
96	Exenatide once-weekly injection for the treatment of type 2 diabetes in Chinese patients: current perspectives. <i>Therapeutics and Clinical Risk Management</i> , 2015, 11, 1153.	0.9	4
97	Elevated Circulating Levels of Irisin and the Effect of Metformin Treatment in Women With Polycystic Ovary Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 1485-1493.	1.8	78
98	Krüppel-like factor 14 increases insulin sensitivity through activation of PI3K/Akt signal pathway. <i>Cellular Signalling</i> , 2015, 27, 2201-2208.	1.7	48
99	Serum C1q/TNF-related protein-3 (CTRP3) levels are decreased in obesity and hypertension and are negatively correlated with parameters of insulin resistance. <i>Diabetology and Metabolic Syndrome</i> , 2015, 7, 33.	1.2	60
100	Glypican-4 is increased in human subjects with impaired glucose tolerance and decreased in patients with newly diagnosed type 2 diabetes. <i>Acta Diabetologica</i> , 2014, 51, 981-990.	1.2	29
101	Silencing of FGF21 expression promotes hepatic gluconeogenesis and glycogenolysis by regulation of the STAT3/SOCS3 signal. <i>FEBS Journal</i> , 2014, 281, 2136-2147.	2.2	28
102	Overexpression of juxtaposed with another zinc finger gene1 reduces proinflammatory cytokine release via inhibition of stress-activated protein kinases and nuclear factor-κB. <i>FEBS Journal</i> , 2014, 281, 3193-3205.	2.2	23
103	Transcutaneous oxygen pressure (TcPO2): A novel diagnostic tool for peripheral neuropathy in type 2 diabetes patients. <i>Diabetes Research and Clinical Practice</i> , 2014, 105, 336-343.	1.1	24
104	Hypothalamic Nesfatin-1/NUCB2 Knockdown Augments Hepatic Gluconeogenesis That Is Correlated With Inhibition of mTOR-STAT3 Signaling Pathway in Rats. <i>Diabetes</i> , 2014, 63, 1234-1247.	0.3	50
105	Serum retinol-binding protein 4 levels are elevated but do not contribute to insulin resistance in newly diagnosed Chinese hypertensive patients. <i>Diabetology and Metabolic Syndrome</i> , 2014, 6, 72.	1.2	15
106	Effects of glucagon-like peptide-1 agents on left ventricular function: Systematic review and meta-analysis. <i>Annals of Medicine</i> , 2014, 46, 664-671.	1.5	12
107	Overexpression of JAZF1 protected ApoE-deficient mice from atherosclerosis by inhibiting hepatic cholesterol synthesis via CREB-dependent mechanisms. <i>International Journal of Cardiology</i> , 2014, 177, 100-110.	0.8	26
108	Efficacy and Safety of Aldose Reductase Inhibitor for the Treatment of Diabetic Cardiovascular Autonomic Neuropathy: Systematic Review and Meta-Analysis. <i>PLoS ONE</i> , 2014, 9, e87096.	1.1	35

#	ARTICLE	IF	CITATIONS
109	Zinc- α 2-Glycoprotein Is Associated With Insulin Resistance in Humans and Is Regulated by Hyperglycemia, Hyperinsulinemia, or Liraglutide Administration. <i>Diabetes Care</i> , 2013, 36, 1074-1082.	4.3	83
110	Circulating Sfrp5 Is a Signature of Obesity-Related Metabolic Disorders and Is Regulated by Glucose and Liraglutide in Humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 290-298.	1.8	103
111	The effects of fibroblast growth factor-21 knockdown and over-expression on its signaling pathway and glucose-lipid metabolism in vitro. <i>Molecular and Cellular Endocrinology</i> , 2012, 348, 21-26.	1.6	45
112	Nesfatin-1 Action in the Brain Increases Insulin Sensitivity Through Akt/AMPK/TORC2 Pathway in Diet-Induced Insulin Resistance. <i>Diabetes</i> , 2012, 61, 1959-1968.	0.3	112
113	Liraglutide Increases FGF-21 Activity and Insulin Sensitivity in High Fat Diet and Adiponectin Knockdown Induced Insulin Resistance. <i>PLoS ONE</i> , 2012, 7, e48392.	1.1	48
114	Elevated circulating vaspin levels were decreased by rosiglitazone therapy in T2DM patients with poor glycemic control on metformin alone. <i>Cytokine</i> , 2011, 56, 399-402.	1.4	9
115	Liraglutide Prevents Hypoadiponectinemia-Induced Insulin Resistance and Alterations of Gene Expression Involved in Glucose and Lipid Metabolism. <i>Molecular Medicine</i> , 2011, 17, 1168-1178.	1.9	49
116	The role of JAZF1 on lipid metabolism and related genes in vitro. <i>Metabolism: Clinical and Experimental</i> , 2011, 60, 523-530.	1.5	48
117	Overexpression of visfatin/PBEF/Nampt alters whole-body insulin sensitivity and lipid profile in rats. <i>Annals of Medicine</i> , 2009, 41, 311-320.	1.5	56
118	Circulating preptin levels in normal, impaired glucose tolerance, and type 2 diabetic subjects. <i>Annals of Medicine</i> , 2009, 41, 52-56.	1.5	55
119	The adipose triglyceride lipase, adiponectin and visfatin are downregulated by tumor necrosis factor- α (TNF- α) in vivo. <i>Cytokine</i> , 2009, 45, 12-19.	1.4	58
120	Short-term pioglitazone treatment prevents free fatty acid-induced hepatic insulin resistance in normal rats: Possible role of the resistin and adiponectin. <i>Biochemical and Biophysical Research Communications</i> , 2006, 339, 1190-1196.	1.0	17
121	Effects of free fatty acids on plasma resistin and insulin resistance in awake rats. <i>Metabolism: Clinical and Experimental</i> , 2005, 54, 1142-1146.	1.5	16
122	In vivo effects of phosphodiesterase III inhibitors on glucose metabolism and insulin sensitivity. <i>Journal of the Chinese Medical Association</i> , 2003, 66, 210-6.	0.6	4
123	GPHB5 Is a Biomarker in Women With Metabolic Syndrome: Results From Cross-Sectional and Intervention Studies. <i>Frontiers in Endocrinology</i> , 0, 13, .	1.5	2