
Gianandrea La Porta

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5489979/publications.pdf Version: 2024-02-01



CIANANDREA LA PORTA

#	Article	IF	CITATIONS
1	Italian odonates in the Pandora's box: A comprehensive DNA barcoding inventory shows taxonomic warnings at the Holarctic scale. Molecular Ecology Resources, 2021, 21, 183-200.	4.8	37
2	Characterization of long-range transported bioaerosols in the Central Mediterranean. Science of the Total Environment, 2021, 763, 143010.	8.0	17
3	Population Ecology and Genetic Diversity of the Invasive Alien Species Procambarus clarkii in Lake Trasimeno (Italy). Biology, 2021, 10, 1059.	2.8	6
4	Movement and demography of Southern damselfly (<i>Coenagrion mercuriale</i> , Odonata) in a Mediterranean lotic ecosystem. Ethology Ecology and Evolution, 2020, 32, 107-121.	1.4	2
5	Evaluation of geochemical baselines and metal enrichment factor values through high ecological quality reference points: a novel methodological approach. Environmental Science and Pollution Research, 2020, 27, 930-940.	5.3	7
6	Use of Larval Morphological Deformities in Chironomus plumosus (Chironomidae: Diptera) as an Indicator of Freshwater Environmental Contamination (Lake Trasimeno, Italy). Water (Switzerland), 2020, 12, 1.	2.7	31
7	Small heat shock proteins in the amphibian Pelophylax bergeri: Cloning and characterization of Hsp27 and Hsp30 cDNAs and their expression analysis in ex vivo skin exposed to abiotic stresses. Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology, 2019, 235, 90-101.	1.8	5
8	Investigation on the declining Southern Damselfly (Coenagrion mercuriale, Odonata) in a Mediterranean population: survival rate and population size. Journal of Insect Conservation, 2019, 23, 667-675.	1.4	2
9	Morphologic and genetic variability in the Barbus fishes (Teleostei, Cyprinidae) of Central Italy. Zoologica Scripta, 2019, 48, 289-301.	1.7	13
10	Longitudinal study of Amphibiocystidium sp. infection in a natural population of the Italian stream frog (Rana italica). Parasitology, 2019, 146, 903-910.	1.5	4
11	Morphological survey as powerful detection tool of pure and local phenotypes in <i>Salmo trutta</i> complex. Knowledge and Management of Aquatic Ecosystems, 2019, , 48.	1.1	9
12	Introgression despite protection: the case of native brown trout in Natura 2000 network in Italy. Conservation Genetics, 2019, 20, 343-356.	1.5	31
13	Ecology and conservation of the Mediterranean trout in the central Apennines (Italy). Journal of Limnology, 2019, 78, .	1.1	6
14	Population status of the native <i>Cottus gobio</i> after removal of the alien <i>Salmo trutta</i> : a case-study in two Mediterranean streams (Italy). Knowledge and Management of Aquatic Ecosystems, 2018, , 22.	1.1	7
15	Airborne bacteria and persistent organic pollutants associated with an intense Saharan dust event in the Central Mediterranean. Science of the Total Environment, 2018, 645, 401-410.	8.0	38
16	Sampling and comparing odonate assemblages by means of exuviae: statistical and methodological aspects. Journal of Insect Conservation, 2017, 21, 207-218.	1.4	13
17	A geographic distribution data set of biodiversity in Italian freshwaters. Biogeographia, 2016, 31, .	0.5	4
18	Long-term changes in diet and trophic niche of the European wildcat (Felis silvestris silvestris) in Italy. Mammal Research, 2016, 61, 109-119.	1.3	27

#	Article	IF	CITATIONS
19	Characterization of the Skin Microbiota in Italian Stream Frogs (<i>Rana italica</i>) Infected and Uninfected by a Cutaneous Parasitic Disease. Microbes and Environments, 2015, 30, 262-269.	1.6	38
20	Short-term cadmium exposure induces stress responses in frog (Pelophylax bergeri) skin organ culture. Ecotoxicology and Environmental Safety, 2015, 122, 221-229.	6.0	16
21	Weak effects of habitat type on susceptibility to invasive freshwater species: an Italian case study. Aquatic Conservation: Marine and Freshwater Ecosystems, 2014, 24, 841-852.	2.0	17
22	Effects of selenium supplemented diets on growth and condition indexes in juvenile red swamp crayfish, Procambarus clarkii. Environmental Toxicology and Pharmacology, 2013, 36, 484-492.	4.0	16
23	Proposed standard weight (Ws) equation for European perch (Perca fluviatilis Linnaeus, 1758). Journal of Applied Ichthyology, 2012, 28, 34-39.	0.7	8
24	Proposed standard mass equations for European chub <i>Leuciscus cephalus</i> in Italy. Journal of Fish Biology, 2011, 78, 1890-1899.	1.6	7
25	BIOLOGY OF PROCAMBARUS CLARKII (GIRARD, 1852) IN LAKE TRASIMENO. Knowledge and Management of Aquatic Ecosystems: an International Journal on Aquatic Ecosystems, 2006, , 1155-1168.	0.4	40
26	The consortium of the sponge Ephydatia fluviatilis (L.) living on the common reed Phragmites australis in Lake Piediluco (central Italy). Hydrobiologia, 2004, 520, 165-178.	2.0	17
27	Larval development of Aeshna cyanea (Müller, 1764) (Odonata: Aeshnidae) in Central Italy. Hydrobiologia, 2001, 457, 149-154.	2.0	12
28	Larval development ofLibellula depressa(Odonata, Libellulidae) from pools in central Italy. Italian Journal of Zoology, 2000, 67, 343-348.	0.6	13
29	Variations in the fish community in lake Piediluco (Italy) caused by changes in the lake's trophic status and the introduction of alien species. Journal of Applied Ichthyology, 0, 26, 53-59.	0.7	17