Jelena M. Aćmović

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5499444/publications.pdf

Version: 2024-02-01

16 papers	196 citations	1040018 9 h-index	1058452 14 g-index
16 all docs	16 docs citations	16 times ranked	263 citing authors

#	Article	IF	CITATIONS
1	The interplay between copper(II), human serum albumin, fatty acids, and carbonylating agent interferes with Cys 34 thiol reactivity and copper binding. Journal of Biological Inorganic Chemistry, 2019, 24, 61-70.	2.6	6
2	Quantification of total content of non-esterified fatty acids bound to human serum albumin. Journal of Pharmaceutical and Biomedical Analysis, 2016, 129, 43-49.	2.8	4
3	Binding of enterolactone and enterodiol to human serum albumin: increase of cysteine-34 thiol group reactivity. Food and Function, 2016, 7, 1217-1226.	4.6	16
4	HSA carbonylation with methylglyoxal and the binding/release of copper(ii) ions. Metallomics, 2015, 7, 1431-1438.	2.4	8
5	How the sialylation level of serum N-acetyl- \hat{l}^2 -D-glucosaminidase a form in type 1 diabetes mellitus influences its activity?. Journal of the Serbian Chemical Society, 2014, 79, 1491-1503.	0.8	O
6	The influence of fatty acids on determination of human serum albumin thiol group. Analytical Biochemistry, 2014, 448, 50-57.	2.4	16
7	Fatty acids binding to human serum albumin: Changes of reactivity and glycation level of Cysteine-34 free thiol group with methylglyoxal. Chemico-Biological Interactions, 2014, 224, 42-50.	4.0	30
8	The efficiency of compounds with \hat{l}_{\pm} -amino- \hat{l}_{\pm} -mercapto-ethane group in protection of human serum albumin carbonylation and cross-linking with methylglyoxal. Molecular BioSystems, 2014, 10, 2166-2175.	2.9	7
9	Monitoring of the human serum albumin carbonylation level through determination of guanidino group content. Analytical Biochemistry, 2013, 433, 162-167.	2.4	6
10	Improving the reliability of human serum albumin-thiol group determination. Analytical Biochemistry, 2013, 439, 17-22.	2.4	15
11	Method for monitoring of the protein amino group changes during carbonylation. Clinical Biochemistry, 2011, 44, 994-999.	1.9	19
12	Influence of the microenvironment of thiol groups in low molecular mass thiols and serum albumin on the reaction with methylglyoxal. Chemico-Biological Interactions, 2010, 188, 21-30.	4.0	20
13	The role of the thiol group in protein modification with methylglyoxal. Journal of the Serbian Chemical Society, 2009, 74, 867-883.	0.8	28
14	The possibility of determining N-acetyl- \hat{l}^2 -d-glucosaminidase isoenzymes under alkaline conditions. Clinical Biochemistry, 2005, 38, 384-389.	1.9	11
15	Influence of pigments and pH of urine on the determination ofN-acetyl-β-D-glucosaminidase activity with 2-methoxy-4-(2′-nitrovinyl)-phenyl-N-acetyl-β-D-glucosaminide. Journal of Clinical Laboratory Analysis, 2005, 19, 260-266.	2.1	1
16	Reactivity of IGF binding protein-3 isoforms towards concanavalin A in healthy adults and subjects with cirrhosis. Addiction Biology, 2003, 8, 81-88.	2.6	9