

# Kazuya Nagasawa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5502653/publications.pdf>

Version: 2024-02-01

161  
papers

1,829  
citations

331670  
21  
h-index

395702  
33  
g-index

163  
all docs

163  
docs citations

163  
times ranked

1270  
citing authors

#	ARTICLE	IF	CITATIONS
1	Thermal limits and ocean migrations of sockeye salmon ( <i>Oncorhynchus nerka</i> ): long-term consequences of global warming. Canadian Journal of Fisheries and Aquatic Sciences, 1998, 55, 937-948.	1.4	123
2	Trophic relations in the subarctic North Pacific ecosystem:possible feeding effect from pink salmon. Marine Ecology - Progress Series, 1997, 150, 75-85.	1.9	99
3	Small subunit rDNA and Bayesian inference reveal <i>Pectenophilus formatus</i> (Copepoda: incertae sedis) as highly transformed Mytilicolidae, and support assignment of Chondracanthidae and Xarifiidae to Lichomolgoidea (Cyclopoida). Biological Journal of the Linnean Society, 2006, 87, 403-425.	1.6	74
4	Recent Changes in Age and Size of Chum Salmon ( <i>Oncorhynchus keta</i> ) in the North Pacific Ocean and Possible Causes. Canadian Journal of Fisheries and Aquatic Sciences, 1993, 50, 290-295.	1.4	67
5	Stock enhancement programme for black sea bream, <i>Acanthopagrus schlegelii</i> (Bleeker), in Hiroshima Bay, Japan: a review. Aquaculture Research, 2008, 39, 1307-1315.	1.8	56
6	Distribution of Pelagic Elasmobranchs Caught by Salmon Research Gillnets in the North Pacific. Fisheries Science, 1996, 62, 860-865.	1.6	52
7	Stock enhancement program for black sea bream ( <i>Acanthopagrus schlegelii</i> ) in Hiroshima Bay: Monitoring the genetic effects. Aquaculture, 2008, 276, 36-43.	3.5	46
8	Epipelagic nekton of the North Pacific Subarctic and Transition Zones. Progress in Oceanography, 1999, 43, 365-397.	3.2	41
9	Interannual variability in stock abundance of the neon flying squid, <i>Ommastrephes bartramii</i> , in the North Pacific Ocean during 1979-1998: impact of driftnet fishing and oceanographic conditions. Fisheries Oceanography, 2000, 9, 163-170.	1.7	41
10	Observations on five species of philometrid nematodes from marine fishes in Japan. Systematic Parasitology, 1998, 40, 67-80.	1.1	32
11	Title is missing!. Hydrobiologia, 2001, 453/454, 411-416.	2.0	30
12	Some anisakid nematodes from marine fishes of Japan and the North Pacific Ocean. Journal of Natural History, 2000, 34, 1555-1574.	0.5	28
13	Regional synchrony in return rates of chum salmon ( <i>Oncorhynchus keta</i> ) in Japan in relation to coastal temperature and size at release. Fisheries Research, 2009, 95, 14-27.	1.7	28
14	Prevalence and abundance of <i>Lepeophtheirus salmonis</i> (Copepoda: Caligidae) on high-seas salmon and trout in the North Pacific Ocean.. Nippon Suisan Gakkaishi, 1987, 53, 2151-2156.	0.1	27
15	Long-term variations in abundance of Pacific herring ( <i>Clupea pallasi</i> ) in Hokkaido and Sakhalin related to changes in environmental conditions. Progress in Oceanography, 2001, 49, 551-564.	3.2	27
16	Caryophyllidean tapeworms (Platyhelminthes: Eucestoda) from freshwater fishes in Japan. Folia Parasitologica, 2001, 48, 275-288.	1.3	27
17	Two New Huffmanela Species, <i>H. japonica</i> n. sp. and <i>H. shikokuensis</i> n. sp. (Nematoda:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 10 0.7 26		
18	Kudoa intestinalis n. sp. (Myxosporea: Multivalvulida) from the Intestinal Musculature of the Striped Mullet, <i>Mugil cephalus</i> , from Japan. Journal of Parasitology, 1993, 79, 190.	0.7	25

#	ARTICLE	IF	CITATIONS
19	Larval Anisakid Nematodes of Japanese Common Squid ( <i>Todarodes pacificus</i> ) from the Sea of Japan. <i>Journal of Parasitology</i> , 1995, 81, 69.	0.7	25
20	Spatial Variability of Living Coccolithophore Distribution in the Western Subarctic Pacific and Western Bering Sea. <i>Journal of Oceanography</i> , 2004, 60, 505-515.	1.7	24
21	On the origin of <i>Anguillicoloides crassus</i> , the invasive nematode of anguillid eels. <i>Aquatic Invasions</i> , 2012, 7, 443-453.	1.6	24
22	Relationship between zooplankton abundance and the early marine life history of juvenile chum salmon <i>Oncorhynchus keta</i> in eastern Hokkaido, Japan. <i>Fisheries Science</i> , 2009, 75, 303-316.	1.6	21
23	Redescription of four species of <i>Hatschekia</i> (Copepoda: Siphonostomatoida: Hatschekiidae) parasitic on tetraodontiform fishes from Japan. <i>Zootaxa</i> , 2009, 2110, 1-21.	0.5	17
24	A mass occurrence of human infection with <i>Diplogonoporus grandis</i> (Cestoda: Diphyllobothriidae) in Shizuoka Prefecture, central Japan. <i>Parasitology International</i> , 2002, 51, 73-79.	1.3	16
25	A COLLECTION OF FISH LEECHES (HIRUDINIDA: PISCICOLIDAE) FROM JAPAN AND SURROUNDING WATERS, INCLUDING REDESCRIPTIONS OF THREE SPECIES. <i>Journal of Parasitology</i> , 2007, 93, 875-883.	0.7	16
26	Two species of philometrid nematodes (Philometridae) from marine fishes off Japan, including <i>Philometroides branchiostegi</i> sp. n. from <i>Branchiostegus japonicus</i> (Malacanthidae). <i>Folia Parasitologica</i> , 2012, 59, 71-78.	1.3	16
27	Redescription of <i>Raphidascaris gigi</i> Fujita, 1928 (Nematoda: Anisakidae), a parasite of freshwater fishes in Japan. <i>Systematic Parasitology</i> , 2002, 52, 193-198.	1.1	15
28	Gyrodactylus medaka n. sp. (Monogenea: Gyrodactylidae) parasitic on wild and laboratory-reared medaka <i>Oryzias latipes</i> (Beloniformes: Adrianichthyidae) in Japan. <i>Parasitology International</i> , 2018, 67, 651-658.	1.3	15
29	Development of Ultrasonic Telemetry Technique for Investigating the Magnetic of Salmonids. <i>Fisheries Science</i> , 1996, 62, 698-704.	1.6	15
30	Morphology of <i>Pectenophilus ornatus</i> , New Genus, New Species, a Copepod Parasite of the Japanese Scallop <i>Patinopecten yessoensis</i> . <i>Journal of Crustacean Biology</i> , 1988, 8, 31.	0.8	14
31	Ergasilid copepods (Poecilostomatoida) in plankton samples from Hokkaido, Japan, with reconsideration of the taxonomic status of <i>Limnoncaea</i> Kokubo, 1914. <i>Journal of Natural History</i> , 2004, 38, 471-498.	0.5	14
32	Parasitic copepods of marine fish cultured in Japan: a review. <i>Journal of Natural History</i> , 2015, 49, 2891-2903.	0.5	14
33	Host Utilization by the Salmon Louse <i>Lepeophtheirus salmonis</i> (Copepoda: Caligidae) in the Sea of Japan. <i>Journal of Parasitology</i> , 1993, 79, 127.	0.7	13
34	Rediscovery in Japan and host association of <i>Salmincola carpionis</i> (Copepoda: Lernaeopodidae), a parasite of wild and reared freshwater salmonids. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 1995, 52, 178-185.	1.4	13
35	Two remarkable nematodes from sharks in Japan. <i>Journal of Natural History</i> , 2000, 34, 1-13.	0.5	13
36	Occurrence of <i>Caligus lalandei</i> Barnard, 1948 (Copepoda, Siphonostomatoida) on Amberjacks ( <i>Seriola</i> ) Tj ETQq0 0 0 rgBT /Overlock 10	0.7	13

#	ARTICLE	IF	CITATIONS
37	Double-netting: an alternative approach for the recovery of parasitic copepods from finfishes. Journal of Natural History, 2013, 47, 529-541.	0.5	13
38	Tidal-Flat Macrofauna as Diets of the Japanese Eel <i>Anguilla japonica</i> in Western Japan, with a Note on the Occurrence of a Parasitic Nematode <i>Heliconema anguillae</i> in Eel Stomachs. Zoological Science, 2016, 33, 50.	0.7	13
39	First record of ostracods as natural intermediate hosts of <i>Anguillicola crassus</i> , a pathogenic swimbladder parasite of eels <i>Anguilla</i> spp.. Diseases of Aquatic Organisms, 2005, 66, 171-173.	1.0	12
40	Reduction in size-at-age of black sea bream ( <i>Acanthopagrus schlegelii</i> ) following intensive releases of cultured juveniles in Hiroshima Bay, Japan. Fisheries Research, 2009, 99, 130-133.	1.7	12
41	Effects of juvenile size at release and early marine growth on adult return rates for Hokkaido chum salmon ( <i>Oncorhynchus keta</i> ) in relation to sea surface temperature. Fisheries Oceanography, 2011, 20, 278-293.	1.7	12
42	Long-term changes in the population size and geographical distribution of <i>Pennella</i> sp. (Copepoda) on the saury, <i>Cololabis saira</i> , in the western North Pacific Ocean and adjacent seas. Hydrobiologia, 1988, 167-168, 571-577.	2.0	11
43	New data on the morphology of <i>Philonema oncorhynchi</i> Kuitunen-Ekbaum, 1933 (Nematoda: Tj ETQq1 1 0.784314 rgBT /Overlock 10 T0 11 Parasitology, 1999, 43, 67-74.	1.1	11
44	Larval anisakid nematodes from four species of squid (Cephalopoda: Teuthoidea) from the central and western North Pacific Ocean. Journal of Natural History, 2002, 36, 883-891.	0.5	11
45	The copepod genus <i>Hatschekia</i> Poche, 1902 (Siphonostomatoida: Hatschekiidae) from triggerfishes (Pisces: Tetraodontiformes: Balistidae) from off the Ryukyu Islands, Japan, with descriptions of eleven new species. Zootaxa, 2010, 2478, 1.	0.5	11
46	Sperm Utilization Patterns and Reproductive Success in Captive Loggerhead Turtles (<i>Caretta</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 30 0.6 11		
47	Chalimus Stages of <i>Caligus latigenitalis</i> (Copepoda: Caligidae) Parasitic on Blackhead Seabream from Japanese Waters, with Discussion of Terminology Used for Developmental Stages of Caligids. Journal of Parasitology, 2011, 97, 221-236.	0.7	11
48	Age, Growth, and Feeding Habits of Lanternfish, &lt;i&gt; <i>Stenobrachius leucopsarus</i> &lt;/i&gt; (Myctophidae), Collected from the Near-surface Layer in the Bering Sea. Fisheries Science, 1999, 65, 11-15.	1.6	10
49	Effects of <i>Acanthocephalus</i> sp. (Acanthocephala: Echinorhynchidae) on the body size and reproduction of isopods ( <i>Asellus hilgendorfi</i> ). Limnology, 2003, 4, 43-46.	1.5	10
50	Three new species of <i>Hatschekia</i> Poche, 1902 (Copepoda: Siphonostomatoida: Hatschekiidae) parasitic on <i>Abalistes filamentosus</i> (Pisces: Tetraodontiformes: Balistidae) from off Okinawa, Japan. Systematic Parasitology, 2009, 74, 225-237.	1.1	10
51	Can research on the early marine life stage of juvenile chum salmon <i>Oncorhynchus keta</i> forecast returns of adult salmon? A case study from eastern Hokkaido, Japan. Fisheries Science, 2010, 76, 909-920.	1.6	10
52	Three new species of <i>Hatschekia</i> Poche, 1902 (Copepoda: Siphonostomatoida: Hatschekiidae) parasitic on boxfishes (Pisces: Tetraodontiformes: Aracanidae and Ostraciidae) in Japanese waters. Systematic Parasitology, 2010, 75, 147-158.	1.1	10
53	Three new species of the family Pennellidae (Copepoda: Siphonostomatoida) from gobiid fishes (Actinopterygii: Perciformes) in coastal waters of the western Pacific Ocean. Zootaxa, 2010, 2687, 29.	0.5	10
54	Four new species of splanchnotrophid copepods (Poecilostomatoida) parasitic on doridacean nudibranchs (Gastropoda, Opistobranchia) from Japan, with proposition of one new genus. ZooKeys, 2012, 247, 1-29.	1.1	10

#	ARTICLE	IF	CITATIONS
55	Redescription of the fish parasite <i>Nerocila japonica</i> Schioedte & Meinert, 1881 (Crustacea: Isopoda) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Parasitology, 2012, 81, 147-157.	1.1	10
56	Growth and reproductive cycle of <i>&lt; i&gt;Hypermastus tokunagai&lt;/i&gt;</i> (Caenogastropoda: Eulimidae), an ectoparasite of the sand dollar <i>&lt; i&gt;Scaphechinus mirabilis&lt;/i&gt;</i> (Clypeasteroida: Scutellidae) in the Seto Inland Sea, Japan. Journal of the Marine Biological Association of the United Kingdom, 2013, 93, 1041-1051.	0.8	10
57	A new species of <i>Dactylogyrus</i> (Monogenea: Dactylogyridae) parasitic on an endangered freshwater fish, <i>Rhodeus atremius</i> , endemic to Japan. Parasitology International, 2016, 65, 483-487.	1.3	10
58	&lt;i&gt; <i>Tachaea chinensis</i> &lt;/i&gt; (Isopoda: Corallanidae), an ectoparasite of freshwater shrimps and prawns, from western Japan, with a list of its known collection localities and hosts. Crustacean Research, 2018, 47, 73-88.	0.8	10
59	The finding of <i>Pennella</i> sp. (Copepoda: Pennellidae) on the saury, <i>Cololabis saira</i> , in the western and central North Pacific Ocean and the Okhotsk Sea.. Fish Pathology, 1984, 18, 205-208.	0.7	9
60	Occurrence and effects of <i>Haemobaphes diceraus</i> (Copepoda:Pennellidae) on brown sole <i>Limanda herzensteini</i> off the Okhotsk coast of Hokkaido.. Nippon Suisan Gakkaishi, 1987, 53, 991-994.	0.1	9
61	Occurrence and effects of the parasitic copepod <i>Salmincola carpionis</i> on salmonids in the Nikko District, central Japan. Journal of Marine Systems, 1998, 15, 269-272.	2.1	9
62	WHY INFESTATION BY LEPEOPHTHEIRUS SALMONIS (COPEPODA:CALICIDAE) IS NOT A PROBLEM IN THE COHO SALMON FARMING INDUSTRY IN JAPAN. Journal of Crustacean Biology, 2001, 21, 954-960.	0.8	9
63	Estimation of sperm storage duration in captive loggerhead turtles ( <i>Caretta caretta</i> ). Journal of Experimental Marine Biology and Ecology, 2013, 439, 136-142.	1.5	9
64	Four new species of <i>&lt; i&gt;Colobomatus&lt;/i&gt;</i> (Copepoda: Philichthyidae) parasitic in the lateral line system of marine finfishes captured off the Ryukyu Islands, Japan, with redescriptions of <i>&lt; i&gt;Colobomatus collettei&lt;/i&gt;</i> Cressey, 1977 and <i>&lt; i&gt;Colobomatus pupa&lt;/i&gt;</i> Izawa, 1974. Journal of Natural History, 2013, 47, 563-580.	0.5	9
65	<i>Rhabdochona angusticaudata</i> sp. n. (Nematoda: Rhabdochonidae) from the Japanese eel <i>Anguilla japonica</i> , and new records of some other nematodes from inland fishes in Japan. Folia Parasitologica, 2018, 65, .	1.3	9
66	Prevalence of visceral adhesions in sockeye salmon, <i>Oncorhynchus nerka</i> , in the central North Pacific Ocean.. Fish Pathology, 1985, 20, 313-321.	0.7	9
67	Effects of <i>Pectenophilus ornatus</i> (Copepoda) on the Biomass of Cultured Japanese Scallop <i>Patinopecten yessoensis</i> . Journal of Parasitology, 1992, 78, 552.	0.7	8
68	Possible Life Cycle of <i>Longicollum pagrosomi</i> , an Acanthocephalan Parasite of Cultured Red Sea Bream.. Fish Pathology, 1996, 31, 235-236.	0.7	8
69	Sarcotretes Longirostris N. Sp. (Copepoda: Pennellidae) Parasitic on Bluefin Driftfish ( <i>Psenes</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Biology, 2007, 27, 116-120.	0.8	8
70	Host Size- and Habitat-Dependent Intensity of <i>Heliconema longissimum</i> (Nematoda: Physalopteridae) In the Japanese Eel ( <i>Anguilla japonica</i> ). Journal of Parasitology, 2011, 97, 994-998.	0.7	8
71	A new species of parasitic copepod, <i>Sarcotretes umitakae</i> sp. n. (Siphonostomatoida, Pennellidae), on the rattail (Actinopterygii, Macrouridae) from the East China Sea, Japan. ZooKeys, 2012, 246, 1-10.	1.1	8
72	New records of philometrids (Nematoda: Philometridae) from marine fishes off Japan, including description of <i>Philometra kidakoi</i> sp. n. and <i>Congerinema japonicum</i> gen. et sp. n.. Folia Parasitologica, 2019, 66, .	1.3	8

#	ARTICLE	IF	CITATIONS
73	The identity of <i>Limnoncaea diuncata</i> Kokubo, 1914 (Copepoda: Poecilostomatoida) from Hokkaido, Japan, with the relegation of <i>Diergasilus</i> Do, 1981 to a junior synonym of <i>Thersitina</i> Norman, 1905. Systematic Parasitology, 2004, 57, 35-44.	1.1	7
74	A New Species of Capillaria (Nematoda: Capillariidae) from the Intestine of the Marine Fish <i>Acanthopagrus schlegelii</i> schlegelii (Sparidae) from Japan. Journal of Parasitology, 2010, 96, 771-774.	0.7	7
75	<i>Caligus Macarovi</i> (Copepoda, Caligidae) from Pacific Bluefin Tuna, <i>Thunnus Orientalis</i> , Cultured in Japan. Crustaceana, 2011, 84, 1145-1147.	0.3	7
76	Four Alien Monogeneans, Including <i>Trinigyrus peregrinus</i>; n. sp., Parasitic on the Invasive Armored Catfish <i>Pterygoplichthys disjunctivus</i>; (Siluriformes: Loricariidae) from Okinawa-jima Island, Okinawa Prefecture, Japan. Species Diversity, 2016, 21, 95-104.	0.4	7
77	First Record of <i>Norileca indica</i> (Isopoda, Cymothoidae) Parasitic on <i>Selar crumenophthalmus</i> and <i>Decapterus kurroides</i> (Perciformes, Carangidae) in the Philippines. Comparative Parasitology, 2017, 84, 60-63.	0.4	7
78	New Record of a Marine Fish Parasite <i>Nerocila trichiura</i> (Crustacea: Isopoda: Cymothoidae) from Japan, with its Confirmed Distribution in the Western North Pacific Ocean. Species Diversity, 2019, 24, 195-201.	0.4	7
79	<i>Heliconema anguillae</i> Yamaguti, 1935, a physalopterid nematode found in Japanese eels: taxonomic resurrection with a note on the third-stage larva from intertidal crabs in western Japan. Folia Parasitologica, 2015, 62, .	1.3	7
80	Ecological factors influencing the infection levels of salmonids by <i>Acanthocephalus opsariichthydis</i> (Acanthocephala: Echinorhynchidae) in Lake Yunoko, Japan.. Fish Pathology, 1983, 18, 53-60.	0.7	6
81	Seasonal and regional occurrence of <i>Acanthocephalus</i> sp. (Acanthocephala: Echinorhynchidae) in fishes and isopods ( <i>Asellus hilgendorfi</i> ) in a lake system in northern Japan. Limnology, 2002, 3, 143-150.	1.5	6
82	An East-West Comparison of the Transition Zone Coastal Pelagic Nekton of the North Pacific Ocean. Journal of Oceanography, 2003, 59, 415-434.	1.7	6
83	<i>Paeonocanthus antarcticensis</i> (Hewitt, 1965): a rare copepod parasite of bathypelagic fish, <i>Bathylagus antarcticus</i> GÄ¼nther, from the Antarctic Ocean. Zoological Science, 2003, 20, 1315-1321.	0.7	6
84	A new species of parasitic copepod of the genus <i>Hatschekia</i> Poche, 1902 (Siphonostomatoida:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 30 Systematic Parasitology, 2010, 76, 53-58.	1.1	6
85	<i>Pseudohatschekiidae</i> , a new family for <i>Pseudohatschekia branchiostegi</i> Yamaguti, 1939 (Crustacea:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 30 Japanese waters, with descriptions of the early developmental stages of <i>P. branchiostegi</i> . Systematic Parasitology, 2010, 77, 215-231.	1.1	6
86	Two new species of the copepod <i>Hatschekia</i> Poche, 1902 (Siphonostomatoida: Hatschekiidae) from angelfishes (Pisces: Perciformes: Pomacanthidae) collected during the KUMEJIMA 2009 Expedition. Zootaxa, 2012, 3367, 49.	0.5	6
87	<i>Choniomyzon inflatus</i> n. sp. (Crustacea: Copepoda: Nicothoidae) associated with <i>Ibacus novemdentatus</i> (Crustacea: Decapoda: Scyllaridae) from Japanese waters. Systematic Parasitology, 2013, 84, 157-165.	1.1	6
88	<i>Nerocila phaiopleura</i>;, a cymothoid isopod parasitic on Pacific bluefin tuna, <i>Thunnus orientalis</i>;, cultured in Japan. Crustacean Research, 2017, 46, 95-101.	0.8	6
89	New host and ocean records for <i>Driocephalus cerebrinoxius</i> (Sphyriidae, Siphonostomatoida) and a reconsideration of phylogeny within Sphyriidae. Acta Ichthyologica Et Piscatoria, 2006, 36, 1-9.	0.7	6
90	<i>Trebias shiinoi</i> n. sp. (Trebidae: Siphonostomatoida: Copepoda) from Uteri and Embryos of the Japanese Angelshark ( <i>Squatina japonica</i> ) and the Clouded Angelshark ( <i>Squatina nebulosa</i> ), and Redescription of <i>Trebias longicaudatus</i> . Journal of Parasitology, 1998, 84, 1218.	0.7	5

#	ARTICLE	IF	CITATIONS
91	Copepod Parasites of the Fatheads (Pisces, Psychrolutidae) Their Implication on the Phylogenetic Relationships of Psychrolutid Genera. <i>Zoological Science</i> , 2005, 22, 411-425.	0.7	5
92	Host and Seasonal Distribution of Fish Acanthocephalans from the Lake Biwa Basin, Japan. <i>Comparative Parasitology</i> , 2007, 74, 244-253.	0.4	5
93	Helminths from the giant mottled eel <i>Anguilla marmorata</i> Quoy & Gaimard in Japan, with a description of <i>Acanthocephalus longiacanthus</i> n. sp. (Acanthocephala: Echinorhynchidae). <i>Systematic Parasitology</i> , 2014, 88, 91-102.	1.1	5
94	Infection of <i>Brachiella thynni</i> (Copepoda, Lernaeopodidae) on Pacific bluefin tuna, <i>Thunnus orientalis</i> (Actinopterygii, Scombridae), cultured in Japan. <i>Crustaceana</i> , 2015, 88, 945-948.	0.3	5
95	A new species of <i>Leposphilus</i> Hesse, 1866 (Copepoda: Philichthyidae) parasitic in the interorbital canals of the whitemouth croaker <i>Micropogonias furnieri</i> (Desmarest) (Sciaenidae) off Brazil with an amended diagnosis of the genus. <i>Systematic Parasitology</i> , 2016, 93, 501-515.	1.1	5
96	<i>Mothocyia collettei</i> Bruce, 1986 (Isopoda, Cymothoidae), a marine fish parasite new to Japan. <i>Crustaceana</i> , 2017, 90, 613-616.	0.3	5
97	Alien Gill Parasites of the Silver Carp <i>Hypophthalmichthys molitrix</i> (Cypriniformes: Cyprinidae) in Tochigi Prefecture, Central Japan. <i>Species Diversity</i> , 2020, 25, 61-73.	0.4	5
98	<i>Gobioecetes longibasais</i> n. sp. (Monogenea: Dactylogyridae) from <i>Rhinogobius similis</i> Gill (Perciformes: Gobiidae) from Okinawa-jima Island, the Ryukyu Archipelago, southern Japan, with a new host record for <i>Gobioecetes biwaensis</i> Ogawa & Itoh, 2017. <i>Systematic Parasitology</i> , 2020, 97, 193-200.	1.1	5
99	&lt;i&gt;Haplocotyle japonica&lt;/i&gt; n. gen., n. sp. (Monogenea: Microbothriidae) Parasitic on &lt;i&gt;Rhinobatos hynnicephalus&lt;/i&gt; (Elasmobranchii: Rajiformes: Rhinobatidae) in Japanese Waters. <i>Species Diversity</i> , 2017, 22, 117-125.	0.4	5
100	Crustacean Parasites of the Saury, <i>Cololabis saira</i> : a Review. <i>Fish Pathology</i> , 1984, 19, 57-63.	0.7	5
101	Infection of the Salmon Louse <i>Lepeophtheirus salmonis</i> (Copepoda: Caligidae) on Seawater-Farmed Salmonids in Northern Honshu, Japan.. <i>Fish Pathology</i> , 1993, 28, 143-144.	0.7	5
102	Role of <i>Leptodora kindti</i> (Cladocera:Leptodoridae) in the life cycle of <i>Raphidascaris biwakoensis</i> (Nematoda:Anisakidae), a fish parasite in Lake Biwa, Japan. <i>Diseases of Aquatic Organisms</i> , 1998, 32, 157-160.	1.0	4
103	Influence of presoaking conditions of spat collectors in seawater containing adult Pacific oysters ( <i>Crassostrea gigas</i> ) on their larval settlement. <i>Nippon Suisan Gakkaishi</i> , 2008, 74, 1017-1023.	0.1	4
104	Four New Species of <i>Peniculisa</i> Wilson, 1917 (Copepoda: Siphonostomatoida: Pennellidae) Parasitic on Coastal Marine Fishes in Japanese Waters. <i>Journal of Parasitology</i> , 2010, 96, 689-702.	0.7	4
105	<i>Echinorhynchus hexagrammi</i> Baeva, 1965 (Acanthocephala: Echinorhynchidae) from marine fishes off Hokkaido, Japan, with morphological observations and new host records. <i>Systematic Parasitology</i> , 2011, 80, 35-40.	1.1	4
106	Influence of host habitat on the occurrence of gill monogeneans <i>Pseudodactylogyrus</i> spp. on wild Japanese eels <i>Anguilla japonica</i> . <i>Diseases of Aquatic Organisms</i> , 2012, 100, 43-49.	1.0	4
107	Spatial distribution of <i>Neoergasilus japonicus</i> (Copepoda: Ergasilidae) on the fins of bluegill (<i>Lepomis macrochirus</i>). <i>Journal of Natural History</i> , 2013, 47, 543-552.	0.5	4
108	<i>Bivaginogyrus obscurus</i> (Monogenea: Dactylogyridae), a Gill Parasite of Two Cyprinids <i>Pseudorasbora pumila</i> and <i>Pseudorasbora parva</i> , New to Japan. <i>Species Diversity</i> , 2014, 19, 167-171.	0.4	4

#	ARTICLE	IF	CITATIONS
109	Temporal changes in demersal fish assemblage structure in the East China Sea and the Yellow Sea. Nippon Suisan Gakkaishi, 2015, 81, 429-437.	0.1	4
110	Species composition of crabs and change in abundance of portunid crabs in the East China Sea and the Yellow Sea from the 1990s to 2000s. Nippon Suisan Gakkaishi, 2015, 81, 43-51.	0.1	4
111	New records of Caligidae (Copepoda, Siphonostomatoidea) from the Philippines. Zootaxa, 2016, 4174, 237.	0.5	4
112	First Record of the Fish Louse <i>Argulus quadristriatus</i> (Branchiura: Arguloida: Argulidae) from Japanese Waters, with Three New Host Records. Species Diversity, 2017, 22, 37-44.	0.4	4
113	Some Spirurid Nematodes (Spirurida) from Freshwater and Brackish-Water Fishes in Okinawa Prefecture, Japan, with Descriptions of Two New Species. Acta Parasitologica, 2021, 66, 163-177.	1.1	4
114	The genus Hatschekia (Copepoda: Hatschekiidae) from pufferfishes (Tetraodontiformes: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 547 Td (redescription of H. pholas. Folia Parasitologica, 2013, 60, 61-74.	1.3	4
115	Infection with <i>Argulus japonicus</i> on Amur Catfish Silurus asotus in an Aquarium. Japanese Journal of Limnology, 2011, 72, 211-214.	0.1	4
116	&lt;i&gt;Argulus japonicus&lt;/i&gt; (Branchiura: Argulidae) parasitic on largemouth bass &lt;i&gt;Micropterus&lt;/i&gt; &lt;i&gt;salmoides&lt;/i&gt; in Japan, with the morphology of the adult female of the argulid. Crustacean Research, 2021, 50, 119-129.	0.8	4
117	Acanthocephalus lucidus van CLEAVE (Acanthocephala: Echinorhynchidae) from cultured rainbow trout <i>Salmo gairdneri</i> Richardson.. Nippon Suisan Gakkaishi, 1981, 47, 1153-1156.	0.1	3
118	New Host and Ocean Records for the Parasitic Copepod Bobkabata kabatabobbus (Lernaeosoleidae:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 0.6	0.6	3
119	New Hosts, Including Marine Fishes Cultured in Japan, of Limnotrachelobdella okae (Hirudinida:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 0.4	0.4	3
120	Species of Taeniacanthus Sumpf, 1871 (Crustacea: Copepoda: Taeniacanthidae) parasitic on boxfishes (Tetraodontiformes: Aracanidae and Ostraciidae) from the Indo-West Pacific region, with descriptions of two new species. Systematic Parasitology, 2011, 80, 141-157.	1.1	3
121	Mate Selection Based on Genetic Relatedness of Loggerhead Turtles in Captivity. Chelonian Conservation and Biology, 2012, 11, 214-222.	0.6	3
122	Redescription of <i>Lepeophtheirus acutus</i> Heegaard, 1943 (Copepoda: Caligidae) parasitic on two elasmobranch hosts off Okinawa-jima Island, Japan. Journal of Natural History, 2013, 47, 581-596.	0.5	3
123	Redescription of <i>Pseudacanthocanthopsis secunda</i> (Yamaguti and Yamasu, 1960) (Copepoda:) Tj ETQq1 1 0.784314 rgBT /Overlock 0.5	0.5	3
124	First Japanese Record of Heteropriapulus heterotylus (Monogenea: Dactylogyridae), from the Alien Catfish Pterygoplichthys disjunctivus (Siluriformes: Loricariidae) in Okinawa. Species Diversity, 2013, 18, 281-284.	0.4	3
125	Year-round infections and complicated demography of a food-transmitted parasite Heliconema anguillae implying the feeding activity of Japanese eels in saline habitats. Fisheries Science, 2016, 82, 863-871.	1.6	3
126	Life-cycle of Choniomyzon inflatus Wakabayashi, Otake, Tanaka & Nagasawa, 2013 (Copepoda:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 1.1	1.1	3
	adult male. Systematic Parasitology, 2016, 93, 145-157.		

#	ARTICLE	IF	CITATIONS
127	Pseudocycnus appendiculatus Heller, 1865, the third copepod species parasitic on Pacific bluefin tuna, <i>Thunnus orientalis</i> (Temminck & Schlegel, 1844), cultured in Japan, with its first Japanese record. <i>Crustaceana</i> , 2017, 90, 511-515.	0.3	3
128	A new species of Furcohaptor Bijukumar & Kearn, 1996 (Monogenea: Diplectanidae) parasitic on <i>Cynoglossus robustus</i> GÄ¼nther (Pleuronectiformes: Cynoglossidae) in the Seto Inland Sea, Japan, with comments on its systematic position and an amended generic diagnosis of Furcohaptor. <i>Systematic Parasitology</i> , 2017, 94, 907-913.	1.1	3
129	Redescription of <i>Dichelyne (Cucullanellus) jialaris</i> (Nematoda: Cucullanidae), a parasite of seabreams (Perciformes: Sparidae) in East Asia. <i>Acta Parasitologica</i> , 2018, 63, 802-807.	1.1	3
130	&lt;i&gt;Norileca&lt;/i&gt; aff. &lt;i&gt;triangulata&lt;/i&gt; (Isopoda: Cymothoidae), a branchial cavity parasite of the African sailfin flyingfish, &lt;i&gt;Parexocoetus mento &lt;/i&gt; (Beloniformes:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 622 Research, 2021, 50, 55-64.	0.8	3
131	< i>Dactylogyrus oryziasi</i> n. sp. (Monogenea: Dactylogyridae) from < i>Oryzias latipes</i> (Beloniformes: Adrianichthyidae) in Japan. <i>Species Diversity</i> , 2017, 22, 1-5.	0.4	3
132	Infection with the Parasitic Copepod <i>Clavella parva</i> (Lernaeopodidae) in Gold-eye Rockfish <i>Sebastes thompsoni</i> Broodstock in Japan. <i>Fish Pathology</i> , 2008, 43, 55-60.	0.7	3
133	Description of <i>Raphidascaris (Ichthyascaris) multipapillata</i> sp. nov., Redescription of <i>Hysterothylacium simile</i> Li et al., 2013 and New Records of Some Other Ascaridoid and Seuratoid Nematodes from Marine Fishes in Japan. <i>Acta Parasitologica</i> , 2021, 66, 461-474.	1.1	3
134	Distribution and Bivalve Hosts of the Parasitic Copepod <i>Pectenophilus Ornatus</i> Nagasawa, Bresciani, and LÃœtzen in Japan. <i>Journal of Crustacean Biology</i> , 1993, 13, 544-550.	0.8	2
135	Description of <i>Naobranchia Denticis</i> N. Sp. (Copepoda, Siphonostomatoidea, Lernaeopodidae) Parasitic on <i>Dentex Hypselosomus</i> (Teleostei, Sparidae) from Japanese Waters, with a Key to the Species of Naobranchia. <i>Crustaceana</i> , 2011, 84, 461-476.	0.3	2
136	The spotfin burrfish ( <i>Chilomycterus reticulatus</i> ), a new host record for <i>Cymothoa pulchra</i> (Isopoda,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	0.3	2
137	Three Species of <i>Dactylogyrus</i> (Monogenea: Dactylogyridae) Parasitic on Japanese Seabass <i>Lateolabrax japonicus</i> (Perciformes: Lateolabracidae) in Japan, with New Country Records for <i>Dactylogyrus gotoi</i> and <i>Dactylogyrus kikuchii</i> . <i>Species Diversity</i> , 2014, 19, 71-79.	0.4	2
138	Crimson seabream <i>Evynnis tumifrons</i> (Temminck & Schlegel, 1843) (Perciformes, Sparidae), a new host for Ceratothoa verrucosa (Schioedte & Meinert, 1883) (Isopoda, Cymothoidae). <i>Crustaceana</i> , 2016, 89, 1229-1232.	0.3	2
139	Occurrence of praniza larvae of a gnathiid (Isopoda, Gnathiidae) on wild-caught orange-spotted grouper, <i>Epinephelus coioides</i> (Hamilton, 1822) (Perciformes, Serranidae), in Lingayen Gulf, Philippines. <i>Crustaceana</i> , 2017, 90, 101-110.	0.3	2
140	First Records of < i>Trachelastes sachalinensis</i> (Copepoda: Lernaeopodidae), a Fin Parasite of Cyprinids, from Japan. <i>Species Diversity</i> , 2017, 22, 167-173.	0.4	2
141	<i>Euryphorus brachypterus</i> (Gerstaecker, 1853), a caligid copepod infecting Pacific bluefin tuna, <i>Thunnus orientalis</i> (Temminck & Schlegel, 1844), cultured in Japan. <i>Crustaceana</i> , 2018, 91, 251-255.	0.3	2
142	<i>Nerocila japonica</i> Schioedte & Meinert, 1881 (Isopoda, Cymothoidae) found in a Japanese culture of <i>Girella</i> Leonina (Richardson, 1846) (Actinopterygii, Kyphosidae). <i>Crustaceana</i> , 2018, 91, 375-377.	0.3	2
143	Some nematodes from eels (Anguilliformes: Anguillidae) in Japan, with descriptions of two new species. <i>Systematic Parasitology</i> , 2020, 97, 691-712.	1.1	2
144	Long-term changes in the population size and geographical distribution of <i>Pennella</i> sp. (Copepoda) on the saury, <i>Cololabis saira</i> , in the western North Pacific Ocean and adjacent seas*, 1988, , 571-577.	2	2

#	ARTICLE	IF	CITATIONS
145	Marine Fish Parasite <i>&lt;Argulus caecus&gt;</i> (Crustacea: Branchiura: Argulidae) Accidentally Collected from a Fixed Net Caught Squid in Northern Japan. Species Diversity, 2021, 26, 289-296.	0.4	2
146	New Geographical Records of Three Gill Monogeneans (Platyhelminthes) Parasitic on <i>&lt;Pseudorasbora parva&gt;</i> (Cypriniformes: Cyprinidae) in Japan. Species Diversity, 2016, 21, 1-7.	0.4	2
147	<i>&lt;Nerocila phaiopleura&gt;</i> (Isopoda: Cymothoidae): a new record from Ibaraki Prefecture, central Japan, with a discussion of its distribution in Japanese waters. Crustacean Research, 2020, 49, 41-47.	0.8	2
148	Host Body Size, Prevalence, and Breeding Season of the Microparasitic Snail <i>Pelseneeria castanea</i> on the Sea Urchin <i>Strongylocentrotus nudus</i> in Coastal Waters of the Sea of Japan off Western Hokkaido, Japan. Japanese Journal of Benthology, 2006, 61, 16-20.	0.1	1
149	A new species of <i>Pseudomacrochiron Reddiah</i> , 1969 (Crustacea: Copepoda: Macrochironidae) associated with scyphistomae of the moon jellyfish <i>Aurelia</i> sp. (Cnidaria: Scyphozoa) off Japan. Systematic Parasitology, 2012, 81, 125-134.	1.1	1
150	Benthic shrimp assemblages associated with water masses in the East China Sea and the Yellow Sea. Nippon Suisan Gakkaishi, 2013, 79, 987-1001.	0.1	1
151	A review of the <i>Taeniacanthus balistae</i> species group (Crustacea: Copepoda: Taeniacanthidae), with descriptions of two new species. Zootaxa, 2016, 4174, 212.	0.5	1
152	<i>Nippoparasitus unoashicola</i> , a new genus and species of philobennid copepod (Cyclopoida) parasitic on the Pacific sugar limpet, <i>Patelloidea saccharina</i> (Linnaeus, 1758) (Patellogastropoda: Lottiidae) from the intertidal zone of eastern Japan. Zootaxa, 2016, 4174, 386.	0.5	1
153	<i>&amp;lt;i&amp;gt;Ceratothoa verrucosa&amp;lt;/i&amp;gt;</i> (Isopoda: Cymothoidae) attached by <i>&amp;lt;i&amp;gt;Choricotyle elongata&amp;lt;/i&amp;gt;</i> (Platyhelminthes: Monogenea) in the mouth cavity of red seabream <i>&amp;lt;i&amp;gt;Pagrus major&amp;lt;/i&amp;gt;</i> . Crustacean Research, 2018, 47, 5-8.	0.8	1
154	<i>Salsuginus seculus</i> (Monogenea: Ancyrocephalidae) Newly Recorded from Japan, Infecting the Introduced Mosquitofish <i>Gambusia affinis</i> . Species Diversity, 2014, 19, 173-178.	0.4	1
155	New host and third western Pacific Ocean records for <i>&lt;Nerocila trichiura&gt;</i> (Isopoda: Tj ETQq1 1 0.784314 rgBT 0.8 /Overlock 10 Tf 50		
156	Do Visceral Adhesions Affect the Growth of Sockeye Salmon in the North Pacific Ocean and Bering Sea?. Fish Pathology, 2022, 57, 41-48.	0.7	1
157	Distribution of the rough pomfret, <i>Taractes asper</i> (Bramidae), in the central North Pacific Ocean. Ichthyological Research, 1998, 45, 419-423.	0.8	0
158	Influence of larval size and feeding on the ratio of their settlement to spat collectors presoaked in seawater containing adult Pacific oysters <i>Crassostrea gigas</i> . Nippon Suisan Gakkaishi, 2009, 75, 1036-1041.	0.1	0
159	Contribution of Ju-shey Ho to the systematics of symbiotic copepods of Japan. Journal of Natural History, 2013, 47, 517-527.	0.5	0
160	<i>&lt;Doridicola similis&gt;</i> (Copepoda: Poecilostomatoida: Rhynchosomatidae) Associated with <i>&lt;Sepioteuthis&gt;</i> sp. (Cephalopoda: Myopsida: Loliginidae), New to Japan. Species Diversity, 2017, 22, 161-166.	0.4	0
161	<i>&amp;lt;i&amp;gt;Argulus matuii&amp;lt;/i&amp;gt;</i> (Branchiura: Argulidae) parasitic on yellowfin seabream <i>&amp;lt;i&amp;gt;Acanthopagrus latus&amp;lt;/i&amp;gt;</i> in Japan, with a note on the body coloration in an ethanol-preserved argulid specimen. Crustacean Research, 2022, 51, 31-38.	0.8	0