

Kazuya Nagasawa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5502653/publications.pdf>

Version: 2024-02-01

161
papers

1,829
citations

331670

21
h-index

395702

33
g-index

163
all docs

163
docs citations

163
times ranked

1270
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>Argulus matuii</i> (Branchiura: Argulidae) parasitic on yellowfin seabream <i>Acanthopagrus latus</i> in Japan, with a note on the body coloration in an ethanol-preserved argulid specimen. <i>Crustacean Research</i> , 2022, 51, 31-38.	0.8	0
2	Do Visceral Adhesions Affect the Growth of Sockeye Salmon in the North Pacific Ocean and Bering Sea?. <i>Fish Pathology</i> , 2022, 57, 41-48.	0.7	1
3	Some Spirurid Nematodes (Spirurida) from Freshwater and Brackish-Water Fishes in Okinawa Prefecture, Japan, with Descriptions of Two New Species. <i>Acta Parasitologica</i> , 2021, 66, 163-177.	1.1	4
4	<i>Norileca</i> aff. <i>triangulata</i> (Isopoda: Cymothoidae), a branchial cavity parasite of the African sailfin flyingfish, <i>Parexocoetus mento</i> (Beloniformes: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 622 Research, 2021, 50, 55-64.	0.8	3
5	Marine Fish Parasite <i>Argulus caecus</i> (Crustacea: Branchiura: Argulidae) Accidentally Collected from a Fixed Net Caught Squid in Northern Japan. <i>Species Diversity</i> , 2021, 26, 289-296.	0.4	2
6	<i>Argulus japonicus</i> (Branchiura: Argulidae) parasitic on largemouth bass <i>Micropterus salmoides</i> in Japan, with the morphology of the adult female of the argulid. <i>Crustacean Research</i> , 2021, 50, 119-129.	0.8	4
7	Description of <i>Raphidascaris</i> (<i>Ichthyascaris</i>) <i>multipapillata</i> sp. nov., Redescription of <i>Hysterothylacium simile</i> Li et al., 2013 and New Records of Some Other Ascaridoid and Seuratoid Nematodes from Marine Fishes in Japan. <i>Acta Parasitologica</i> , 2021, 66, 461-474.	1.1	3
8	Some nematodes from eels (Anguilliformes: Anguillidae) in Japan, with descriptions of two new species. <i>Systematic Parasitology</i> , 2020, 97, 691-712.	1.1	2
9	Alien Gill Parasites of the Silver Carp <i>Hypophthalmichthys molitrix</i> (Cypriniformes: Cyprinidae) in Tochigi Prefecture, Central Japan. <i>Species Diversity</i> , 2020, 25, 61-73.	0.4	5
10	<i>Gobioecetes longibasais</i> n. sp. (Monogenea: Dactylogyridae) from <i>Rhinogobius similis</i> Gill (Perciformes: Gobiidae) from Okinawa-jima Island, the Ryukyu Archipelago, southern Japan, with a new host record for <i>Gobioecetes biwaensis</i> Ogawa & Itoh, 2017. <i>Systematic Parasitology</i> , 2020, 97, 193-200.	1.1	5
11	<i>Nerocila phaiopleura</i> (Isopoda: Cymothoidae): a new record from Ibaraki Prefecture, central Japan, with a discussion of its distribution in Japanese waters. <i>Crustacean Research</i> , 2020, 49, 41-47.	0.8	2
12	New host and third western Pacific Ocean records for <i>Nerocila trichiura</i> (Isopoda: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 302 Td (0.8	1
13	New Record of a Marine Fish Parasite <i>Nerocila trichiura</i> (Crustacea: Isopoda: Cymothoidae) from Japan, with its Confirmed Distribution in the Western North Pacific Ocean. <i>Species Diversity</i> , 2019, 24, 195-201.	0.4	7
14	New records of philometrids (Nematoda: Philometridae) from marine fishes off Japan, including description of <i>Philometra kidakoi</i> sp. n. and <i>Congerinema japonicum</i> gen. et sp. n.. <i>Folia Parasitologica</i> , 2019, 66, .	1.3	8
15	<i>Euryphorus brachypterus</i> (Gerstaecker, 1853), a caligid copepod infecting Pacific bluefin tuna, <i>Thunnus orientalis</i> (Temminck & Schlegel, 1844), cultured in Japan. <i>Crustaceana</i> , 2018, 91, 251-255.	0.3	2
16	<i>Nerocila japonica</i> Schioedte & Meinert, 1881 (Isopoda: Cymothoidae) found in a Japanese culture of <i>Girella leonina</i> (Richardson, 1846) (Actinopterygii: Kyphosidae). <i>Crustaceana</i> , 2018, 91, 375-377.	0.3	2
17	<i>Tachaea chinensis</i> (Isopoda: Corallanidae), an ectoparasite of freshwater shrimps and prawns, from western Japan, with a list of its known collection localities and hosts. <i>Crustacean Research</i> , 2018, 47, 73-88.	0.8	10
18	Redescription of <i>Dichelyne</i> (<i>Cucullanellus</i>) <i>jialaris</i> (Nematoda: Cucullanidae), a parasite of seabreams (Perciformes: Sparidae) in East Asia. <i>Acta Parasitologica</i> , 2018, 63, 802-807.	1.1	3

#	ARTICLE	IF	CITATIONS
19	<i>Cyrodactylus medaka</i> n. sp. (Monogenea: Cyrodactylidae) parasitic on wild and laboratory-reared medaka <i>Oryzias latipes</i> (Beloniformes: Adrianichthyidae) in Japan. <i>Parasitology International</i> , 2018, 67, 651-658.	1.3	15
20	<i>Ceratothoa verrucosa</i> (Isopoda: Cymothoidae) attached by <i>Choricotyle elongata</i> (Platyhelminthes: Monogenea) in the mouth cavity of red seabream <i>Pagrus major</i>. <i>Crustacean Research</i> , 2018, 47, 5-8.	0.8	1
21	<i>Rhabdochona angusticaudata</i> sp. n. (Nematoda: Rhabdochonidae) from the Japanese eel <i>Anguilla japonica</i> , and new records of some other nematodes from inland fishes in Japan. <i>Folia Parasitologica</i> , 2018, 65, .	1.3	9
22	<i>Mothocya collettei</i> Bruce, 1986 (Isopoda, Cymothoidae), a marine fish parasite new to Japan. <i>Crustaceana</i> , 2017, 90, 613-616.	0.3	5
23	<i>Pseudocycnus appendiculatus</i> Heller, 1865, the third copepod species parasitic on Pacific bluefin tuna, <i>Thunnus orientalis</i> (Temminck & Schlegel, 1844), cultured in Japan, with its first Japanese record. <i>Crustaceana</i> , 2017, 90, 511-515.	0.3	3
24	Occurrence of praniza larvae of a gnathiid (Isopoda, Gnathiidae) on wild-caught orange-spotted grouper, <i>Epinephelus coioides</i> (Hamilton, 1822) (Perciformes, Serranidae), in Lingayen Gulf, Philippines. <i>Crustaceana</i> , 2017, 90, 101-110.	0.3	2
25	First Record of <i>Norileca indica</i> (Isopoda, Cymothoidae) Parasitic on <i>Selar crumenophthalmus</i> and <i>Decapterus kurroides</i> (Perciformes, Carangidae) in the Philippines. <i>Comparative Parasitology</i> , 2017, 84, 60-63.	0.4	7
26	A new species of Furcuhaptor <i>Bijukumar & Kearns</i> , 1996 (Monogenea: Diplectanidae) parasitic on <i>Cynoglossus robustus</i> (Pleuronectiformes: Cynoglossidae) in the Seto Inland Sea, Japan, with comments on its systematic position and an amended generic diagnosis of Furcuhaptor. <i>Systematic Parasitology</i> , 2017, 94, 907-913.	1.1	3
27	<i>Nerocila phaiopleura</i>, a cymothoid isopod parasitic on Pacific bluefin tuna, <i>Thunnus orientalis</i>, cultured in Japan. <i>Crustacean Research</i> , 2017, 46, 95-101.	0.8	6
28	<i>Doridicola similis</i> (Copepoda: Poecilostomatoida: Rhynchomolgidae) Associated with <i>Sepioteuthis</i> sp. (Cephalopoda: Myopsida: Loliginidae), New to Japan. <i>Species Diversity</i> , 2017, 22, 161-166.	0.4	0
29	First Records of <i>Tracheliastes sachalinensis</i> (Copepoda: Lernaepodidae), a Fin Parasite of Cyprinids, from Japan. <i>Species Diversity</i> , 2017, 22, 167-173.	0.4	2
30	First Record of the Fish Louse <i>Argulus quadristriatus</i> (Branchiura: Arguloida: Argulidae) from Japanese Waters, with Three New Host Records. <i>Species Diversity</i> , 2017, 22, 37-44.	0.4	4
31	<i>Dactylogyrus oryziasi</i> n. sp. (Monogenea: Dactylogyridae) from <i>Oryzias latipes</i> (Beloniformes: Adrianichthyidae) in Japan. <i>Species Diversity</i> , 2017, 22, 1-5.	0.4	3
32	<i>Haplocotyle japonica</i> n. gen., n. sp. (Monogenea: Microbothriidae) Parasitic on <i>Rhinobatos hynnicephalus</i> (Elasmobranchii: Rajiformes: Rhinobatidae) in Japanese Waters. <i>Species Diversity</i> , 2017, 22, 117-125.	0.4	5
33	Four Alien Monogeneans, Including <i>Trinigyris peregrinus</i> n. sp., Parasitic on the Invasive Armored Catfish <i>Pterygoplichthys disjunctivus</i> (Siluriformes: Loricariidae) from Okinawa-jima Island, Okinawa Prefecture, Japan. <i>Species Diversity</i> , 2016, 21, 95-104.	0.4	7
34	Year-round infections and complicated demography of a food-transmitted parasite <i>Heliconema anguillae</i> implying the feeding activity of Japanese eels in saline habitats. <i>Fisheries Science</i> , 2016, 82, 863-871.	1.6	3
35	A new species of <i>Leposphilus</i> Hesse, 1866 (Copepoda: Philichthyidae) parasitic in the interorbital canals of the whitemouth croaker <i>Micropogonias furnieri</i> (Desmarest) (Sciaenidae) off Brazil with an amended diagnosis of the genus. <i>Systematic Parasitology</i> , 2016, 93, 501-515.	1.1	5
36	A review of the <i>Taeniacanthus balistae</i> species group (Crustacea: Copepoda: Taeniacanthidae), with descriptions of two new species. <i>Zootaxa</i> , 2016, 4174, 212.	0.5	1

#	ARTICLE	IF	CITATIONS
37	New records of Caligidae (Copepoda, Siphonostomatoida) from the Philippines. <i>Zootaxa</i> , 2016, 4174, 237.	0.5	4
38	<i>Nippoparasitus unoashicola</i> , a new genus and species of philoblenniid copepod (Cyclopoida) parasitic on the Pacific sugar limpet, <i>Patelloida saccharina</i> (Linnaeus, 1758) (Patellogastropoda: Lottiidae) from the intertidal zone of eastern Japan. <i>Zootaxa</i> , 2016, 4174, 386.	0.5	1
39	A new species of <i>Dactylogyrus</i> (Monogenea: Dactylogyridae) parasitic on an endangered freshwater fish, <i>Rhodeus atremius atremius</i> , endemic to Japan. <i>Parasitology International</i> , 2016, 65, 483-487.	1.3	10
40	Crimson seabream <i>Eyvynnis tumifrons</i> (Temminck & Schlegel, 1843) (Perciformes, Sparidae), a new host for <i>Ceratothoa verrucosa</i> (Schioedte & Meinert, 1883) (Isopoda, Cymothoidae). <i>Crustaceana</i> , 2016, 89, 1229-1232.	0.3	2
41	Life-cycle of <i>Choniomyzon inflatus</i> Wakabayashi, Otake, Tanaka & Nagasawa, 2013 (Copepoda: Tj ETQq1 1 0.784314 rgBT /Ove adult male. <i>Systematic Parasitology</i> , 2016, 93, 145-157.	1.1	3
42	Tidal-Flat Macrobenthos as Diets of the Japanese Eel <i>Anguilla japonica</i> in Western Japan, with a Note on the Occurrence of a Parasitic Nematode <i>Heliconema anguillae</i> in Eel Stomachs. <i>Zoological Science</i> , 2016, 33, 50.	0.7	13
43	New Geographical Records of Three Gill Monogeneans (Platyhelminthes) Parasitic on <i>Pseudorasbora parva</i> (Cypriniformes: Cyprinidae) in Japan. <i>Species Diversity</i> , 2016, 21, 1-7.	0.4	2
44	Temporal changes in demersal fish assemblage structure in the East China Sea and the Yellow Sea. <i>Nippon Suisan Gakkaishi</i> , 2015, 81, 429-437.	0.1	4
45	Species composition of crabs and change in abundance of portunid crabs in the East China Sea and the Yellow Sea from the 1990s to 2000s. <i>Nippon Suisan Gakkaishi</i> , 2015, 81, 43-51.	0.1	4
46	Infection of <i>Brachiella thynni</i> (Copepoda, Lernaepodidae) on Pacific bluefin tuna, <i>Thunnus orientalis</i> (Actinopterygii, Scombridae), cultured in Japan. <i>Crustaceana</i> , 2015, 88, 945-948.	0.3	5
47	Parasitic copepods of marine fish cultured in Japan: a review. <i>Journal of Natural History</i> , 2015, 49, 2891-2903.	0.5	14
48	<i>Heliconema anguillae</i> Yamaguti, 1935, a physalopterid nematode found in Japanese eels: taxonomic resurrection with a note on the third-stage larva from intertidal crabs in western Japan. <i>Folia Parasitologica</i> , 2015, 62, .	1.3	7
49	Three Species of <i>Dactylogyrus</i> (Monogenea: Dactylogyridae) Parasitic on Japanese Seabass <i>Lateolabrax japonicus</i> (Perciformes: Lateolabracidae) in Japan, with New Country Records for <i>Dactylogyrus gotoi</i> and <i>Dactylogyrus kikuchii</i> . <i>Species Diversity</i> , 2014, 19, 71-79.	0.4	2
50	<i>Bivaginogyrus obscurus</i> (Monogenea: Dactylogyridae), a Gill Parasite of Two Cyprinids <i>Pseudorasbora pumila pumila</i> and <i>Pseudorasbora parva</i> , New to Japan. <i>Species Diversity</i> , 2014, 19, 167-171.	0.4	4
51	Helminths from the giant mottled eel <i>Anguilla marmorata</i> Quoy & Gaimard in Japan, with a description of <i>Acanthocephalus longiacanthus</i> n. sp. (Acanthocephala: Echinorhynchidae). <i>Systematic Parasitology</i> , 2014, 88, 91-102.	1.1	5
52	<i>Salsuginus seculus</i> (Monogenea: Ancyrocephalidae) Newly Recorded from Japan, Infecting the Introduced Mosquitofish <i>Gambusia affinis</i> . <i>Species Diversity</i> , 2014, 19, 173-178.	0.4	1
53	<i>Choniomyzon inflatus</i> n. sp. (Crustacea: Copepoda: Nicothoidae) associated with <i>Ibacus novemdentatus</i> (Crustacea: Decapoda: Scyllaridae) from Japanese waters. <i>Systematic Parasitology</i> , 2013, 84, 157-165.	1.1	6
54	Double-netting: an alternative approach for the recovery of parasitic copepods from finfishes. <i>Journal of Natural History</i> , 2013, 47, 529-541.	0.5	13

#	ARTICLE	IF	CITATIONS
55	Estimation of sperm storage duration in captive loggerhead turtles (<i>Caretta caretta</i>). <i>Journal of Experimental Marine Biology and Ecology</i> , 2013, 439, 136-142.	1.5	9
56	Growth and reproductive cycle of <i>Hypermastus tokunagai</i> (Caenogastropoda: Eulimidae), an ectoparasite of the sand dollar <i>Scaphechinus mirabilis</i> (Clypeasteroidea: Scutellidae) in the Seto Inland Sea, Japan. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2013, 93, 1041-1051.	0.8	10
57	Redescription of <i>Lepeophtheirus acutus</i> Heegaard, 1943 (Copepoda: Caligidae) parasitic on two elasmobranch hosts off Okinawa-jima Island, Japan. <i>Journal of Natural History</i> , 2013, 47, 581-596.	0.5	3
58	Redescription of <i>Pseudacanthocanthopsis secunda</i> (Yamaguti and Yamasu, 1960) (Copepoda: Tj ETQq0 0 0 rgBT /Overlock 10 Tf off Japan and Korea. <i>Journal of Natural History</i> , 2013, 47, 553-562.	0.5	3
59	Spatial distribution of <i>Neoergasilus japonicus</i> (Copepoda: Ergasilidae) on the fins of bluegill (<i>Lepomis macrochirus</i>). <i>Journal of Natural History</i> , 2013, 47, 543-552.	0.5	4
60	Four new species of <i>Colobomatus</i> (Copepoda: Philichthyidae) parasitic in the lateral line system of marine finfishes captured off the Ryukyu Islands, Japan, with redescrptions of <i>Colobomatus collettei</i> Cressey, 1977 and <i>Colobomatus pupa</i> Izawa, 1974. <i>Journal of Natural History</i> , 2013, 47, 563-580.	0.5	9
61	Contribution of Ju-shey Ho to the systematics of symbiotic copepods of Japan. <i>Journal of Natural History</i> , 2013, 47, 517-527.	0.5	0
62	Benthic shrimp assemblages associated with water masses in the East China Sea and the Yellow Sea. <i>Nippon Suisan Gakkaishi</i> , 2013, 79, 987-1001.	0.1	1
63	First Japanese Record of <i>Heteropriapulus heterotylus</i> (Monogenea: Dactylogyridae), from the Alien Catfish <i>Pterygoplichthys disjunctivus</i> (Siluriformes: Loricariidae) in Okinawa. <i>Species Diversity</i> , 2013, 18, 281-284.	0.4	3
64	The genus <i>Hatschekia</i> (Copepoda: Hatschekiidae) from pufferfishes (Tetraodontiformes: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 387 Td (redescription of <i>H. pholas</i> . <i>Folia Parasitologica</i> , 2013, 60, 61-74.	1.3	4
65	Four new species of splanchnotrophid copepods (Poecilostomatoida) parasitic on doridacean nudibranchs (Gastropoda, Opisthobranchia) from Japan, with proposition of one new genus. <i>ZooKeys</i> , 2012, 247, 1-29.	1.1	10
66	The spotfin burrfish (<i>Chilomycterus reticulatus</i>), a new host record for <i>Cymothoa pulchra</i> (Isopoda, Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 387 Td (0.3	2
67	Mate Selection Based on Genetic Relatedness of Loggerhead Turtles in Captivity. <i>Chelonian Conservation and Biology</i> , 2012, 11, 214-222.	0.6	3
68	A new species of parasitic copepod, <i>Sarcotretes umitakae</i> sp. n. (Siphonostomatoida, Pennellidae), on the rattail (Actinopterygii, Macrouridae) from the East China Sea, Japan. <i>ZooKeys</i> , 2012, 246, 1-10.	1.1	8
69	Influence of host habitat on the occurrence of gill monogeneans <i>Pseudodactylogyrus</i> spp. on wild Japanese eels <i>Anguilla japonica</i> . <i>Diseases of Aquatic Organisms</i> , 2012, 100, 43-49.	1.0	4
70	Two new species of the copepod <i>Hatschekia</i> Poche, 1902 (Siphonostomatoida: Hatschekiidae) from angelfishes (Pisces: Perciformes: Pomacanthidae) collected during the KUMEJIMA 2009 Expedition. <i>Zootaxa</i> , 2012, 3367, 49.	0.5	6
71	A new species of <i>Pseudomacrchiron</i> Reddiah, 1969 (Crustacea: Copepoda: Macrochironidae) associated with scyphistomae of the moon jellyfish <i>Aurelia</i> sp. (Cnidaria: Scyphozoa) off Japan. <i>Systematic Parasitology</i> , 2012, 81, 125-134.	1.1	1
72	Redescription of the fish parasite <i>Nerocila japonica</i> Schioedte & Meinert, 1881 (Crustacea: Isopoda: Tj ETQq0 0 0 rgBT /Overlock 10 Tf Parasitology, 2012, 81, 147-157.	1.1	10

#	ARTICLE	IF	CITATIONS
73	Two species of philometrid nematodes (Philometridae) from marine fishes off Japan, including <i>Philometroides branchiostegi</i> sp. n. from <i>Branchiostegus japonicus</i> (Malacanthidae). <i>Folia Parasitologica</i> , 2012, 59, 71-78.	1.3	16
74	On the origin of <i>Anguillicoloides crassus</i> , the invasive nematode of anguillid eels. <i>Aquatic Invasions</i> , 2012, 7, 443-453.	1.6	24
75	Sperm Utilization Patterns and Reproductive Success in Captive Loggerhead Turtles (<i>Caretta</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 22	0.6	11
76	Chalimus Stages of <i>Caligus latigenitalis</i> (Copepoda: Caligidae) Parasitic on Blackhead Seabream from Japanese Waters, with Discussion of Terminology Used for Developmental Stages of Caligids. <i>Journal of Parasitology</i> , 2011, 97, 221-236.	0.7	11
77	Effects of juvenile size at release and early marine growth on adult return rates for Hokkaido chum salmon (<i>Oncorhynchus keta</i>) in relation to sea surface temperature. <i>Fisheries Oceanography</i> , 2011, 20, 278-293.	1.7	12
78	Host Size- and Habitat-Dependent Intensity of <i>Heliconema longissimum</i> (Nematoda: Physalopteridae) In the Japanese Eel (<i>Anguilla japonica</i>). <i>Journal of Parasitology</i> , 2011, 97, 994-998.	0.7	8
79	<i>Echinorhynchus hexagrammi</i> Baeva, 1965 (Acanthocephala: Echinorhynchidae) from marine fishes off Hokkaido, Japan, with morphological observations and new host records. <i>Systematic Parasitology</i> , 2011, 80, 35-40.	1.1	4
80	Species of <i>Taeniacanthus</i> Sumpf, 1871 (Crustacea: Copepoda: Taeniacanthidae) parasitic on boxfishes (Tetraodontiformes: Aracanidae and Ostraciidae) from the Indo-West Pacific region, with descriptions of two new species. <i>Systematic Parasitology</i> , 2011, 80, 141-157.	1.1	3
81	<i>Caligus Macarovi</i> (Copepoda, Caligidae) from Pacific Bluefin Tuna, <i>Thunnus Orientalis</i> , Cultured in Japan. <i>Crustaceana</i> , 2011, 84, 1145-1147.	0.3	7
82	Description of <i>Naobranchia Denticis</i> N. Sp. (Copepoda, Siphonostomatoida, Lernaepodidae) Parasitic on <i>Dentex Hypselosomus</i> (Teleostei, Sparidae) from Japanese Waters, with a Key to the Species of <i>Naobranchia</i> . <i>Crustaceana</i> , 2011, 84, 461-476.	0.3	2
83	Infection with <i>Argulus japonicus</i> on Amur Catfish <i>Silurus asotus</i> in an Aquarium. <i>Japanese Journal of Limnology</i> , 2011, 72, 211-214.	0.1	4
84	Can research on the early marine life stage of juvenile chum salmon <i>Oncorhynchus keta</i> forecast returns of adult salmon? A case study from eastern Hokkaido, Japan. <i>Fisheries Science</i> , 2010, 76, 909-920.	1.6	10
85	Three new species of <i>Hatschekia</i> Poche, 1902 (Copepoda: Siphonostomatoida: Hatschekiidae) parasitic on boxfishes (Pisces: Tetraodontiformes: Aracanidae and Ostraciidae) in Japanese waters. <i>Systematic Parasitology</i> , 2010, 75, 147-158.	1.1	10
86	A new species of parasitic copepod of the genus <i>Hatschekia</i> Poche, 1902 (Siphonostomatoida:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 22 <i>Systematic Parasitology</i> , 2010, 76, 53-58.	1.1	6
87	<i>Pseudohatschekiidae</i> , a new family for <i>Pseudohatschekia branchiostegi</i> Yamaguti, 1939 (Crustacea:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 22 Japanese waters, with descriptions of the early developmental stages of <i>P. branchiostegi</i> . <i>Systematic Parasitology</i> , 2010, 77, 215-231.	1.1	6
88	Three new species of the family Pennellidae (Copepoda: Siphonostomatoida) from gobiid fishes (Actinopterygii: Perciformes) in coastal waters of the western Pacific Ocean. <i>Zootaxa</i> , 2010, 2687, 29.	0.5	10
89	The copepod genus <i>Hatschekia</i> Poche, 1902 (Siphonostomatoida: Hatschekiidae) from triggerfishes (Pisces: Tetraodontiformes: Balistidae) from off the Ryukyu Islands, Japan, with descriptions of eleven new species. <i>Zootaxa</i> , 2010, 2478, 1.	0.5	11
90	Four New Species of <i>Peniculisa</i> Wilson, 1917 (Copepoda: Siphonostomatoida: Pennellidae) Parasitic on Coastal Marine Fishes in Japanese Waters. <i>Journal of Parasitology</i> , 2010, 96, 689-702.	0.7	4

#	ARTICLE	IF	CITATIONS
91	A New Species of Capillaria (Nematoda: Capillariidae) from the Intestine of the Marine Fish <i>Acanthopagrus schlegelii</i> (Sparidae) from Japan. <i>Journal of Parasitology</i> , 2010, 96, 771-774.	0.7	7
92	Redescription of four species of Hatschekia (Copepoda: Siphonostomatoida: Hatschekiidae) parasitic on tetraodontiform fishes from Japan. <i>Zootaxa</i> , 2009, 2110, 1-21.	0.5	17
93	Three new species of Hatschekia Poche, 1902 (Copepoda: Siphonostomatoida: Hatschekiidae) parasitic on <i>Abalistes filamentosus</i> (Pisces: Tetraodontiformes: Balistidae) from off Okinawa, Japan. <i>Systematic Parasitology</i> , 2009, 74, 225-237.	1.1	10
94	Relationship between zooplankton abundance and the early marine life history of juvenile chum salmon <i>Oncorhynchus keta</i> in eastern Hokkaido, Japan. <i>Fisheries Science</i> , 2009, 75, 303-316.	1.6	21
95	Regional synchrony in return rates of chum salmon (<i>Oncorhynchus keta</i>) in Japan in relation to coastal temperature and size at release. <i>Fisheries Research</i> , 2009, 95, 14-27.	1.7	28
96	Reduction in size-at-age of black sea bream (<i>Acanthopagrus schlegelii</i>) following intensive releases of cultured juveniles in Hiroshima Bay, Japan. <i>Fisheries Research</i> , 2009, 99, 130-133.	1.7	12
97	New Hosts, Including Marine Fishes Cultured in Japan, of <i>Limnotrachelobdella okae</i> (Hirudinida: Tj ETQq1 1 0.784314 rgBT /Overlock 10 0,4 3		
98	Influence of larval size and feeding on the ratio of their settlement to spat collectors presoaked in seawater containing adult Pacific oysters <i>Crassostrea gigas</i> . <i>Nippon Suisan Gakkaishi</i> , 2009, 75, 1036-1041.	0.1	0
99	Stock enhancement programme for black sea bream, <i>Acanthopagrus schlegelii</i> (Bleeker), in Hiroshima Bay, Japan: a review. <i>Aquaculture Research</i> , 2008, 39, 1307-1315.	1.8	56
100	Stock enhancement program for black sea bream (<i>Acanthopagrus schlegelii</i>) in Hiroshima Bay: Monitoring the genetic effects. <i>Aquaculture</i> , 2008, 276, 36-43.	3.5	46
101	Influence of presoaking conditions of spat collectors in seawater containing adult Pacific oysters (<i>Crassostrea gigas</i>) on their larval settlement. <i>Nippon Suisan Gakkaishi</i> , 2008, 74, 1017-1023.	0.1	4
102	Infection with the Parasitic Copepod <i>Clavella parva</i> (Lernaeopodidae) in Gold-eye Rockfish <i>Sebastes thompsoni</i> Broodstock in Japan. <i>Fish Pathology</i> , 2008, 43, 55-60.	0.7	3
103	<i>Sarcotretes Longirostris</i> N. Sp. (Copepoda: Pennellidae) Parasitic on Bluefin Driftfish (<i>Psenes</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 0.8 8 <i>Biology</i> , 2007, 27, 116-120.		
104	A COLLECTION OF FISH LEECHES (HIRUDINIDA: PISCICOLIDAE) FROM JAPAN AND SURROUNDING WATERS, INCLUDING REDESCRIPTIONS OF THREE SPECIES. <i>Journal of Parasitology</i> , 2007, 93, 875-883.	0.7	16
105	Host and Seasonal Distribution of Fish Acanthocephalans from the Lake Biwa Basin, Japan. <i>Comparative Parasitology</i> , 2007, 74, 244-253.	0.4	5
106	Host Body Size, Prevalence, and Breeding Season of the Microparasitic Snail <i>Pelseneria castanea</i> on the Sea Urchin <i>Strongylocentrotus nudus</i> in Coastal Waters of the Sea of Japan off Western Hokkaido, Japan. <i>Japanese Journal of Benthology</i> , 2006, 61, 16-20.	0.1	1
107	Small subunit rDNA and Bayesian inference reveal <i>Pectenophilus formatus</i> (Copepoda: incertae sedis) as highly transformed Mytilicolidae, and support assignment of Chondracanthidae and Xarifiidae to Lichomolgoidea (Cyclopoida). <i>Biological Journal of the Linnean Society</i> , 2006, 87, 403-425.	1.6	74
108	New host and ocean records for <i>Driocephalus cerebrinoxius</i> (Sphyrriidae, Siphonostomatoida) and a reconsideration of phylogeny within Sphyrriidae. <i>Acta Ichthyologica Et Piscatoria</i> , 2006, 36, 1-9.	0.7	6

#	ARTICLE	IF	CITATIONS
109	First record of ostracods as natural intermediate hosts of <i>Anguillicola crassus</i> , a pathogenic swimbladder parasite of eels <i>Anguilla</i> spp.. <i>Diseases of Aquatic Organisms</i> , 2005, 66, 171-173.	1.0	12
110	Copepod Parasites of the Fatheads (Pisces, Psychrolutidae) Their Implication on the Phylogenetic Relationships of Psychrolutid Genera. <i>Zoological Science</i> , 2005, 22, 411-425.	0.7	5
111	Spatial Variability of Living Coccolithophore Distribution in the Western Subarctic Pacific and Western Bering Sea. <i>Journal of Oceanography</i> , 2004, 60, 505-515.	1.7	24
112	The identity of <i>Limnoncaea diuncata</i> Kokubo, 1914 (Copepoda: Poecilostomatoida) from Hokkaido, Japan, with the relegation of <i>Diergasilus</i> Do, 1981 to a junior synonym of <i>Thersitina</i> Norman, 1905. <i>Systematic Parasitology</i> , 2004, 57, 35-44.	1.1	7
113	Ergasilid copepods (Poecilostomatoida) in plankton samples from Hokkaido, Japan, with reconsideration of the taxonomic status of <i>Limnoncaea</i> Kokubo, 1914. <i>Journal of Natural History</i> , 2004, 38, 471-498.	0.5	14
114	An East-West Comparison of the Transition Zone Coastal Pelagic Nekton of the North Pacific Ocean. <i>Journal of Oceanography</i> , 2003, 59, 415-434.	1.7	6
115	Effects of <i>Acanthocephalus</i> sp. (Acanthocephala: Echinorhynchidae) on the body size and reproduction of isopods (<i>Asellus hilgendorfi</i>). <i>Limnology</i> , 2003, 4, 43-46.	1.5	10
116	<i>Paeonocanthus antarcticensis</i> (Hewitt, 1965): a rare copepod parasite of bathypelagic fish, <i>Bathylagus antarcticus</i> Günther, from the Antarctic Ocean. <i>Zoological Science</i> , 2003, 20, 1315-1321.	0.7	6
117	Larval anisakid nematodes from four species of squid (Cephalopoda: Teuthoidea) from the central and western North Pacific Ocean. <i>Journal of Natural History</i> , 2002, 36, 883-891.	0.5	11
118	A mass occurrence of human infection with <i>Diplogonoporus grandis</i> (Cestoda: Diphyllbothriidae) in Shizuoka Prefecture, central Japan. <i>Parasitology International</i> , 2002, 51, 73-79.	1.3	16
119	New Host and Ocean Records for the Parasitic Copepod <i>Bobkabata kabatabobbus</i> (Lernaeosoleidae:). <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10</i>	0.6	9
120	Seasonal and regional occurrence of <i>Acanthocephalus</i> sp. (Acanthocephala: Echinorhynchidae) in fishes and isopods (<i>Asellus hilgendorfi</i>) in a lake system in northern Japan. <i>Limnology</i> , 2002, 3, 143-150.	1.5	6
121	Redescription of <i>Raphidascaaris gigi</i> Fujita, 1928 (Nematoda: Anisakidae), a parasite of freshwater fishes in Japan. <i>Systematic Parasitology</i> , 2002, 52, 193-198.	1.1	15
122	Occurrence of <i>Caligus lalandei</i> Barnard, 1948 (Copepoda, Siphonostomatoida) on Amberjacks (<i>Seriola</i>) <i>Tj ETQq0 0 0 rgBT /Overlock 10</i>	0.7	18
123	Title is missing!. <i>Hydrobiologia</i> , 2001, 453/454, 411-416.	2.0	30
124	Long-term variations in abundance of Pacific herring (<i>Clupea pallasii</i>) in Hokkaido and Sakhalin related to changes in environmental conditions. <i>Progress in Oceanography</i> , 2001, 49, 551-564.	3.2	27
125	WHY INFESTATION BY <i>LEPEOPHTHEIRUS SALMONIS</i> (COPEPODA:CALIGIDAE) IS NOT A PROBLEM IN THE COHO SALMON FARMING INDUSTRY IN JAPAN. <i>Journal of Crustacean Biology</i> , 2001, 21, 954-960.	0.8	9
126	Caryophyllidean tapeworms (Platyhelminthes: Eucestoda) from freshwater fishes in Japan. <i>Folia Parasitologica</i> , 2001, 48, 275-288.	1.3	27

#	ARTICLE	IF	CITATIONS
127	Interannual variability in stock abundance of the neon flying squid, <i>Ommastrephes bartramii</i> , in the North Pacific Ocean during 1979-1998: impact of driftnet fishing and oceanographic conditions. <i>Fisheries Oceanography</i> , 2000, 9, 163-170.	1.7	41
128	Some anisakid nematodes from marine fishes of Japan and the North Pacific Ocean. <i>Journal of Natural History</i> , 2000, 34, 1555-1574.	0.5	28
129	Two remarkable nematodes from sharks in Japan. <i>Journal of Natural History</i> , 2000, 34, 1-13.	0.5	13
130	Age, Growth, and Feeding Habits of Lanternfish, <i>Stenobrachius leucopsarus</i> (Myctophidae), Collected from the Near-surface Layer in the Bering Sea. <i>Fisheries Science</i> , 1999, 65, 11-15.	1.6	10
131	New data on the morphology of <i>Philonema oncorhynchi</i> Kuitunen-Ekbaum, 1933 (Nematoda: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 503-26). <i>Parasitology</i> , 1999, 43, 67-74.	1.1	11
132	Epipelagic nekton of the North Pacific Subarctic and Transition Zones. <i>Progress in Oceanography</i> , 1999, 43, 365-397.	3.2	41
133	Observations on five species of philometrid nematodes from marine fishes in Japan. <i>Systematic Parasitology</i> , 1998, 40, 67-80.	1.1	32
134	Occurrence and effects of the parasitic copepod <i>Salmincola carpionis</i> on salmonids in the Nikko District, central Japan. <i>Journal of Marine Systems</i> , 1998, 15, 269-272.	2.1	9
135	Distribution of the rough pomfret, <i>Taractes asper</i> (Bramidae), in the central North Pacific Ocean. <i>Ichthyological Research</i> , 1998, 45, 419-423.	0.8	0
136	Thermal limits and ocean migrations of sockeye salmon (<i>Oncorhynchus nerka</i>): long-term consequences of global warming. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 1998, 55, 937-948.	1.4	123
137	Two New Huffmanella Species, <i>H. japonica</i> n. sp. and <i>H. shikokuensis</i> n. sp. (Nematoda: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 503-26).	0.7	26
138	<i>Trebius shiinoi</i> n. sp. (Trebiidae: Siphonostomatoida: Copepoda) from Uteri and Embryos of the Japanese Angelshark (<i>Squatina japonica</i>) and the Clouded Angelshark (<i>Squatina nebulosa</i>), and Redescription of <i>Trebius longicaudatus</i> . <i>Journal of Parasitology</i> , 1998, 84, 1218.	0.7	5
139	Role of <i>Leptodora kindti</i> (Cladocera:Leptodoridae) in the life cycle of <i>Raphidascaris biwakoensis</i> (Nematoda:Anisakidae), a fish parasite in Lake Biwa, Japan. <i>Diseases of Aquatic Organisms</i> , 1998, 32, 157-160.	1.0	4
140	Trophic relations in the subarctic North Pacific ecosystem:possible feeding effect from pink salmon. <i>Marine Ecology - Progress Series</i> , 1997, 150, 75-85.	1.9	99
141	Possible Life Cycle of <i>Longicollum pagrosomi</i> , an Acanthocephalan Parasite of Cultured Red Sea Bream.. <i>Fish Pathology</i> , 1996, 31, 235-236.	0.7	8
142	Distribution of Pelagic Elasmobranchs Caught by Salmon Research Gillnets in the North Pacific. <i>Fisheries Science</i> , 1996, 62, 860-865.	1.6	52
143	Development of Ultrasonic Telemetry Technique for Investigating the Magnetic of Salmonids. <i>Fisheries Science</i> , 1996, 62, 698-704.	1.6	15
144	Larval Anisakid Nematodes of Japanese Common Squid (<i>Todarodes pacificus</i>) from the Sea of Japan. <i>Journal of Parasitology</i> , 1995, 81, 69.	0.7	25

#	ARTICLE	IF	CITATIONS
145	Rediscovery in Japan and host association of <i>Salmincola carpionis</i> (Copepoda: Lernaepodidae), a parasite of wild and reared freshwater salmonids. Canadian Journal of Fisheries and Aquatic Sciences, 1995, 52, 178-185.	1.4	13
146	Recent Changes in Age and Size of Chum Salmon (<i>Oncorhynchus keta</i>) in the North Pacific Ocean and Possible Causes. Canadian Journal of Fisheries and Aquatic Sciences, 1993, 50, 290-295.	1.4	67
147	Distribution and Bivalve Hosts of the Parasitic Copepod <i>Pectenophilus Ornatus</i> Nagasawa, Bresciani, and L��tzen in Japan. Journal of Crustacean Biology, 1993, 13, 544-550.	0.8	2
148	Host Utilization by the Salmon Louse <i>Lepeophtheirus salmonis</i> (Copepoda: Caligidae) in the Sea of Japan. Journal of Parasitology, 1993, 79, 127.	0.7	13
149	<i>Kudoa intestinalis</i> n. sp. (Myxosporea: Multivalvulida) from the Intestinal Musculature of the Striped Mullet, <i>Mugil cephalus</i> , from Japan. Journal of Parasitology, 1993, 79, 190.	0.7	25
150	Infection of the Salmon Louse <i>Lepeophtheirus salmonis</i> (Copepoda: Caligidae) on Seawater-Farmed Salmonids in Northern Honshu, Japan.. Fish Pathology, 1993, 28, 143-144.	0.7	5
151	Effects of <i>Pectenophilus ornatus</i> (Copepoda) on the Biomass of Cultured Japanese Scallop <i>Patinopecten yessoensis</i> . Journal of Parasitology, 1992, 78, 552.	0.7	8
152	Long-term changes in the population size and geographical distribution of <i>Pennella</i> sp. (Copepoda) on the saury, <i>Cololabis saira</i> , in the western North Pacific Ocean and adjacent seas. Hydrobiologia, 1988, 167-168, 571-577.	2.0	11
153	Morphology of <i>Pectenophilus ornatus</i> , New Genus, New Species, a Copepod Parasite of the Japanese Scallop <i>Patinopecten yessoensis</i> . Journal of Crustacean Biology, 1988, 8, 31.	0.8	14
154	Long-term changes in the population size and geographical distribution of <i>Pennella</i> sp. (Copepoda) on the saury, <i>Cololabis saira</i> , in the western North Pacific Ocean and adjacent seas*. , 1988, , 571-577.		2
155	Prevalence and abundance of <i>Lepeophtheirus salmonis</i> (Copepoda: Caligidae) on high-seas salmon and trout in the North Pacific Ocean.. Nippon Suisan Gakkaishi, 1987, 53, 2151-2156.	0.1	27
156	Occurrence and effects of <i>Haemobaphes diceraus</i> (Copepoda:Pennellidae) on brown sole <i>Limanda herzensteini</i> off the Okhotsk coast of Hokkaido.. Nippon Suisan Gakkaishi, 1987, 53, 991-994.	0.1	9
157	Prevalence of visceral adhesions in sockeye salmon, <i>Oncorhynchus nerka</i> , in the central North Pacific Ocean.. Fish Pathology, 1985, 20, 313-321.	0.7	9
158	The finding of <i>Pennella</i> sp. (Copepoda: Pennellidae) on the saury, <i>Cololabis saira</i> , in the western and central North Pacific Ocean and the Okhotsk Sea.. Fish Pathology, 1984, 18, 205-208.	0.7	9
159	Crustacean Parasites of the Saury, <i>Cololabis saira</i> : a Review. Fish Pathology, 1984, 19, 57-63.	0.7	5
160	Ecological factors influencing the infection levels of salmonids by <i>Acanthocephalus opsariichthydis</i> (Acanthocephala: Echinorhynchidae) in Lake Yunoko, Japan.. Fish Pathology, 1983, 18, 53-60.	0.7	6
161	<i>Acanthocephalus lucidus</i> van CLEAVE (Acanthocephala: Echinorhynchidae) from cultured rainbow trout <i>Salmo gairdneri</i> Richardson.. Nippon Suisan Gakkaishi, 1981, 47, 1153-1156.	0.1	3