

# Wenbing Sun

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5502751/publications.pdf>

Version: 2024-02-01

13  
papers

255  
citations

1464605

7  
h-index

1255698

13  
g-index

13  
all docs

13  
docs citations

13  
times ranked

385  
citing authors

#	ARTICLE	IF	CITATIONS
1	Arsenic trioxide inhibits angiogenesis of hepatocellular carcinoma after insufficient radiofrequency ablation via blocking paracrine angiopoietin-1 and angiopoietin-2. <i>International Journal of Hyperthermia</i> , 2022, 39, 888-896.	1.1	6
2	<p></p>miR-133a-3p Regulates Hepatocellular Carcinoma Progression Through Targeting CORO1C</p>. <i>Cancer Management and Research</i> , 2020, Volume 12, 8685-8693.	0.9	18
3	Platelet lysates in Hepatocellular Carcinoma patients after radiofrequency ablation facilitate tumor proliferation, invasion and vasculogenic mimicry. <i>International Journal of Medical Sciences</i> , 2020, 17, 2104-2112.	1.1	7
4	Downregulation of serum exosomal miRâ€³20d predicts poor prognosis in hepatocellular carcinoma. <i>Journal of Clinical Laboratory Analysis</i> , 2020, 34, e23239.	0.9	42
5	ATPase Inhibitory Factor 1 Promotes Hepatocellular Carcinoma Progression After Insufficient Radiofrequency Ablation, and Attenuates Cell Sensitivity to Sorafenib Therapy. <i>Frontiers in Oncology</i> , 2020, 10, 1080.	1.3	21
6	FO XK2 downregulation suppresses EMT in hepatocellular carcinoma. <i>Open Medicine (Poland)</i> , 2020, 15, 702-708.	0.6	5
7	Endothelial pyroptosis underlies systemic inflammatory response following radiofrequency ablation of hepatic hemangiomas. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2019, 79, 619-628.	0.6	11
8	HMGB1-induced endothelial cell pyroptosis is involved in systemic inflammatory response syndrome following radiofrequency ablation of hepatic hemangiomas. <i>American Journal of Translational Research (discontinued)</i> , 2019, 11, 7555-7567.	0.0	4
9	ATPase inhibitory factor 1 inhibition improves the antitumor of YCâ€™1 against hepatocellular carcinoma. <i>Oncology Letters</i> , 2018, 16, 5230-5236.	0.8	4
10	Metformin exhibits the anti-proliferation and anti-invasion effects in hepatocellular carcinoma cells after insufficient radiofrequency ablation. <i>Cancer Cell International</i> , 2017, 17, 48.	1.8	19
11	Sorafenib suppresses the epithelial-mesenchymal transition of hepatocellular carcinoma cells after insufficient radiofrequency ablation. <i>BMC Cancer</i> , 2015, 15, 939.	1.1	37
12	Positive feedback between oncogenic KRAS and&nbsp;HIF-1&alpha; confers drug resistance in colorectal cancer. <i>OncoTargets and Therapy</i> , 2015, 8, 1229.	1.0	15
13	Insufficient radiofrequency ablation promotes epithelial-mesenchymal transition of hepatocellular carcinoma cells through Akt and ERK signaling pathways. <i>Journal of Translational Medicine</i> , 2013, 11, 273.	1.8	66