

Kelig Mahe

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5509250/kelig-mahe-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

43
papers

617
citations

13
h-index

24
g-index

49
ext. papers

746
ext. citations

2.6
avg, IF

3.59
L-index

#	Paper	IF	Citations
43	Fluctuating Asymmetry in Asteriscii Otoliths of Common Carp (<i>Cyprinus carpio</i>) Collected from Three Localities in Iraqi Rivers Linked to Environmental Factors. <i>Fishes</i> , 2022 , 7, 91	2.5	0
42	Directional Bilateral Asymmetry in Fish Otolith: A Potential Tool to Evaluate Stock Boundaries?. <i>Symmetry</i> , 2021 , 13, 987	2.7	3
41	Reconstructing freshwater fishing seasonality in a neotropical savanna: First application of swamp eel (<i>Synbranchus marmoratus</i>) sclerochronology to a pre-Columbian Amazonian site (Loma Salvatierra, Bolivia). <i>Journal of Archaeological Science: Reports</i> , 2021 , 37, 102880	0.7	1
40	New scale analyses reveal centenarian African coelacanths. <i>Current Biology</i> , 2021 , 31, 3621-3628.e4	6.3	3
39	Relationship between Body and Otolith Morphological Characteristics of Sabre Squirrelfish (<i>Sargocentron spiniferum</i>) from the Southern Red Sea: Difference between Right and Left Otoliths. <i>Oceans</i> , 2021 , 2, 624-633	1.3	2
38	Development in the European flounder (<i>Platichthys flesus</i>) of a q-PCR assay for the measurement of telomere length, a potential biomarker of pollutant effects for biomonitoring studies. <i>Marine Pollution Bulletin</i> , 2021 , 170, 112610	6.7	
37	Age, growth and mortality of the striped seabream <i>Lithognathus mormyrus</i> (Linnaeus, 1758) in the Gulf of Tunis (Central Mediterranean Sea). <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2021 , 101, 159-167	1.1	
36	Characterizing seasonal fishing patterns and growth dynamics during the Middle and Late Holocene in the Strait of Magellan (Chilean Patagonia): Sclerochronological analysis of tadpole codling (<i>Salilota australis</i>) vertebrae. <i>Journal of Island and Coastal Archaeology</i> , 2020 , 1-20	1.2	2
35	Complementarity and discriminatory power of genotype and otolith shape in describing the fine-scale population structure of an exploited fish, the common sole of the Eastern English Channel. <i>PLoS ONE</i> , 2020 , 15, e0241429	3.7	6
34	Size-effect, asymmetry, and small-scale spatial variation in otolith shape of juvenile sole in the Southern North Sea. <i>Hydrobiologia</i> , 2019 , 845, 95-108	2.4	4
33	Use of fish otoliths as a temporal biomarker of field uranium exposure. <i>Science of the Total Environment</i> , 2019 , 690, 511-521	10.2	7
32	Do environmental conditions (temperature and food composition) affect otolith shape during fish early-juvenile phase? An experimental approach applied to European Seabass (<i>Dicentrarchus labrax</i>). <i>Journal of Experimental Marine Biology and Ecology</i> , 2019 , 521, 151239	2.1	7
31	Explorative analysis on red mullet (<i>Mullus barbatus</i>) ageing data variability in the Mediterranean. <i>Scientia Marina</i> , 2019 , 83, 271	1.8	3
30	Directional bilateral asymmetry in otolith morphology may affect fish stock discrimination based on otolith shape analysis. <i>ICES Journal of Marine Science</i> , 2019 , 76, 232-243	2.7	13
29	Evidence of a relationship between weight and total length of marine fish in the North-eastern Atlantic Ocean: physiological, spatial and temporal variations. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2018 , 98, 617-625	1.1	7
28	Age and growth of the striped seabream, <i>Lithognathus mormyrus</i> (Actinopterygii: Perciformes: Sparidae), in the central coast of Algeria, Mediterranean Sea. <i>Acta Ichthyologica Et Piscatoria</i> , 2018 , 48, 319-328	1.8	3
27	Age, growth and mortality of <i>Sciaena umbra</i> (Sciaenidae) in the Gulf of Tunis. <i>Scientia Marina</i> , 2018 , 82, 17	1.8	4

26	Utility of mixed effects models to inform the stock structure of whiting in the Northeast Atlantic Ocean. <i>Fisheries Research</i> , 2017 , 190, 132-139	2.3	4
25	Estimation of Age, Growth and Fishing Season of a Palaeolithic Population of Grayling (<i>Thymallus thymallus</i>) Using Scale Analysis. <i>International Journal of Osteoarchaeology</i> , 2017 , 27, 683-692	1.1	4
24	Relationship between somatic growth and otolith growth: a case study of the ornate jobfish <i>Pristipomoides argyrogrammicus</i> from the coast of Réunion (SW Indian Ocean). <i>African Journal of Marine Science</i> , 2017 , 39, 145-151	0.8	3
23	Use of otolith-shape analysis for stock discrimination of Boops boops along the Algerian coast (southwestern Mediterranean Sea). <i>African Journal of Marine Science</i> , 2017 , 39, 251-258	0.8	11
22	Ontogenetic changes in the larval condition of Downs herring: use of a multi-index approach at an individual scale. <i>Marine Biology</i> , 2017 , 164, 1	2.5	3
21	Lost in the North: The first record of <i>Diretmichthys parini</i> (Post and QuEo, 1981) in the northern North Sea. <i>Marine Pollution Bulletin</i> , 2017 , 115, 439-443	6.7	1
20	Comparing growth in surface and cave morphs of the species : insights from scales. <i>EvoDevo</i> , 2017 , 8, 23	3.2	15
19	Otolith shape as a valuable tool to evaluate the stock structure of swordfish <i>Xiphias gladius</i> in the Indian Ocean. <i>African Journal of Marine Science</i> , 2016 , 38, 457-464	0.8	6
18	Growth of an Inshore Antarctic fish, <i>Trematomus newnesi</i> (Nototheniidae), off Adelie Land. <i>Polar Science</i> , 2016 , 10, 167-172	2.3	2
17	Identifying blue whiting (<i>Micromesistius poutassou</i>) stock structure in the Northeast Atlantic by otolith shape analysis. <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 2016 , 73, 1363-1371	2.4	18
16	Diet is correlated with otolith shape in marine fish. <i>Marine Ecology - Progress Series</i> , 2016 , 555, 167-184	2.6	25
15	Variability of PCB burden in 5 fish and sharks species of the French Mediterranean continental slope. <i>Environmental Pollution</i> , 2016 , 212, 374-381	9.3	7
14	Sagittal otolith morphogenesis asymmetry in marine fishes. <i>Journal of Fish Biology</i> , 2015 , 87, 646-63	1.9	29
13	Habitat discrimination of big-scale sand smelt <i>Atherina boyeri</i> Risso, 1810 (Atheriniformes: Atherinidae) in eastern Algeria using somatic morphology and otolith shape. <i>Italian Journal of Zoology</i> , 2015 , 82, 446-453		6
12	Are red mullet efficient as bio-indicators of mercury contamination? A case study from the French Mediterranean. <i>Marine Pollution Bulletin</i> , 2015 , 91, 191-9	6.7	22
11	Otolith growth and age estimation of bastard grunt, <i>Pomadasyus incisus</i> (Actinopterygii: Perciformes: Haemulidae), in the Gulf of Tunis (Central Mediterranean). <i>Acta Ichthyologica Et Piscatoria</i> , 2015 , 45, 57-63	1.8	2
10	Morphological variability of the shape of striped red mullet <i>Mullus surmuletus</i> in relation to stock discrimination between the Bay of Biscay and the eastern English Channel. <i>Journal of Fish Biology</i> , 2014 , 84, 1063-73	1.9	10
9	Black scorpionfish <i>Scorpaena porcus</i> (Scorpaenidae): a first record in the eastern English Channel. <i>Marine Biodiversity Records</i> , 2014 , 7,	2	2

8	Striped red mullet (<i>Mullus surmuletus</i> , Linnaeus, 1758) in the eastern English Channel and southern North Sea: growth and reproductive biology. <i>Journal of Applied Ichthyology</i> , 2013 , 29, 1067-1072	0.9	9
7	Discriminating stocks of striped red mullet (<i>Mullus surmuletus</i>) in the Northwest European seas using three automatic shape classification methods. <i>Fisheries Research</i> , 2013 , 143, 153-160	2.3	20
6	Shell Shape Analysis and Spatial Allometry Patterns of Manila Clam (<i>Ruditapes philippinarum</i>) in a Mesotidal Coastal Lagoon. <i>Journal of Marine Biology</i> , 2012 , 2012, 1-11	1	21
5	Calcein and manganese experiments for marking the shell of the common cockle (<i>Cerastoderma edule</i>): tidal rhythm validation of increments formation. <i>Aquatic Living Resources</i> , 2010 , 23, 239-245	1.5	22
4	Ecosystem trends: evidence for agreement between fishers' perceptions and scientific information. <i>ICES Journal of Marine Science</i> , 2008 , 65, 1057-1068	2.7	61
3	Ontogenetic and spatial variation in the diet of hake (<i>Merluccius merluccius</i>) in the Bay of Biscay and the Celtic Sea. <i>ICES Journal of Marine Science</i> , 2007 , 64, 1210-1219	2.7	54
2	Growth, feeding and distribution of the solenette <i>Buglossidium luteum</i> with particular reference to its habitat preference. <i>Journal of Sea Research</i> , 2004 , 51, 211-217	1.9	18
1	Effects of temperature on growth and metabolism in a Mediterranean population of European sea bass, <i>Dicentrarchus labrax</i> . <i>Aquaculture</i> , 2004 , 237, 269-280	4.4	176