

Lisa Connolly

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5515024/lisa-connolly-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

76
papers

1,496
citations

23
h-index

35
g-index

82
ext. papers

1,770
ext. citations

5.5
avg. IF

4.6
L-index

#	Paper	IF	Citations
76	Microplastics and Their Impact on Reproduction-Can we Learn From the Model?. <i>Frontiers in Toxicology</i> , 2022 , 4, 748912	1.6	3
75	Obesity II: Establishing Causal Links Between Chemical Exposures and Obesity.. <i>Biochemical Pharmacology</i> , 2022 , 115015	6	6
74	A human relevant mixture of persistent organic pollutants (POPs) and perfluorooctane sulfonic acid (PFOS) enhance nerve growth factor (NGF)-induced neurite outgrowth in PC12 cells. <i>Toxicology Letters</i> , 2021 , 338, 85-96	4.4	3
73	Effects of Defined Mixtures of Persistent Organic Pollutants (POPs) on Pre-lethal Cytotoxicity in the Human A-498 Kidney Cell Line In Vitro. <i>Exposure and Health</i> , 2021 , 13, 465-475	8.8	1
72	Hormonal activity in commonly used Black hair care products: evaluating hormone disruption as a plausible contribution to health disparities. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2021 , 31, 476-486	6.7	6
71	The insulin-like growth factor system: A target for endocrine disruptors?. <i>Environment International</i> , 2021 , 147, 106311	12.9	8
70	Investigating the pre-lethal cytotoxic effects of bis(2,4-di-tert-butylphenyl)phosphate on Chinese hamster ovary cells using high content analysis. <i>Journal of Biotechnology</i> , 2021 , 328, 59-71	3.7	
69	Assessing the chemical-induced estrogenicity using in silico and in vitro methods. <i>Environmental Toxicology and Pharmacology</i> , 2021 , 87, 103688	5.8	1
68	In vitro differential responses of rat and human aryl hydrocarbon receptor to two distinct ligands and to different polyphenols. <i>Environmental Pollution</i> , 2020 , 265, 114966	9.3	4
67	Safeguarding Female Reproductive Health against Endocrine Disrupting Chemicals-The FREIA Project. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	8
66	Low Doses of Mycotoxin Mixtures below EU Regulatory Limits Can Negatively Affect the Performance of Broiler Chickens: A Longitudinal Study. <i>Toxins</i> , 2020 , 12,	4.9	12
65	Putative adverse outcome pathways for female reproductive disorders to improve testing and regulation of chemicals. <i>Archives of Toxicology</i> , 2020 , 94, 3359-3379	5.8	15
64	Public Awareness and Risk Perceptions of Endocrine Disrupting Chemicals: A Qualitative Study. <i>International Journal of Environmental Research and Public Health</i> , 2020 , 17,	4.6	3
63	Potential adverse effects on animal health and performance caused by the addition of mineral adsorbents to feeds to reduce mycotoxin exposure. <i>Mycotoxin Research</i> , 2020 , 36, 115-126	4	26
62	Human-Based Exposure Levels of Perfluoroalkyl Acids May Induce Harmful Effects to Health by Disrupting Major Components of Androgen Receptor Signalling In Vitro. <i>Exposure and Health</i> , 2020 , 12, 527-538	8.8	3
61	Investigation of In Vitro Endocrine Activities of and Cyanobacterial Strains. <i>Toxins</i> , 2020 , 12,	4.9	6
60	A Human Relevant Defined Mixture of Persistent Organic Pollutants (POPs) Affects In Vitro Secretion of Glucagon-Like Peptide 1 (GLP-1), but Does Not Affect Translocation of Its Receptor. <i>Toxicological Sciences</i> , 2019 , 172, 359-367	4.4	3

59	Human blood-based exposure levels of persistent organic pollutant (POP) mixtures antagonise androgen receptor transactivation and translocation. <i>Environment International</i> , 2019 , 132, 105083	12.9	9
58	A mixture of persistent organic pollutants relevant for human exposure inhibits the transactivation activity of the aryl hydrocarbon receptor in vitro. <i>Environmental Pollution</i> , 2019 , 254, 113098	9.3	14
57	The effect of individual and mixtures of mycotoxins and persistent organochloride pesticides on oestrogen receptor transcriptional activation using in vitro reporter gene assays. <i>Food and Chemical Toxicology</i> , 2019 , 130, 68-78	4.7	6
56	Improvements in single-use bioreactor film material composition leads to robust and reliable Chinese hamster ovary cell performance. <i>Biotechnology Progress</i> , 2019 , 35, e2824	2.8	5
55	In vitro bioassay investigations of suspected obesogen monosodium glutamate at the level of nuclear receptor binding and steroidogenesis. <i>Toxicology Letters</i> , 2019 , 301, 11-16	4.4	9
54	Comparative In Vitro Assessment of a Range of Commercial Feed Additives with Multiple Mycotoxin Binding Claims. <i>Toxins</i> , 2019 , 11,	4.9	15
53	The origin of in-vitro estrogen-like activity in oregano herb extracts. <i>Toxicology in Vitro</i> , 2019 , 56, 101-109	9.6	6
52	Mycotoxins as potential cause of human infertility – a review of evidence from animal and cellular models. <i>Acta Horticulturae</i> , 2018 , 513-525	0.3	
51	Response to comments raised by Fernstrom and Smiga (TOXLET-D-17-00079) on our recent article Shannon M et al. [Toxicol. Lett. 265 (2017) 97]. <i>Toxicology Letters</i> , 2017 , 272, 103-105	4.4	
50	The endocrine disrupting potential of monosodium glutamate (MSG) on secretion of the glucagon-like peptide-1 (GLP-1) gut hormone and GLP-1 receptor interaction. <i>Toxicology Letters</i> , 2017 , 265, 97-105	4.4	24
49	Effects of defined mixtures of persistent organic pollutants (POPs) on multiple cellular responses in the human hepatocarcinoma cell line, HepG2, using high content analysis screening. <i>Toxicology and Applied Pharmacology</i> , 2016 , 294, 21-31	4.6	31
48	Label-free based quantitative proteomics analysis of primary neonatal porcine Leydig cells exposed to the persistent contaminant 3-methylsulfonyl-DDE. <i>Journal of Proteomics</i> , 2016 , 137, 68-82	3.9	7
47	In vitro bioassay investigations of the endocrine disrupting potential of steviol glycosides and their metabolite steviol, components of the natural sweetener Stevia. <i>Molecular and Cellular Endocrinology</i> , 2016 , 427, 65-72	4.4	15
46	Do persistent organic pollutants interact with the stress response? Individual compounds, and their mixtures, interaction with the glucocorticoid receptor. <i>Toxicology Letters</i> , 2016 , 241, 121-32	4.4	18
45	An in vitro investigation on the cytotoxic and nuclear receptor transcriptional activity of the mycotoxins fumonisin B1 and beauvericin. <i>Toxicology Letters</i> , 2016 , 257, 1-10	4.4	25
44	Challenging conventional risk assessment with respect to human exposure to multiple food contaminants in food: A case study using maize. <i>Toxicology Letters</i> , 2015 , 238, 54-64	4.4	24
43	Endocrine disruptor activity of multiple environmental food chain contaminants. <i>Toxicology in Vitro</i> , 2015 , 29, 211-20	3.6	31
42	Validation of an ultra high performance liquid chromatography tandem mass spectrometry method for detection and quantitation of 19 endocrine disruptors in milk. <i>Food Control</i> , 2015 , 48, 48-55	6.2	24

41	Biotransformation of zearalenone and zearalenols to their major glucuronide metabolites reduces estrogenic activity. <i>Toxicology in Vitro</i> , 2015 , 29, 575-81	3.6	50
40	High content analysis: a sensitive tool to detect and quantify the cytotoxic, synergistic and antagonistic effects of chemical contaminants in foods. <i>Toxicology Letters</i> , 2015 , 233, 278-86	4.4	22
39	An investigation of the endocrine disrupting potential of enniatin B using in vitro bioassays. <i>Toxicology Letters</i> , 2015 , 233, 84-94	4.4	14
38	Estrogenic endocrine disruptors present in sports supplements. A risk assessment for human health. <i>Food Chemistry</i> , 2014 , 159, 157-65	8.5	19
37	Effects of the mycotoxin patulin at the level of nuclear receptor transcriptional activity and steroidogenesis in vitro. <i>Toxicology Letters</i> , 2014 , 229, 366-73	4.4	15
36	Cytotoxic assessment of the regulated, co-existing mycotoxins aflatoxin B1, fumonisin B1 and ochratoxin, in single, binary and tertiary mixtures. <i>Toxicon</i> , 2014 , 90, 70-81	2.8	38
35	Validation and application of a reporter gene assay for the determination of estrogenic endocrine disruptor activity in milk. <i>Food and Chemical Toxicology</i> , 2014 , 69, 260-6	4.7	13
34	Using SILAC proteomics to investigate the effect of the mycotoxin, alternariol, in the human H295R steroidogenesis model. <i>Cell Biology and Toxicology</i> , 2014 , 30, 361-76	7.4	6
33	Endocrine disruptor activity in bottled mineral and flavoured water. <i>Food Chemistry</i> , 2013 , 136, 1590-6	8.5	35
32	An in vitro investigation of endocrine disrupting effects of the mycotoxin alternariol. <i>Toxicology and Applied Pharmacology</i> , 2013 , 271, 64-71	4.6	49
31	Endocrine disrupting effects of ochratoxin A at the level of nuclear receptor activation and steroidogenesis. <i>Toxicology Letters</i> , 2013 , 217, 243-50	4.4	30
30	Removal of natural hormones in dairy farm wastewater using reactive and sorptive materials. <i>Science of the Total Environment</i> , 2013 , 461-462, 1-9	10.2	13
29	Cytosol protein regulation in H295R steroidogenesis model induced by the zearalenone metabolites, β and β zearalenol. <i>Toxicon</i> , 2012 , 59, 17-24	2.8	13
28	Treatment of estrogens and androgens in dairy wastewater by a constructed wetland system. <i>Water Research</i> , 2012 , 46, 2333-43	12.5	37
27	An in vitro investigation of endocrine disrupting effects of trichothecenes deoxynivalenol (DON), T-2 and HT-2 toxins. <i>Toxicology Letters</i> , 2012 , 214, 268-78	4.4	43
26	Validation and application of reporter gene assays for the determination of estrogenic and androgenic endocrine disruptor activity in sport supplements. <i>Analytical and Bioanalytical Chemistry</i> , 2012 , 403, 3057-67	4.4	10
25	Production of a monoclonal antibody and its application in an optical biosensor based assay for the quantitative measurement of pantothenic acid (vitamin B5) in foodstuffs. <i>Food Chemistry</i> , 2012 , 134, 540-545	8.5	10
24	Relative quantification of the proteomic changes associated with the mycotoxin zearalenone in the H295R steroidogenesis model. <i>Toxicon</i> , 2011 , 58, 533-42	2.8	13

23	Endocrine disrupting effects of zearalenone, alpha- and beta-zearalenol at the level of nuclear receptor binding and steroidogenesis. <i>Toxicology Letters</i> , 2011 , 206, 210-7	4.4	157
22	The application of reporter gene assays for the detection of endocrine disruptors in sport supplements. <i>Analytica Chimica Acta</i> , 2011 , 700, 34-40	6.6	17
21	Development and validation of a fast monoclonal based disequilibrium enzyme-linked immunosorbent assay for the detection of triphenylmethane dyes and their metabolites in fish. <i>Analytica Chimica Acta</i> , 2011 , 698, 51-60	6.6	27
20	In vitro bioassays for the study of endocrine-disrupting food additives and contaminants. <i>TrAC - Trends in Analytical Chemistry</i> , 2011 , 30, 227-238	14.6	39
19	Immunochemical and mass spectrometric analysis of N ϵ (carboxymethyl)lysine content of AGE-BSA systems prepared with and without selected antiglycation agents. <i>Journal of Agricultural and Food Chemistry</i> , 2010 , 58, 11955-61	5.7	7
18	Effect of inhibitor compounds on N ϵ (carboxymethyl)lysine (CML) and N ϵ (carboxyethyl)lysine (CEL) formation in model foods. <i>Journal of Agricultural and Food Chemistry</i> , 2010 , 58, 12036-41	5.7	67
17	Detection of glucocorticoid bioactivity in bovine urine samples using a reporter gene assay. <i>Analytica Chimica Acta</i> , 2009 , 637, 321-7	6.6	5
16	Expression of multidrug resistance markers ABCB1 (MDR-1/P-gp) and ABCC1 (MRP-1) in renal cell carcinoma. <i>BMC Urology</i> , 2009 , 9, 6	2.2	65
15	Development of a monoclonal antibody binding okadaic acid and dinophysistoxins-1, -2 in proportion to their toxicity equivalence factors. <i>Toxicol</i> , 2009 , 54, 491-8	2.8	52
14	Development and single-laboratory validation of a pseudofunctional biosensor immunoassay for the detection of the okadaic acid group of toxins. <i>Analytical Chemistry</i> , 2009 , 81, 10208-14	7.8	43
13	Endocrine-disrupting chemicals: origins, fates and transmission into the food chain 2009 , 103-125		5
12	The development of a multi-nitroimidazole residue analysis assay by optical biosensor via a proof of concept project to develop and assess a prototype test kit. <i>Analytica Chimica Acta</i> , 2007 , 598, 155-61	6.6	21
11	Rapid screening method for halofuginone residues in poultry eggs and liver using time-resolved fluorometry combined with the all-in-one dry chemistry assay concept. <i>Analytica Chimica Acta</i> , 2005 , 529, 21-25	6.6	16
10	Increased anti-tumour efficacy of doxorubicin when combined with sulindac in a xenograft model of an MRP-1-positive human lung cancer. <i>Anticancer Research</i> , 2004 , 24, 457-64	2.3	22
9	Production and characterisation of polyclonal antibodies to a range of nitroimidazoles. <i>Analytica Chimica Acta</i> , 2003 , 483, 193-200	6.6	16
8	The production and characterisation of an antibody to detect the coccidiostat toltrazuril and its metabolite ponazuril. <i>Analyst</i> , 2003 , 128, 459-61	5	5
7	The production and characterisation of dinitrocarbanilide antibodies raised using antigen mimics. <i>Journal of Immunological Methods</i> , 2002 , 264, 45-51	2.5	20
6	A new monoclonal antibody, P2A8(6), that specifically recognizes a novel epitope on the multidrug resistance-associated protein 1 (MRP1), but not on MRP2 nor MRP3. <i>Hybridoma</i> , 2001 , 20, 333-41		7

5	Selection with melphalan or paclitaxel (Taxol) yields variants with different patterns of multidrug resistance, integrin expression and in vitro invasiveness. <i>European Journal of Cancer</i> , 2001 , 37, 1041-52	7.5	61
4	A new monoclonal antibody that specifically recognises the MDR-3-encoded gene product. <i>International Journal of Cancer</i> , 1999 , 80, 265-71	7.5	6
3	Recent developments in drug resistance and apoptosis research. <i>Critical Reviews in Oncology/Hematology</i> , 1998 , 28, 181-205	7	12
2	Isolation from a human MDR lung cell line of multiple clonal subpopulations which exhibit significantly different drug resistance. <i>International Journal of Cancer</i> , 1997 , 71, 907-15	7.5	24
1	Lipogenic Potency of Individual Perfluorinated Alkyl Acids (PFAAs) and Persistent Organic Pollutant (POP) Mixtures at Human Blood-Based Exposure Levels on Adipogenesis in 3T3-L1 Cells. <i>Exposure and Health</i> ,1	8.8	1