

Gene Lin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5521040/publications.pdf>

Version: 2024-02-01

10
papers

451
citations

1477746

6
h-index

1872312

6
g-index

10
all docs

10
docs citations

10
times ranked

704
citing authors

#	ARTICLE	IF	CITATIONS
1	A2AR Adenosine Signaling Suppresses Natural Killer Cell Maturation in the Tumor Microenvironment. <i>Cancer Research</i> , 2018, 78, 1003-1016.	0.4	269
2	Pediatric tolerogenic DCs expressing CD4 and immunoglobulin-like transcript receptor (ILT)4 secrete IL10 in response to Fc and adenosine. <i>European Journal of Immunology</i> , 2018, 48, 482-491.	1.6	15
3	Induction of antiinflammatory purinergic signaling in activated human iNKT cells. <i>JCI Insight</i> , 2018, 3, .	2.3	14
4	Randomized phase 2 trial of regadenoson for treatment of acute vaso-occlusive crises in sickle cell disease. <i>Blood Advances</i> , 2017, 1, 1645-1649.	2.5	38
5	Sickle cell vaso-occlusion causes activation of iNKT cells that is decreased by the adenosine A2A receptor agonist regadenoson. <i>Blood</i> , 2013, 121, 3329-3334.	0.6	87
6	NF- κ B Is Activated in CD4+ iNKT Cells by Sickle Cell Disease and Mediates Rapid Induction of Adenosine A2A Receptors. <i>PLoS ONE</i> , 2013, 8, e74664.	1.1	28
7	NF- κ B Activation Mediates Induction Of Anti-Inflammatory Adenosine A2A Receptors In iNKT Cells Of Sickle Cell Patients During Vaso-Occlusive Episodes and Upon Activation Of Cultured Human iNKT Cells. <i>Blood</i> , 2013, 122, 975-975.	0.6	0
8	Human Sickle Cell Disease Increases Numbers and Activation Of Peripheral Blood Myeloid Dendritic Cells, Monocytes, and Neutrophils. <i>Blood</i> , 2013, 122, 1033-1033.	0.6	0
9	The Role Of NF- κ B In The Activation Of Human iNKT Cells In Sickle Cell Disease Patients and In Vitro. <i>Blood</i> , 2013, 122, 2291-2291.	0.6	0
10	Regadenoson, An Adenosine 2A Receptor Agonist, Is Safe and Inhibits Invariant NKT Cells in Sickle Cell Disease. <i>Blood</i> , 2011, 118, 849-849.	0.6	0