## Mariam Alaverdashvili

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5521285/publications.pdf

Version: 2024-02-01

26 papers 1,024 citations

566801 15 h-index 26 g-index

26 all docs

26 does citations

26 times ranked

1269 citing authors

#	Article	IF	CITATIONS
1	Positive allosteric modulation of type 1 cannabinoid receptors reduces spike-and-wave discharges in Genetic Absence Epilepsy Rats from Strasbourg. Neuropharmacology, 2021, 190, 108553.	2.0	22
2	Preventing protein-energy malnutrition after cortical stroke enhances recovery of symmetry in forelimb use during spontaneous exploration. Applied Physiology, Nutrition and Metabolism, 2020, 45, 1015-1021.	0.9	1
3	Allosteric and orthosteric pharmacology of cannabidiol and cannabidiolâ€dimethylheptyl at the type 1 and type 2 cannabinoid receptors. British Journal of Pharmacology, 2019, 176, 1455-1469.	2.7	207
4	A longitudinal exploration of the impact of social anxiety and individual attachment on depression severity. Journal of Affective Disorders, 2019, 257, 250-256.	2.0	8
5	Protein-Energy Malnutrition Exacerbates Stroke-Induced Forelimb Abnormalities and Dampens Neuroinflammation. Translational Stroke Research, 2018, 9, 622-630.	2.3	12
6	The future of type 1 cannabinoid receptor allosteric ligands. Drug Metabolism Reviews, 2018, 50, 14-25.	1.5	26
7	Parallel changes in cortical neuron biochemistry and motor function in protein-energy malnourished adult rats. Neurolmage, 2017, 149, 275-284.	2.1	6
8	Mapping the dynamics of cortical neuroplasticity of skilled motor learning using micro X-ray fluorescence and histofluorescence imaging of zinc in the rat. Behavioural Brain Research, 2017, 318, 52-60.	1.2	4
9	Manganese-Enhanced Magnetic Resonance Imaging and Studies of Rat Behavior: Transient Motor Deficit in Skilled Reaching, Rears, and Activity in Rats After a Single Dose of MnCl <sub>2</sub> . Magnetic Resonance Insights, 2017, 10, 1178623X1770687.	2.5	4
10	Concurrent Glycogen and Lactate Imaging with FTIR Spectroscopy To Spatially Localize Metabolic Parameters of the Glial Response Following Brain Ischemia. Analytical Chemistry, 2016, 88, 10949-10956.	3.2	31
11	Laser system refinements to reduce variability in infarct size in the rat photothrombotic stroke model. Journal of Neuroscience Methods, 2015, 247, 58-66.	1.3	11
12	Protein-Energy Malnutrition Causes Deficits in Motor Function in Adult Male Rats. Journal of Nutrition, 2015, 145, 2503-2511.	1.3	13
13	Laminar-specific distribution of zinc: Evidence for presence of layer IV in forelimb motor cortex in the rat. Neurolmage, 2014, 103, 502-510.	2.1	14
14	A behavioral method for identifying recovery and compensation: Hand use in a preclinical stroke model using the single pellet reaching task. Neuroscience and Biobehavioral Reviews, 2013, 37, 950-967.	2.9	82
15	Compensation aids skilled reaching in aging and in recovery from forelimb motor cortex stroke in the rat. Neuroscience, 2010, 167, 21-30.	1.1	49
16	Thinning, movement, and volume loss of residual cortical tissue occurs after stroke in the adult rat as identified by histological and magnetic resonance imaging analysis. Neuroscience, 2010, 170, 123-137.	1.1	27
17	Both compensation and recovery of skilled reaching following small photothrombotic stroke to motor cortex in the rat. Experimental Neurology, 2009, 218, 145-153.	2.0	70
18	Nicotine does not improve recovery from learned nonuse nor enhance constraint-induced therapy after motor cortex stroke in the rat. Behavioural Brain Research, 2009, 198, 411-419.	1.2	8

#	Article	IF	CITATIONS
19	Similar hand shaping in reaching-for-food (skilled reaching) in rats and humans provides evidence of homology in release, collection, and manipulation movements. Behavioural Brain Research, 2009, 204, 153-161.	1.2	113
20	Motor cortex stroke impairs individual digit movement in skilled reaching by the rat. European Journal of Neuroscience, 2008, 28, 311-322.	1.2	58
21	Acute but not chronic differences in skilled reaching for food following motor cortex devascularization vs. photothrombotic stroke in the rat. Neuroscience, 2008, 157, 297-308.	1.1	34
22	"Learned baduse―limits recovery of skilled reaching for food after forelimb motor cortex stroke in rats: A new analysis of the effect of gestures on success. Behavioural Brain Research, 2008, 188, 281-290.	1.2	73
23	The problem of relating plasticity and skilled reaching after motor cortex stroke in the rat. Behavioural Brain Research, 2008, 192, 124-136.	1.2	76
24	Cineradiographic (video X-ray) analysis of skilled reaching in a single pellet reaching task provides insight into relative contribution of body, head, oral, and forelimb movement in rats. Behavioural Brain Research, 2008, 192, 232-247.	1.2	33
25	No improvement by amphetamine on learned non-use, attempts, success or movement in skilled reaching by the rat after motor cortex stroke. European Journal of Neuroscience, 2007, 25, 3442-3452.	1.2	26
26	Motor performance and behavior of immature rats are not compromised by a high dose of topiramate. Epilepsy and Behavior, 2005, 7, 222-230.	0.9	16