

Kang-Yi Su

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5524897/kang-yi-su-publications-by-citations.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

65

papers

1,896

citations

23

h-index

42

g-index

75

ext. papers

2,299

ext. citations

5.7

avg. IF

4.22

L-index

#	Paper	IF	Citations
65	Pretreatment epidermal growth factor receptor (EGFR) T790M mutation predicts shorter EGFR tyrosine kinase inhibitor response duration in patients with non-small-cell lung cancer. <i>Journal of Clinical Oncology</i> , 2012 , 30, 433-40	2.2	420
64	Slug confers resistance to the epidermal growth factor receptor tyrosine kinase inhibitor. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2011 , 183, 1071-9	10.2	129
63	Enterovirus-induced miR-141 contributes to shutoff of host protein translation by targeting the translation initiation factor eIF4E. <i>Cell Host and Microbe</i> , 2011 , 9, 58-69	23.4	121
62	Outcomes in patients with non-small-cell lung cancer and acquired Thr790Met mutation treated with osimertinib: a genomic study. <i>Lancet Respiratory Medicine</i> , 2018 , 6, 107-116	35.1	86
61	Mice deficient in collapsin response mediator protein-1 exhibit impaired long-term potentiation and impaired spatial learning and memory. <i>Journal of Neuroscience</i> , 2007 , 27, 2513-24	6.6	76
60	Proteogenomics of Non-smoking Lung Cancer in East Asia Delineates Molecular Signatures of Pathogenesis and Progression. <i>Cell</i> , 2020 , 182, 226-244.e17	56.2	65
59	R331W Missense Mutation of Oncogene YAP1 Is a Germline Risk Allele for Lung Adenocarcinoma With Medical Actionability. <i>Journal of Clinical Oncology</i> , 2015 , 33, 2303-10	2.2	62
58	Identification of five driver gene mutations in patients with treatment-naïve lung adenocarcinoma in Taiwan. <i>PLoS ONE</i> , 2015 , 10, e0120852	3.7	58
57	Dynamic plasma EGFR mutation status as a predictor of EGFR-TKI efficacy in patients with EGFR-mutant lung adenocarcinoma. <i>Journal of Thoracic Oncology</i> , 2015 , 10, 603-10	8.9	55
56	Coexistence of EGFR T790M mutation and common activating mutations in pretreatment non-small cell lung cancer: A systematic review and meta-analysis. <i>Lung Cancer</i> , 2016 , 94, 46-53	5.9	48
55	Rutin, a flavonoid and principal component of <i>saussurea involucreata</i> , attenuates physical fatigue in a forced swimming mouse model. <i>International Journal of Medical Sciences</i> , 2014 , 11, 528-37	3.7	48
54	EGFR mutation, smoking, and gender in advanced lung adenocarcinoma. <i>Oncotarget</i> , 2017 , 8, 98384-98393	3.3	43
53	3,4-Dihydroxytoluene, a metabolite of rutin, inhibits inflammatory responses in lipopolysaccharide-activated macrophages by reducing the activation of NF- κ B signaling. <i>BMC Complementary and Alternative Medicine</i> , 2014 , 14, 21	4.7	36
52	High PD-L1 expression correlates with primary resistance to EGFR-TKIs in treatment naïve advanced EGFR-mutant lung adenocarcinoma patients. <i>Lung Cancer</i> , 2019 , 127, 37-43	5.9	35
51	Tumor PD-L1 Expression and Clinical Outcomes in Advanced-stage Non-Small Cell Lung Cancer Patients Treated with Nivolumab or Pembrolizumab: Real-World Data in Taiwan. <i>Journal of Cancer</i> , 2018 , 9, 1813-1820	4.5	34
50	Clustered genomic alterations in chromosome 7p dictate outcomes and targeted treatment responses of lung adenocarcinoma with EGFR-activating mutations. <i>Journal of Clinical Oncology</i> , 2011 , 29, 3435-42	2.2	28
49	The Association of Acquired T790M Mutation with Clinical Characteristics after Resistance to First-Line Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor in Lung Adenocarcinoma. <i>Cancer Research and Treatment</i> , 2018 , 50, 1294-1303	5.2	28

48	Association between programmed death-ligand 1 expression, immune microenvironments, and clinical outcomes in epidermal growth factor receptor mutant lung adenocarcinoma patients treated with tyrosine kinase inhibitors. <i>European Journal of Cancer</i> , 2020 , 124, 110-122	7.5	28
47	Shisa3 is associated with prolonged survival through promoting Eatenin degradation in lung cancer. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2014 , 190, 433-44	10.2	27
46	Serine protease hepsin regulates hepatocyte size and hemodynamic retention of tumor cells by hepatocyte growth factor signaling in mice. <i>Hepatology</i> , 2012 , 56, 1913-23	11.2	25
45	Rutin, a Flavonoid That Is a Main Component of <i>Saussurea involucreta</i> , Attenuates the Senescence Effect in D-Galactose Aging Mouse Model. <i>Evidence-based Complementary and Alternative Medicine</i> , 2012 , 2012, 980276	2.3	25
44	Taiwanese Green Propolis and Propolin G Protect the Liver from the Pathogenesis of Fibrosis via Eliminating TGF- β -Induced Smad2/3 Phosphorylation. <i>Journal of Agricultural and Food Chemistry</i> , 2014 , 62, 3192-3201	5.7	24
43	HOXA5 inhibits metastasis via regulating cytoskeletal remodelling and associates with prolonged survival in non-small-cell lung carcinoma. <i>PLoS ONE</i> , 2015 , 10, e0124191	3.7	24
42	Clinicopathological and genomic comparisons between different histologic components in combined small cell lung cancer and non-small cell lung cancer. <i>Lung Cancer</i> , 2018 , 125, 282-290	5.9	23
41	Potential Therapeutic Role of Hispidulin in Gastric Cancer through Induction of Apoptosis via NAG-1 Signaling. <i>Evidence-based Complementary and Alternative Medicine</i> , 2013 , 2013, 518301	2.3	22
40	JAG1 Is Associated with Poor Survival through Inducing Metastasis in Lung Cancer. <i>PLoS ONE</i> , 2016 , 11, e0150355	3.7	22
39	SPANXA suppresses EMT by inhibiting c-JUN/SNAI2 signaling in lung adenocarcinoma. <i>Oncotarget</i> , 2016 , 7, 44417-44429	3.3	21
38	Decreased expressions of hepsin in human hepatocellular carcinomas. <i>Liver International</i> , 2006 , 26, 774-809	8.0	17
37	The emergence of T790M mutation in EGFR-mutant lung adenocarcinoma patients having a history of acquired resistance to EGFR-TKI: focus on rebiopsy timing and long-term existence of T790M. <i>Oncotarget</i> , 2016 , 7, 48059-48069	3.3	17
36	Characteristics and Predictive Value of PD-L1 Status in Real-World Non-Small Cell Lung Cancer Patients. <i>Journal of Immunotherapy</i> , 2018 , 41, 292-299	5	16
35	Molecular gene signature and prognosis of non-small cell lung cancer. <i>Oncotarget</i> , 2016 , 7, 51898-51907	3.3	15
34	Small Molecule T315 Promotes Casitas B-Lineage Lymphoma-Dependent Degradation of Epidermal Growth Factor Receptor via Y1045 Autophosphorylation. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2016 , 193, 753-66	10.2	14
33	FAM198B Is Associated with Prolonged Survival and Inhibits Metastasis in Lung Adenocarcinoma via Blockage of ERK-Mediated MMP-1 Expression. <i>Clinical Cancer Research</i> , 2018 , 24, 916-926	12.9	14
32	Prior EGFR tyrosine-kinase inhibitor therapy did not influence the efficacy of subsequent pemetrexed plus platinum in advanced chemonaïve patients with EGFR-mutant lung adenocarcinoma. <i>OncoTargets and Therapy</i> , 2014 , 7, 799-805	4.4	12
31	Potential Therapeutic Role of Z-Isochaihulactone in Lung Cancer through Induction of Apoptosis via Notch Signaling. <i>Evidence-based Complementary and Alternative Medicine</i> , 2012 , 2012, 809204	2.3	12

30	Perinatal polychlorinated biphenyls and polychlorinated dibenzofurans exposure are associated with DNA methylation changes lasting to early adulthood: Findings from Yucheng second generation. <i>Environmental Research</i> , 2019 , 170, 481-486	7.9	12
29	Association of Programmed Death-Ligand 1 Expression with Fusion Variants and Clinical Outcomes in Patients with Anaplastic Lymphoma Kinase-Positive Lung Adenocarcinoma Receiving Crizotinib. <i>Oncologist</i> , 2020 , 25, 702-711	5.7	11
28	EGFR mutation and lobar location of lung adenocarcinoma. <i>Carcinogenesis</i> , 2016 , 37, 157-162	4.6	10
27	R26R-GR: a Cre-activable dual fluorescent protein reporter mouse. <i>PLoS ONE</i> , 2012 , 7, e46171	3.7	10
26	Impact of EGFR mutation detection methods on the efficacy of erlotinib in patients with advanced EGFR-wild type lung adenocarcinoma. <i>PLoS ONE</i> , 2014 , 9, e107160	3.7	10
25	Mutational monitoring of EGFR T790M in cfDNA for clinical outcome prediction in EGFR-mutant lung adenocarcinoma. <i>PLoS ONE</i> , 2018 , 13, e0207001	3.7	10
24	Gene expression of MAGE-A3 and PRAME tumor antigens and EGFR mutational status in Taiwanese non-small cell lung cancer patients. <i>Asia-Pacific Journal of Clinical Oncology</i> , 2017 , 13, e212-e223	1.9	9
23	Dichloromethane extracts of propolis protect cell from oxygen-glucose deprivation-induced oxidative stress via reducing apoptosis. <i>Food and Nutrition Research</i> , 2016 , 60, 30081	3.1	9
22	Implementation and Quality Control of Lung Cancer EGFR Genetic Testing by MALDI-TOF Mass Spectrometry in Taiwan Clinical Practice. <i>Scientific Reports</i> , 2016 , 6, 30944	4.9	9
21	Novel therapeutic drug identification and gene correlation for fatty liver disease using high-content screening: Proof of concept. <i>European Journal of Pharmaceutical Sciences</i> , 2018 , 121, 106-117	5.7	9
20	ALK variants, PD-L1 expression, and their association with outcomes in ALK-positive NSCLC patients. <i>Scientific Reports</i> , 2020 , 10, 21063	4.9	8
19	Rapid Sputum Multiplex Detection of the M. tuberculosis Complex (MTBC) and Resistance Mutations for Eight Antibiotics by Nucleotide MALDI-TOF MS. <i>Scientific Reports</i> , 2017 , 7, 41486	4.9	7
18	DNA polymerase I proofreading exonuclease activity is required for endonuclease V repair pathway both in vitro and in vivo. <i>DNA Repair</i> , 2018 , 64, 59-67	4.3	7
17	Higher frequency but random distribution of EGFR mutation subtypes in familial lung cancer patients. <i>Oncotarget</i> , 2016 , 7, 53299-53308	3.3	6
16	Metabolomics Investigation of Voriconazole-Induced Hepatotoxicity in Mice. <i>Chemical Research in Toxicology</i> , 2019 , 32, 1840-1849	4	5
15	Improved Dried Blood Spot-Based Metabolomics Analysis by a Postcolumn Infused-Internal Standard Assisted Liquid Chromatography-Electrospray Ionization Mass Spectrometry Method. <i>Analytical Chemistry</i> , 2019 , 91, 10702-10712	7.8	4
14	Intercalated Treatment Following Rebiopsy Is Associated with a Shorter Progression-Free Survival of Osimertinib Treatment. <i>Cancer Research and Treatment</i> , 2018 , 50, 1164-1174	5.2	4
13	Primary Tumor Resection Is Associated with a Better Outcome among Advanced EGFR-Mutant Lung Adenocarcinoma Patients Receiving EGFR-TKI Treatment. <i>Oncology</i> , 2021 , 99, 32-40	3.6	4

12	Measurement of uracil-DNA glycosylase activity by matrix assisted laser desorption/ionization time-of-flight mass spectrometry technique. <i>DNA Repair</i> , 2021 , 97, 103028	4.3	4
11	Application of single nucleotide extension and MALDI-TOF mass spectrometry in proofreading and DNA repair assay. <i>DNA Repair</i> , 2018 , 61, 63-75	4.3	4
10	Copy Number Alterations of Depressed Colorectal Neoplasm Predict the Survival and Response to Oxaliplatin in Proximal Colon Cancer. <i>Cancers</i> , 2020 , 12,	6.6	2
9	The impact of different first-line EGFR-TKIs on the clinical outcome of sequential osimertinib treatment in advanced NSCLC with secondary T790M. <i>Scientific Reports</i> , 2021 , 11, 12084	4.9	2
8	Prognostic Characteristics and Immunotherapy Response of Patients With Nonsquamous NSCLC With Mutation in East Asian Populations: A Single-Center Cohort Study in Taiwan. <i>JTO Clinical and Research Reports</i> , 2021 , 2, 100140	1.4	2
7	Proofreading and DNA Repair Assay Using Single Nucleotide Extension and MALDI-TOF Mass Spectrometry Analysis. <i>Journal of Visualized Experiments</i> , 2018 ,	1.6	2
6	Proofreading of single nucleotide insertion/deletion replication errors analyzed by MALDI-TOF mass spectrometry assay. <i>DNA Repair</i> , 2020 , 88, 102810	4.3	1
5	Predilection of contralateral upper lung metastasis in upper lobe lung adenocarcinoma patients. <i>Journal of Thoracic Disease</i> , 2016 , 8, 86-92	2.6	1
4	Transthyretin as a Biomarker to Predict and Monitor Major Depressive Disorder Identified by Whole-Genome Transcriptomic Analysis in Mouse Models. <i>Biomedicines</i> , 2021 , 9,	4.8	1
3	The Difference in Clinical Outcomes Between Osimertinib and Afatinib for First-Line Treatment in Patients with Advanced and Recurrent EGFR-Mutant Non-Small Cell Lung Cancer in Taiwan.. <i>Targeted Oncology</i> , 2022 , 1	5	1
2	Improvement of NGS uniformity of single cell sequencing in CTCs by WGA QC and increased sequencing throughput.. <i>Journal of Clinical Oncology</i> , 2018 , 36, e21176-e21176	2.2	
1	Multiplex ALK, RET, and ROS1 fusion mutation detection in FFPE from lung cancer patients by MALDI-TOF mass spectrometry.. <i>Journal of Clinical Oncology</i> , 2017 , 35, e13103-e13103	2.2	