Hidetomo Murakami

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5530903/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Face pareidolia is associated with right striatal dysfunction in drug-naÃ⁻ve patients with Parkinson's disease. Neurological Sciences, 2021, 42, 5327-5334.	1.9	6
2	Differences in correlations of depression and anhedonia with cardiovascular sympathetic functions during a head-up tilt test in drug-naÃ⁻ve Parkinson's disease patients. Neurological Sciences, 2020, 41, 2825-2830.	1.9	4
3	Mini Review: Correlations of Cognitive Domains With Cerebrospinal Fluid α-Synuclein Levels in Patients With Parkinson's Disease. Frontiers in Aging Neuroscience, 2020, 12, 616357.	3.4	3
4	Correlated levels of cerebrospinal fluid pathogenic proteins in drug-naÃ⁻ve Parkinson's disease. BMC Neurology, 2019, 19, 113.	1.8	11
5	Effect of istradefylline on mood disorders in Parkinson's disease. Journal of the Neurological Sciences, 2019, 396, 78-83.	0.6	35
6	Japanese multicenter database of healthy controls for [1231]FP-CIT SPECT. European Journal of Nuclear Medicine and Molecular Imaging, 2018, 45, 1405-1416.	6.4	80
7	P3â€412: ¹²³ lâ€MIBG MYOCARDIAL SCINTIGRAPHY CAN BE A MARKER OF LANGUAGE FUNCTION IN <i>DE NOVO</i> PARKINSON'S DISEASE. Alzheimer's and Dementia, 2018, 14, P1263.	^V 0.8	0
8	Usefulness Differs Between the Visual Assessment and Specific Binding Ratio of 123I-Ioflupane SPECT in Assessing Clinical Symptoms of Drug-NaÃīve Parkinson's Disease Patients. Frontiers in Aging Neuroscience, 2018, 10, 412.	3.4	7
9	The subjective perception of past, present, and future time in patients with Alzheimer's disease: a qualitative study. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 3185-3192.	2.2	4
10	Cerebrospinal fluid 5-HIAA concentrations correlate with cardiac uptake of 123I-MIBG during myocardial scintigraphy in drug naÃ⁻ve Parkinson's disease. Journal of Neural Transmission, 2018, 125, 1511-1514.	2.8	2
11	New mode of burst spinal cord stimulation improved mental status as well as motor function in a patient with Parkinson's disease. Parkinsonism and Related Disorders, 2018, 57, 82-83.	2.2	23
12	[P3–289]: COGNITIVE DOMAIN CORRELATES WITH STRIATAL ACCUMULATION OF DOPAMINE TRANSPORTER SCINTIGRAPHY IN DRUG NAÃVE PARKINSON'S DISEASE. Alzheimer's and Dementia, 2017, 13, P1053.	0.8	0
13	Accumulation of 123I-Ioflupane Is a Useful Marker of the Efficacy of Selegiline Monotherapy in Drug-NaÃ⁻ve Parkinson's Disease. Frontiers in Aging Neuroscience, 2017, 9, 321.	3.4	4
14	Effects of dopaminergic drug adjustment on executive function in different clinical stages of Parkinson's disease. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 2719-2726.	2.2	10
15	Improvement in Language Function Correlates with Gait Improvement in Drug-naÃīve Parkinson's Disease Patients Taking Dopaminergic Medication. Journal of Parkinson's Disease, 2016, 6, 209-217.	2.8	3
16	Increased detection of mild cognitive impairment with type 2 diabetes mellitus using the Japanese version of the Montreal Cognitive Assessment: A pilot study. Neurology and Clinical Neuroscience, 2015, 3, 89-93.	0.4	7
17	Modified <scp>S</scp> ix <scp>E</scp> lements <scp>T</scp> est: <scp>E</scp> arlier diagnosis of the correlation between motor and executive dysfunction in <scp>P</scp> arkinson's disease without dementia. Neurology and Clinical Neuroscience, 2015, 3, 209-214.	0.4	3
18	Correlation between motor and cognitive functions in the progressive course of Parkinson's disease. Neurology and Clinical Neuroscience, 2013, 1, 172-176.	0.4	9

#	Article	IF	CITATIONS
19	The Montreal Cognitive Assessment and Neurobehavioral Cognitive Status Examination are useful for screening mild cognitive impairment in Japanese patients with Parkinson's disease. Neurology and Clinical Neuroscience, 2013, 1, 103-108.	0.4	14
20	Altered Fibrinogen and Prothrombin mRNA Expression in Streptozotocin-induced Diabetic Rats. The Showa University Journal of Medical Sciences, 2000, 12, 295-302.	0.1	2