Yongxiang Wei

List of Publications by Year
in descending order

Source: https://exaly.com/author-pdf/5533546/publications.pdf

Version: 2024-02-01

10 papers	102 citations	1683354 5 h-index	1372195 10 g-index
10 all docs	10 docs citations	10 times ranked	119 citing authors

#	Article	IF	Citations
1	ESMâ€1 promotes adhesion between monocytes and endothelial cells under intermittent hypoxia. Journal of Cellular Physiology, 2019, 234, 1512-1521.	2.0	43
2	Obstructive sleep apnea increases the risk of cardiovascular damage: a systematic review and meta-analysis of imaging studies. Systematic Reviews, 2021, 10, 212.	2.5	16
3	Circulating ESM-1 levels are correlated with the presence of coronary artery disease in patients with obstructive sleep apnea. Respiratory Research, 2019, 20, 188.	1.4	11
4	Targeted Sequencing Analysis of the Leptin Receptor Gene Identifies Variants Associated with Obstructive Sleep Apnoea in Chinese Han Population. Lung, 2019, 197, 577-584.	1.4	6
5	Targeted sequencing analysis of the adiponectin gene identifies variants associated with obstructive sleep apnoea in Chinese Han population. Medicine (United States), 2019, 98, e15219.	0.4	5
6	Combined Association Between ADIPOQ, PPARG, and TNF Genes Variants and Obstructive Sleep Apnea in Chinese Han Population. Nature and Science of Sleep, 2022, Volume 14, 363-372.	1.4	5
7	Increasing circulating ESM-1 and adhesion molecules are associated with earlystage atherosclerosis in OSA patients:A cross-sectional study. Sleep Medicine, 2022, , .	0.8	5
8	Targeted sequencing analysis of PPARG identifies a risk variant associated with obstructive sleep apnea in Chinese Han subjects. Sleep and Breathing, 2020, 24, 167-174.	0.9	4
9	EMMPRIN: A potential biomarker for predicting the presence of obstructive sleep apnea. Clinica Chimica Acta, 2020, 510, 317-322.	0.5	4
10	Increased levels of VCAM-1 is associated with higher occurrence of coronary artery disease in adults with moderate to severe obstructive sleep apnea. Sleep Medicine, 2021, 85, 131-137.	0.8	3