

Sarah K Kirschner

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5548223/publications.pdf>

Version: 2024-02-01

9
papers

61
citations

1937685

4
h-index

1720034

7
g-index

9
all docs

9
docs citations

9
times ranked

64
citing authors

#	ARTICLE	IF	CITATIONS
1	Intestinal function is impaired in patients with Chronic Obstructive Pulmonary Disease. <i>Clinical Nutrition</i> , 2021, 40, 2270-2277.	5.0	23
2	Risk Factors for Postural and Functional Balance Impairment in Patients with Chronic Obstructive Pulmonary Disease. <i>Journal of Clinical Medicine</i> , 2020, 9, 609.	2.4	13
3	Impaired intestinal function is associated with lower muscle and cognitive health and well-being in patients with congestive heart failure. <i>Journal of Parenteral and Enteral Nutrition</i> , 2022, 46, 660-670.	2.6	11
4	Transorgan short-chain fatty acid fluxes in the fasted and postprandial state in the pig. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2021, 321, E665-E673.	3.5	8
5	Intestinal dysfunction in chronic disease. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2021, 24, 464-472.	2.5	2
6	Plasma Short-Chain Fatty Acid (SCFA) Concentrations Are Related to Cognition and Adipose Tissue Mass but Not Muscle Function in Older Adults. <i>Current Developments in Nutrition</i> , 2022, 6, 33.	0.3	2
7	Lower Plasma Short-Chain Fatty Acids Are Associated With Increased Leg Muscle Fatigue in (Morbidly) Obese Adults. <i>Current Developments in Nutrition</i> , 2021, 5, 1258.	0.3	1
8	Whole Body Production of Butyrate, but Not of Acetate or Propionate, Is Reduced in Adults With Prediabetes. <i>Current Developments in Nutrition</i> , 2022, 6, 35.	0.3	1
9	Reduced Plasma Short-Chain Fatty Acid (SCFA) Concentrations Are Not Associated With Markers of Muscle Health in Non-Small Cell Lung Cancer. <i>Current Developments in Nutrition</i> , 2021, 5, 270.	0.3	0