Celine Melo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5555899/publications.pdf

Version: 2024-02-01

933447 839539 34 375 10 18 citations h-index g-index papers 34 34 34 419 citing authors docs citations times ranked all docs

#	Article	IF	CITATIONS
1	Predators of Quail Eggs, and the Evidence of the Remains: Implications for Nest Predation Studies. Condor, 1998, 100, 395-399.	1.6	78
2	Confirmation of Infanticide in the Communally Breeding Guira Cuckoo. Auk, 1999, 116, 847-851.	1.4	41
3	Micronucleus as biomarker of genotoxicity in birds from Brazilian Cerrado. Ecotoxicology and Environmental Safety, 2015, 115, 223-228.	6.0	34
4	Avian frugivory in <i>Miconia</i> (Melastomataceae): contrasting fruiting times promote habitat complementarity between savanna and palm swamp. Journal of Tropical Ecology, 2013, 29, 99-109.	1.1	25
5	What is on the menu for frugivorous birds in the Cerrado? Fruiting phenology and nutritional traits highlight the importance of habitat complementarity. Acta Botanica Brasilica, 2019, 33, 572-583.	0.8	19
6	Frugivory and dispersal of Faramea cyanea (Rubiaceae) in Cerrado woody plant formations. Brazilian Journal of Biology, 2003, 63, 75-82.	0.9	18
7	Ticks on birds in a savanna (Cerrado) reserve on the outskirts of Uberlândia, Minas Gerais, Brazil. Brazilian Journal of Veterinary Parasitology, 2013, 22, 46-52.	0.7	18
8	Dominant frequency of songs in tropical bird species is higher in sites with high noise pollution. Environmental Pollution, 2018, 235, 983-992.	7.5	16
9	Effect of urbanization on the micronucleus frequency in birds from forest fragments. Ecotoxicology and Environmental Safety, 2019, 171, 631-637.	6.0	14
10	Individual–resource networks reveal distinct fruit preferences of selective individuals from a generalist population of the Helmeted Manakin. Ibis, 2020, 162, 713-722.	1.9	12
11	Modularity in ecological networks between frugivorous birds and congeneric plant species. Journal of Tropical Ecology, 2016, 32, 526-535.	1.1	11
12	Effects of urbanisation and pollution on the heterophil/lymphocyte ratio in birds from Brazilian Cerrado. Environmental Science and Pollution Research, 2022, 29, 40204-40212.	5.3	8
13	Frugivory and potential of birds as dispersers of Siparuna guianensis. Brazilian Journal of Biology, 2015, 75, 300-304.	0.9	7
14	Micronucleus and different nuclear abnormalities in wild birds in the Cerrado, Brazil. Environmental Science and Pollution Research, 2022, 29, 14279-14287.	5. 3	7
15	Frugivory in Lacistema hasslerianum Chodat (Lacistemaceae), a gallery forest understory treelet in Central Brazil. Brazilian Journal of Biology, 2009, 69, 201-207.	0.9	6
16	Plastic responses in tree architecture and specific leaf area of Xylopia aromatica (Annonaceae): adaptations to environments with different light intensities. Revista Brasileira De Botanica, 2013, 36, 279-283.	1.3	6
17	Overlap and resource sharing in coteries of fruit-eating birds. Journal of Tropical Ecology, 2013, 29, 409-416.	1.1	6
18	Effects of urban proximity and the occurrence of erythroplastids in Antilophia galeata. Environmental Science and Pollution Research, 2020, 27, 44650-44655.	5. 3	6

#	Article	IF	CITATIONS
19	Artificial roosts as seed dispersal nuclei in a cerrado area in Triângulo Mineiro, Brazil. Bioscience Journal, 2016, 32, 514-523.	0.4	6
20	Keeping safe and fed: large heterospecific shorebird flocks to decrease intraspecific competition. Journal of Avian Biology, 2020, 51, .	1.2	5
21	Frugivory and seed dispersal of Miconia theaezans (Bonpl.) Cogniaux (Melastomataceae) by birds in a transition palm swamp: gallery forest in Central Brazil. Brazilian Journal of Biology, 2012, 72, 25-31.	0.9	5
22	Impact of Forpus xanthopterygius (Spix, 1824) (Aves, Psittacidae) on flowers of Handroanthus serratifolius (Vahl.) S. O. Grose (Bignoniaceae). Brazilian Journal of Biology, 2009, 69, 1149-1151.	0.9	4
23	Richness, composition and detectability of Psittacidae (Aves) in three palm swamps of the Cerrado sensu lato in central Brazil. Revista Chilena De Historia Natural, 2012, 85, 171-178.	1.2	4
24	Male Helmeted Manakins (Antilophia galeata) with more colorful crowns have better body conditions. Wilson Journal of Ornithology, 2017, 129, 158-163.	0.2	3
25	Fruit colour and edge effects poorly explains frugivorous bird-plant interactions in disturbed semideciduous forests. Acta Scientiarum - Biological Sciences, 2018, 40, 40041.	0.3	3
26	Seasonal consistency of the assembling rules in a bird-fruiting plant network. Ecological Complexity, 2020, 44, 100869.	2.9	3
27	Disentangling abiotic and biotic mechanisms behind the formation of heterospecific Nearctic-Neotropical shorebird flocks. Evolutionary Ecology, 2020, 34, 1047-1061.	1.2	3
28	Bills Favor Mining and Threaten Conservation of Brazilian Merganser (MergusÂoctosetaceus) at Serra da Canastra National Park, Minas Gerais, Brazil. Natureza A Conservacao, 2012, 10, 64-71.	2.5	2
29	Haemosporidian parasites prevalence associated with physical conditioning of avian species from the Brazilian Cerrado. Ciência E Natura, 0, 42, e50.	0.0	2
30	First record of microfilariae in Antilophia galeata (Aves: Pipridae). Acta Brasiliensis, 2020, 4, 106.	0.2	1
31	Leukocyte profile of the helmeted manakin, Antilophia galeata (Passeriformes: Pipridae) in a Cerrado forest fragment. Zoologia, 0, 37, 1-9.	0.5	1
32	Seasonality drives variation in the use of forest strata by adult males of a dimorphic frugivorous bird species. Austral Ecology, 0, , .	1.5	1
33	POPULATION ABUNDANCE OF Fluvicola nengeta (LINNAEUS, 1766) (AVES: TYRANNIDAE) IN A URBAN PARK. Oecologia Australis, 2017, 21, 444-449.	0.2	0
34	Forest dependent birds are the main frugivorous species in mutualistic networks from the Brazilian Cerrado. Ecoscience, 0, , 1-9.	1.4	0