

# Celine Melo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5555899/publications.pdf>

Version: 2024-02-01

34  
papers

375  
citations

933447

10  
h-index

839539

18  
g-index

34  
all docs

34  
docs citations

34  
times ranked

419  
citing authors

#	ARTICLE	IF	CITATIONS
1	Micronucleus and different nuclear abnormalities in wild birds in the Cerrado, Brazil. <i>Environmental Science and Pollution Research</i> , 2022, 29, 14279-14287.	5.3	7
2	Effects of urbanisation and pollution on the heterophil/lymphocyte ratio in birds from Brazilian Cerrado. <i>Environmental Science and Pollution Research</i> , 2022, 29, 40204-40212.	5.3	8
3	Individual "resource networks reveal distinct fruit preferences of selective individuals from a generalist population of the Helmeted Manakin. <i>Ibis</i> , 2020, 162, 713-722.	1.9	12
4	Keeping safe and fed: large heterospecific shorebird flocks to decrease intraspecific competition. <i>Journal of Avian Biology</i> , 2020, 51, .	1.2	5
5	Seasonal consistency of the assembling rules in a bird-fruited plant network. <i>Ecological Complexity</i> , 2020, 44, 100869.	2.9	3
6	Effects of urban proximity and the occurrence of erythroplastids in <i>Antilophia galeata</i> . <i>Environmental Science and Pollution Research</i> , 2020, 27, 44650-44655.	5.3	6
7	Disentangling abiotic and biotic mechanisms behind the formation of heterospecific Nearctic-Neotropical shorebird flocks. <i>Evolutionary Ecology</i> , 2020, 34, 1047-1061.	1.2	3
8	First record of microfilariae in <i>Antilophia galeata</i> (Aves: Pipridae). <i>Acta Brasiliensis</i> , 2020, 4, 106.	0.2	1
9	Effect of urbanization on the micronucleus frequency in birds from forest fragments. <i>Ecotoxicology and Environmental Safety</i> , 2019, 171, 631-637.	6.0	14
10	What is on the menu for frugivorous birds in the Cerrado? Fruiting phenology and nutritional traits highlight the importance of habitat complementarity. <i>Acta Botanica Brasilica</i> , 2019, 33, 572-583.	0.8	19
11	Dominant frequency of songs in tropical bird species is higher in sites with high noise pollution. <i>Environmental Pollution</i> , 2018, 235, 983-992.	7.5	16
12	&lt;b&gt;Fruit colour and edge effects poorly explains frugivorous bird-plant interactions in disturbed semideciduous forests. <i>Acta Scientiarum - Biological Sciences</i> , 2018, 40, 40041.	0.3	3
13	Male Helmeted Manakins ( <i>Antilophia galeata</i> ) with more colorful crowns have better body conditions. <i>Wilson Journal of Ornithology</i> , 2017, 129, 158-163.	0.2	3
14	POPULATION ABUNDANCE OF <i>Fluvicola nengeta</i> (LINNAEUS, 1766) (AVES: TYRANNIDAE) IN A URBAN PARK. <i>Oecologia Australis</i> , 2017, 21, 444-449.	0.2	0
15	Modularity in ecological networks between frugivorous birds and congeneric plant species. <i>Journal of Tropical Ecology</i> , 2016, 32, 526-535.	1.1	11
16	Artificial roosts as seed dispersal nuclei in a cerrado area in Triângulo Mineiro, Brazil. <i>Bioscience Journal</i> , 2016, 32, 514-523.	0.4	6
17	Frugivory and potential of birds as dispersers of <i>Siparuna guianensis</i> . <i>Brazilian Journal of Biology</i> , 2015, 75, 300-304.	0.9	7
18	Micronucleus as biomarker of genotoxicity in birds from Brazilian Cerrado. <i>Ecotoxicology and Environmental Safety</i> , 2015, 115, 223-228.	6.0	34

#	ARTICLE	IF	CITATIONS
19	Plastic responses in tree architecture and specific leaf area of <i>Xylopia aromatica</i> (Annonaceae): adaptations to environments with different light intensities. <i>Revista Brasileira De Botanica</i> , 2013, 36, 279-283.	1.3	6
20	Avian frugivory in <i>Miconia</i> (Melastomataceae): contrasting fruiting times promote habitat complementarity between savanna and palm swamp. <i>Journal of Tropical Ecology</i> , 2013, 29, 99-109.	1.1	25
21	Overlap and resource sharing in coterries of fruit-eating birds. <i>Journal of Tropical Ecology</i> , 2013, 29, 409-416.	1.1	6
22	Ticks on birds in a savanna (Cerrado) reserve on the outskirts of Uberlândia, Minas Gerais, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2013, 22, 46-52.	0.7	18
23	Richness, composition and detectability of Psittacidae (Aves) in three palm swamps of the Cerrado sensu lato in central Brazil. <i>Revista Chilena De Historia Natural</i> , 2012, 85, 171-178.	1.2	4
24	Frugivory and seed dispersal of <i>Miconia theaezans</i> (Bonpl.) Cogniaux (Melastomataceae) by birds in a transition palm swamp: gallery forest in Central Brazil. <i>Brazilian Journal of Biology</i> , 2012, 72, 25-31.	0.9	5
25	Bills Favor Mining and Threaten Conservation of Brazilian Merganser ( <i>Mergus octosetaceus</i> ) at Serra da Canastra National Park, Minas Gerais, Brazil. <i>Natureza A Conservacao</i> , 2012, 10, 64-71.	2.5	2
26	Frugivory in <i>Lacistema hasslerianum</i> Chodat (Lacistemaceae), a gallery forest understory treelet in Central Brazil. <i>Brazilian Journal of Biology</i> , 2009, 69, 201-207.	0.9	6
27	Impact of <i>Forpus xanthopterygius</i> (Spix, 1824) (Aves, Psittacidae) on flowers of <i>Handroanthus serratifolius</i> (Vahl.) S. O. Grose (Bignoniaceae). <i>Brazilian Journal of Biology</i> , 2009, 69, 1149-1151.	0.9	4
28	Frugivory and dispersal of <i>Faramea cyanea</i> (Rubiaceae) in Cerrado woody plant formations. <i>Brazilian Journal of Biology</i> , 2003, 63, 75-82.	0.9	18
29	Confirmation of Infanticide in the Communally Breeding Guira Cuckoo. <i>Auk</i> , 1999, 116, 847-851.	1.4	41
30	Predators of Quail Eggs, and the Evidence of the Remains: Implications for Nest Predation Studies. <i>Condor</i> , 1998, 100, 395-399.	1.6	78
31	Haemosporidian parasites prevalence associated with physical conditioning of avian species from the Brazilian Cerrado. <i>Ciência E Natura</i> , 0, 42, e50.	0.0	2
32	Leukocyte profile of the helmeted manakin, <i>Antilophia galeata</i> (Passeriformes: Pipridae) in a Cerrado forest fragment. <i>Zoologia</i> , 0, 37, 1-9.	0.5	1
33	Forest dependent birds are the main frugivorous species in mutualistic networks from the Brazilian Cerrado. <i>Écoscience</i> , 0, , 1-9.	1.4	0
34	Seasonality drives variation in the use of forest strata by adult males of a dimorphic frugivorous bird species. <i>Austral Ecology</i> , 0, , .	1.5	1