

Fang Zeng

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5557597/fang-zeng-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

152
papers

6,166
citations

46
h-index

72
g-index

155
ext. papers

7,012
ext. citations

7.2
avg. IF

6.22
L-index

#	Paper	IF	Citations
152	Biomarker-activatable probes based on smart AIEgens for fluorescence and optoacoustic imaging. <i>Coordination Chemistry Reviews</i> , 2022 , 458, 214438	23.2	8
151	Targeted and activatable nanosystem for fluorescent and optoacoustic imaging of immune-mediated inflammatory diseases and therapy via inhibiting NF- κ B/NLRP3 pathways.. <i>Bioactive Materials</i> , 2022 , 10, 79-92	16.7	2
150	An AIEgen-based oral-administration nanosystem for detection and therapy of ulcerative colitis via 3D-MSOT/NIR-II fluorescent imaging and inhibiting NLRP3 inflammasome.. <i>Biomaterials</i> , 2022 , 283, 121468	15.6	4
149	An activatable probe for detection and therapy of food-additive-related hepatic injury via NIR-II fluorescence/optoacoustic imaging and biomarker-triggered drug release.. <i>Analytica Chimica Acta</i> , 2022 , 1208, 339831	6.6	4
148	A HO-activatable nanoprobe for diagnosing interstitial cystitis and liver ischemia-reperfusion injury via multispectral optoacoustic tomography and NIR-II fluorescent imaging. <i>Nature Communications</i> , 2021 , 12, 6870	17.4	13
147	Fluorophore-Dapagliflozin Dyad for Detecting Diabetic Liver/Kidney Damages via Fluorescent Imaging and Treating Diabetes via Inhibiting SGLT2. <i>Analytical Chemistry</i> , 2021 , 93, 4647-4656	7.8	3
146	An Activatable Probe with Aggregation-Induced Emission for Detecting and Imaging Herbal Medicine Induced Liver Injury with Optoacoustic Imaging and NIR-II Fluorescence Imaging. <i>Advanced Healthcare Materials</i> , 2021 , e2100867	10.1	9
145	A dopamine-precursor-based nanoprodrug for in-situ drug release and treatment of acute liver failure by inhibiting NLRP3 inflammasome and facilitating liver regeneration. <i>Biomaterials</i> , 2021 , 268, 120573	15.6	14
144	Emerging contrast agents for multispectral optoacoustic imaging and their biomedical applications. <i>Chemical Society Reviews</i> , 2021 , 50, 7924-7940	58.5	18
143	A turn-on probe for detecting antituberculous drug-induced liver injury in mice NIR-II fluorescence/optoacoustic imaging. <i>Chemical Communications</i> , 2021 , 57, 7842-7845	5.8	5
142	A Targeted Nanosystem for Detection of Inflammatory Diseases via Fluorescent/Optoacoustic Imaging and Therapy via Modulating Nrf2/NF- κ B Pathways. <i>Small</i> , 2021 , 17, e2102598	11	5
141	Refashioning benzothiadiazole dye as an activatable nanoprobe for biomarker detection with NIR-II fluorescence/optoacoustic imaging. <i>Cell Reports Physical Science</i> , 2021 , 100570	6.1	3
140	Biomarker-responsive nanoprobe with aggregation-induced emission for locating and guiding resection of deep-seated tumors via optoacoustic and NIR fluorescence imaging. <i>Materials Chemistry Frontiers</i> , 2021 , 5, 1962-1970	7.8	3
139	Synthesis of NQO1-activatable Optoacoustic Probe and Its Imaging of Breast Cancer. <i>Acta Chimica Sinica</i> , 2021 , 79, 331	3.3	2
138	ALP-activated probe for diagnosis of liver injury by multispectral optoacoustic tomography. <i>Methods in Enzymology</i> , 2021 , 657, 301-330	1.7	0
137	Activatable Nanocomposite Probe for Preoperative Location and Intraoperative Navigation for Orthotopic Hepatic Tumor Resection via MSOT and Aggregation-Induced Near-IR-I/II Fluorescence Imaging. <i>Analytical Chemistry</i> , 2020 , 92, 9257-9264	7.8	28
136	Activatable fluorescent probe based on aggregation-induced emission for detecting hypoxia-related pathological conditions. <i>Analytica Chimica Acta</i> , 2020 , 1125, 152-161	6.6	16

135	Nanoaggregate Probe for Breast Cancer Metastasis through Multispectral Optoacoustic Tomography and Aggregation-Induced NIR-I/II Fluorescence Imaging. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 10111-10121	16.4	87
134	Tetranitrile-anthracene as a probe for fluorescence detection of viscosity in fluid drinks via aggregation-induced emission. <i>Analyst, The</i> , 2020 , 145, 844-850	5	13
133	Nanoaggregate Probe for Breast Cancer Metastasis through Multispectral Optoacoustic Tomography and Aggregation-Induced NIR-I/II Fluorescence Imaging. <i>Angewandte Chemie</i> , 2020 , 132, 10197-10207	3.6	18
132	A biopolymer-based and inflammation-responsive nanodrug for rheumatoid arthritis treatment inhibiting JAK-STAT and JNK signalling pathways. <i>Nanoscale</i> , 2020 , 12, 23013-23027	7.7	8
131	An activatable probe for detecting alcoholic liver injury via multispectral optoacoustic tomography and fluorescence imaging. <i>Chemical Communications</i> , 2020 , 56, 11102-11105	5.8	13
130	An Activatable Nano-Prodrug for Treating Tyrosine-Kinase-Inhibitor-Resistant Non-Small Cell Lung Cancer and for Optoacoustic and Fluorescent Imaging. <i>Small</i> , 2020 , 16, e2003451	11	12
129	Near-Infrared Fluorescent Nanoprobe for Detecting Hydrogen Peroxide in Inflammation and Ischemic Kidney Injury. <i>Chinese Journal of Chemistry</i> , 2020 , 38, 1304-1310	4.9	3
128	Tetrazine-Mediated Bioorthogonal System for Prodrug Activation, Photothermal Therapy, and Optoacoustic Imaging. <i>ACS Applied Materials & Interfaces</i> , 2019 , 11, 41875-41888	9.5	6
127	A Gold Nanocage/Cluster Hybrid Structure for Whole-Body Multispectral Optoacoustic Tomography Imaging, EGFR Inhibitor Delivery, and Photothermal Therapy. <i>Small</i> , 2019 , 15, e1900309	11	46
126	Diagnosing Drug-Induced Liver Injury by Multispectral Optoacoustic Tomography and Fluorescence Imaging Using a Leucine-Aminopeptidase-Activated Probe. <i>Analytical Chemistry</i> , 2019 , 91, 8085-8092	7.8	39
125	An Activatable Near-Infrared Chromophore for Multispectral Optoacoustic Imaging of Tumor Hypoxia and for Tumor Inhibition. <i>Theranostics</i> , 2019 , 9, 7313-7324	12.1	45
124	An Unsymmetrical Squaraine-Based Activatable Probe for Imaging Lymphatic Metastasis by Responding to Tumor Hypoxia with MSOT and Aggregation-Enhanced Fluorescent Imaging. <i>Chemistry - A European Journal</i> , 2019 , 25, 16740	4.8	16
123	Bioorthogonal Nanosystem for Near-Infrared Fluorescence Imaging and Prodrug Activation in Mouse Model 2019 , 1, 549-557		11
122	A fluorescent probe based on aggregation-induced emission for hydrogen sulfide-specific assaying in food and biological systems. <i>Analyst, The</i> , 2019 , 144, 6570-6577	5	13
121	A Fluorescent Probe with Aggregation-Induced Emission for Detecting Alkaline Phosphatase and Cell Imaging. <i>Chemistry - an Asian Journal</i> , 2019 , 14, 802-808	4.5	32
120	A Nanoprobe for Diagnosing and Mapping Lymphatic Metastasis of Tumor Using 3D Multispectral Optoacoustic Tomography Owing to Aggregation/Deaggregation Induced Spectral Change. <i>Advanced Functional Materials</i> , 2019 , 29, 1807960	15.6	34
119	A conjugated-polymer-based ratiometric nanoprobe for evaluating in-vivo hepatotoxicity induced by herbal medicine via MSOT imaging. <i>Photoacoustics</i> , 2019 , 13, 6-17	9	21
118	A Turn-On Optoacoustic Probe for Imaging Metformin-Induced Upregulation of Hepatic Hydrogen Sulfide and Subsequent Liver Injury. <i>Theranostics</i> , 2019 , 9, 77-89	12.1	44

117	Tumor Inhibition Achieved by Targeting and Regulating Multiple Key Elements in EGFR Signaling Pathway Using a Self-Assembled Nanoprodrug. <i>Advanced Functional Materials</i> , 2018 , 28, 1800692	15.6	34
116	Therapeutic Nanosystem Consisting of Singlet-Oxygen-Responsive Prodrug and Photosensitizer Excited by Two-Photon Light. <i>ACS Medicinal Chemistry Letters</i> , 2018 , 9, 23-27	4.3	9
115	A sequential enzyme-activated and light-triggered pro-prodrug nanosystem for cancer detection and therapy. <i>Journal of Materials Chemistry B</i> , 2018 , 6, 2547-2556	7.3	21
114	A Fluorescent Probe for Early Detection of Melanoma and Its Metastasis by Specifically Imaging Tyrosinase Activity in a Mouse Model. <i>Analytical Chemistry</i> , 2018 , 90, 8807-8815	7.8	39
113	Oligo(ethylene glycol)-Functionalized Squaraine Fluorophore as a Near-Infrared-Fluorescent Probe for the In Vivo Detection of Diagnostic Enzymes. <i>Analytical Chemistry</i> , 2018 , 90, 9359-9365	7.8	26
112	A turn-on fluorescence probe based on aggregation-induced emission for leucine aminopeptidase in living cells and tumor tissue. <i>Analytica Chimica Acta</i> , 2018 , 1031, 169-177	6.6	34
111	Real-Time Monitoring of Endogenous Cysteine Levels In Vivo by near-Infrared Turn-on Fluorescent Probe with Large Stokes Shift. <i>Analytical Chemistry</i> , 2018 , 90, 1014-1020	7.8	155
110	Activatable probes for diagnosing and positioning liver injury and metastatic tumors by multispectral optoacoustic tomography. <i>Nature Communications</i> , 2018 , 9, 3983	17.4	95
109	A self-immolative and DT-diaphorase-activatable prodrug for drug-release tracking and therapy. <i>Journal of Materials Chemistry B</i> , 2017 , 5, 2635-2643	7.3	27
108	Fluorescent nanoprobe for in-vivo ratiometric imaging of endogenous hydrogen peroxide resulted from drug-induced organ damages. <i>Biosensors and Bioelectronics</i> , 2017 , 94, 278-285	11.8	40
107	A self-immolative prodrug nanosystem capable of releasing a drug and a NIR reporter for in vivo imaging and therapy. <i>Biomaterials</i> , 2017 , 139, 139-150	15.6	40
106	A bioorthogonal nanosystem for imaging and in vivo tumor inhibition. <i>Biomaterials</i> , 2017 , 138, 57-68	15.6	33
105	AIE fluorophore with enhanced cellular uptake for tracking esterase-activated release of taurine and ROS scavenging. <i>Faraday Discussions</i> , 2017 , 196, 335-350	3.6	8
104	An AIE-based fluorescent test strip for the portable detection of gaseous phosgene. <i>Chemical Communications</i> , 2017 , 53, 9813-9816	5.8	70
103	A two-photon-activated prodrug for therapy and drug release monitoring. <i>Journal of Materials Chemistry B</i> , 2017 , 5, 7538-7546	7.3	18
102	A theranostic prodrug based on FRET for real-time drug release monitoring in response to biothiols. <i>Materials Science and Engineering C</i> , 2017 , 72, 77-85	8.3	13
101	A highly selective fluorescent nanoprobe based on AIE and ESIPT for imaging hydrogen sulfide in live cells and zebrafish. <i>Materials Chemistry Frontiers</i> , 2017 , 1, 838-845	7.8	87
100	An anthracenecarboximide fluorescent probe for in vitro and in vivo ratiometric imaging of endogenous alpha-L-fucosidase for hepatocellular carcinoma diagnosis. <i>Materials Chemistry Frontiers</i> , 2017 , 1, 660-667	7.8	18

99	A ratiometric fluorescent probe for aluminum ions based-on monomer/excimer conversion and its applications to real samples. <i>Talanta</i> , 2016 , 151, 8-13	6.2	27
98	A two-photon fluorescent sensor revealing drug-induced liver injury via tracking Eglutamyltranspeptidase (GGT) level in vivo. <i>Biomaterials</i> , 2016 , 80, 46-56	15.6	102
97	A fluorescent probe for simultaneous discrimination of GSH and Cys/Hcy in human serum samples via distinctly-separated emissions with independent excitations. <i>Biosensors and Bioelectronics</i> , 2016 , 81, 341-348	11.8	115
96	A ratiometric fluorescent probe for hyaluronidase detection via hyaluronan-induced formation of red-light emitting excimers. <i>Biosensors and Bioelectronics</i> , 2016 , 79, 776-83	11.8	24
95	An ICT-based ratiometric fluorescent probe for hydrazine detection and its application in living cells and in vivo. <i>Sensors and Actuators B: Chemical</i> , 2016 , 227, 411-418	8.5	79
94	Pyrene Derivative Emitting Red or near-Infrared Light with Monomer/Excimer Conversion and Its Application to Ratiometric Detection of Hypochlorite. <i>ACS Applied Materials & Interfaces</i> , 2016 , 8, 1511-9	9.5	162
93	A fluorescent probe capable of discriminately and simultaneously detecting dl-dithiothreitol and single sulfhydryl-containing thiols. <i>Sensors and Actuators B: Chemical</i> , 2016 , 224, 88-94	8.5	8
92	Preparation of a Multifunctional Nano-carrier System Based on Carbon Dots with pH-Triggered Drug Release. <i>Acta Chimica Sinica</i> , 2016 , 74, 241	3.3	8
91	NIR AIE System for Tracking Release of Taurine and ROS Scavenging. <i>Acta Chimica Sinica</i> , 2016 , 74, 910	3.3	6
90	Temperature-responsive behavior of polymer fluorescent system via electrostatic interaction mediated aggregation/deaggregation. <i>Chinese Journal of Polymer Science (English Edition)</i> , 2016 , 34, 830-837	3.5	7
89	A fluorescent assay for Eglutamyltranspeptidase via aggregation induced emission and its applications in real samples. <i>Biosensors and Bioelectronics</i> , 2016 , 85, 317-323	11.8	52
88	Ratiometric detection and imaging of endogenous hypochlorite in live cells and in vivo achieved by using an aggregation induced emission (AIE)-based nanoprobe. <i>Chemical Communications</i> , 2016 , 52, 7288-91	5.8	132
87	Handy ratiometric detection of gaseous nerve agents with AIE-fluorophore-based solid test strips. <i>Journal of Materials Chemistry C</i> , 2016 , 4, 10105-10110	7.1	43
86	A Nanosystem Capable of Releasing a Photosensitizer Bioprecursor under Two-Photon Irradiation for Photodynamic Therapy. <i>Advanced Science</i> , 2016 , 3, 1500254	13.6	33
85	A ratiometric fluorescent system for carboxylesterase detection with AIE dots as FRET donors. <i>Chemical Communications</i> , 2015 , 51, 12791-4	5.8	86
84	Dual-targeting nanosystem for enhancing photodynamic therapy efficiency. <i>ACS Applied Materials & Interfaces</i> , 2015 , 7, 9287-96	9.5	81
83	A DT-diaphorase responsive theranostic prodrug for diagnosis, drug release monitoring and therapy. <i>Chemical Communications</i> , 2015 , 51, 9567-70	5.8	62
82	A ratiometric fluorescent probe for in vivo tracking of alkaline phosphatase level variation resulting from drug-induced organ damage. <i>Journal of Materials Chemistry B</i> , 2015 , 3, 1042-1048	7.3	49

81	A dual-targeting strategy to enhance photodynamic efficacy using a pH-responsive polymeric micelles. <i>Journal of Controlled Release</i> , 2015 , 213, e49	11.7	
80	A mitochondrial-targeting and NO-based anticancer nanosystem with enhanced photo-controllability and low dark-toxicity. <i>Journal of Materials Chemistry B</i> , 2015 , 3, 4904-4912	7.3	25
79	A logic gate-based fluorescent sensor for detecting H ₂ S and NO in aqueous media and inside live cells. <i>Chemical Communications</i> , 2015 , 51, 4414-6	5.8	72
78	Ratiometric fluorescence assay for α -glutamyltranspeptidase detection based on a single fluorophore via analyte-induced variation of substitution. <i>Chemical Communications</i> , 2014 , 50, 3417-20	5.8	65
77	Ratiometric fluorescent biosensor for hyaluronidase with hyaluronan as both nanoparticle scaffold and substrate for enzymatic reaction. <i>Biomacromolecules</i> , 2014 , 15, 3383-9	6.9	47
76	Low molecular weight PEIs modified by hydrazone-based crosslinker and betaine as improved gene carriers. <i>Colloids and Surfaces B: Biointerfaces</i> , 2014 , 122, 472-481	6	23
75	Ratiometric fluorescent probe for alkaline phosphatase based on betaine-modified polyethylenimine via excimer/monomer conversion. <i>Analytical Chemistry</i> , 2014 , 86, 9873-9	7.8	106
74	A targeted and FRET-based ratiometric fluorescent nanoprobe for imaging mitochondrial hydrogen peroxide in living cells. <i>Small</i> , 2014 , 10, 964-72	11	118
73	Preparation of a mitochondria-targeted and NO-releasing nanoplatform and its enhanced pro-apoptotic effect on cancer cells. <i>Small</i> , 2014 , 10, 3750-60	11	90
72	A novel ratiometric fluorescent probe through aggregation-induced emission and analyte-induced excimer dissociation. <i>Sensors and Actuators B: Chemical</i> , 2014 , 203, 504-510	8.5	18
71	A ratiometric fluorescent nanoprobe for HO sensing and in vivo detection of drug-induced oxidative damage to the digestive system. <i>Journal of Materials Chemistry B</i> , 2014 , 2, 8528-8537	7.3	40
70	Targeted anti-cancer prodrug based on carbon nanotube with photodynamic therapeutic effect and pH-triggered drug release. <i>Journal of Nanoparticle Research</i> , 2013 , 15, 1	2.3	17
69	Carbon-dot-based ratiometric fluorescent sensor for detecting hydrogen sulfide in aqueous media and inside live cells. <i>Chemical Communications</i> , 2013 , 49, 403-5	5.8	400
68	A water-soluble and specific BODIPY-based fluorescent probe for hypochlorite detection and cell imaging. <i>Analytical Methods</i> , 2013 , 5, 5589	3.2	61
67	Carbon dots-based fluorescent probes for sensitive and selective detection of iodide. <i>Mikrochimica Acta</i> , 2013 , 180, 453-460	5.8	139
66	Grafting zwitterionic polymer chains onto PEI as a convenient strategy to enhance gene delivery performance. <i>Polymer Chemistry</i> , 2013 , 4, 5810	4.9	28
65	A polylysine-based fluorescent probe for sulfite anion detection in aqueous media via analyte-induced charge generation and complexation. <i>Polymer Chemistry</i> , 2013 , 4, 5416	4.9	40
64	A low cytotoxic and ratiometric fluorescent nanosensor based on carbon-dots for intracellular pH sensing and mapping. <i>Nanotechnology</i> , 2013 , 24, 365101	3.4	86

63	A water-soluble, low-cytotoxic and sensitive fluorescent probe based on poly(ethylene glycol) for detecting sulfide anion in aqueous media and imaging inside live cells. <i>Polymer</i> , 2013 , 54, 5691-5697	3.9	27
62	A fluorescent ratiometric nanosensor for detecting NO in aqueous media and imaging exogenous and endogenous NO in live cells. <i>Journal of Materials Chemistry B</i> , 2013 , 1, 4152-4159	7.3	69
61	Conjugation with betaine: a facile and effective approach to significant improvement of gene delivery properties of PEI. <i>Biomacromolecules</i> , 2013 , 14, 728-36	6.9	54
60	A PEGylated fluorescent turn-on sensor for detecting fluoride ions in totally aqueous media and its imaging in live cells. <i>Chemistry - A European Journal</i> , 2013 , 19, 936-42	4.8	92
59	Water-dispersible fullerene aggregates as a targeted anticancer prodrug with both chemo- and photodynamic therapeutic actions. <i>Small</i> , 2013 , 9, 613-21	11	81
58	A robust, water-soluble and low cytotoxic fluorescent probe for sulfide anion achieved through incorporation of betaine. <i>Sensors and Actuators B: Chemical</i> , 2013 , 188, 1012-1018	8.5	20
57	One-pot fabrication of polymer nanoparticle-based chemosensors for Cu ²⁺ detection in aqueous media. <i>Polymer Chemistry</i> , 2013 , 4, 2325	4.9	41
56	Ratiometric sensing of mercury(II) based on a FRET process on silica core-shell nanoparticles acting as vehicles. <i>Mikrochimica Acta</i> , 2013 , 180, 845-853	5.8	29
55	Hyperbranched polyester-based fluorescent probe for histone deacetylase via aggregation-induced emission. <i>Biomacromolecules</i> , 2013 , 14, 4507-14	6.9	64
54	Polymer micelle with pH-triggered hydrophobic-hydrophilic transition and de-cross-linking process in the core and its application for targeted anticancer drug delivery. <i>Biomacromolecules</i> , 2012 , 13, 4126-37	6.9	88
53	A nanoparticle-supported fluorescence resonance energy transfer system formed via layer-by-layer approach as a ratiometric sensor for mercury ions in water. <i>Analytica Chimica Acta</i> , 2012 , 734, 69-78	6.6	31
52	A fast-responding fluorescent turn-on sensor for sensitive and selective detection of sulfite anions. <i>Analytical Methods</i> , 2012 , 4, 2638	3.2	74
51	A silica nanoparticle-based sensor for selective fluorescent detection of homocysteine via interaction differences between thiols and particle-surface-bound polymers. <i>Nanotechnology</i> , 2012 , 23, 305503	3.4	20
50	A facile approach for sensitive, reversible and ratiometric detection of biothiols based on thymine-mediated excimer-monomer transformation. <i>Chemical Communications</i> , 2012 , 48, 6007-9	5.8	34
49	CORE-SHELL POLYMER PARTICLES AS A FRET-BASED RATIOMETRIC SENSOR FOR MERCURY ION DETECTION IN WATER. <i>Acta Polymerica Sinica</i> , 2012 , 012, 666-672		3
48	Mesoporous silica particles for selective detection of dopamine with β -cyclodextrin as the selective barricade. <i>Chemical Communications</i> , 2011 , 47, 9086-8	5.8	41
47	FRET-based ratiometric detection system for mercury ions in water with polymeric particles as scaffolds. <i>Journal of Physical Chemistry B</i> , 2011 , 115, 874-82	3.4	98
46	A low dielectric constant polyimide/polyoxometalate composite. <i>Polymers for Advanced Technologies</i> , 2011 , 22, 209-214	3.2	12

45	A fluorescence turn-on sensor for iodide based on a thymine-Hg(II)-thymine complex. <i>Chemistry - A European Journal</i> , 2011 , 17, 14844-50	4.8	113
44	A FRET-based ratiometric sensor for mercury ions in water with multi-layered silica nanoparticles as the scaffold. <i>Chemical Communications</i> , 2011 , 47, 8913-5	5.8	82
43	Targeted anticancer prodrug with mesoporous silica nanoparticles as vehicles. <i>Nanotechnology</i> , 2011 , 22, 455102	3.4	65
42	Cyclodextrin as the vehicle for forming ratiometric mercury ion sensor usable in aqueous media, biological fluids, and live cells. <i>Langmuir</i> , 2010 , 26, 17764-71	4	63
41	Cyclodextrin supramolecular complex as a water-soluble ratiometric sensor for ferric ion sensing. <i>Langmuir</i> , 2010 , 26, 4529-34	4	133
40	Nanosized diblock copolymer micelles as a scaffold for constructing a ratiometric fluorescent sensor for metal ion detection in aqueous media. <i>Nanotechnology</i> , 2010 , 21, 195501	3.4	46
39	Construction of energy transfer systems within nanosized polymer micelles and their fluorescence modulation properties. <i>ChemPhysChem</i> , 2010 , 11, 1036-43	3.2	25
38	Reusable polymer film chemosensor for ratiometric fluorescence sensing in aqueous media. <i>Sensors and Actuators B: Chemical</i> , 2010 , 145, 451-456	8.5	71
37	Polyimide/polyoxometalate copolymer thin films: synthesis, thermal and dielectric properties. <i>Polymers for Advanced Technologies</i> , 2009 , 21, n/a-n/a	3.2	1
36	Supramolecular motif and macroscopic pattern in alpha-methylferrocenemethanol films cast from organic solution. <i>Applied Surface Science</i> , 2009 , 255, 4304-4308	6.7	
35	A facile approach for cupric ion detection in aqueous media using polyethyleneimine/PMMA core-shell fluorescent nanoparticles. <i>Nanotechnology</i> , 2009 , 20, 365502	3.4	41
34	Interactions Between Gold Nanoparticles and Polymer Bearing 3-Styryl Thiophene Chromophores. <i>Journal of Inorganic and Organometallic Polymers and Materials</i> , 2008 , 18, 463-471	3.2	
33	Synthesis and photochromic property of nanoparticles with spiropyran moieties via one-step miniemulsion polymerization. <i>Polymer Bulletin</i> , 2008 , 61, 425-434	2.4	31
32	A core-shell nanoparticle approach to photoreversible fluorescence modulation of a hydrophobic dye in aqueous media. <i>Chemistry - A European Journal</i> , 2008 , 14, 4851-60	4.8	83
31	Photoreversible fluorescence modulation of a rhodamine dye by supramolecular complexation with photosensitive cyclodextrin. <i>Angewandte Chemie - International Edition</i> , 2007 , 46, 7015-8	16.4	59
30	Photoreversible Fluorescence Modulation of a Rhodamine Dye by Supramolecular Complexation with Photosensitive Cyclodextrin. <i>Angewandte Chemie</i> , 2007 , 119, 7145-7148	3.6	8
29	Modulation of Fluorescence of a Terbium-Complex-Containing Polymer by Gold Nanoparticles through Energy Transfer. <i>Journal of Inorganic and Organometallic Polymers and Materials</i> , 2007 , 17, 679-685	3.2	7
28	Photoinduced Formation of Microscopic Ordering and Macroscopic Pattern in Spiropyran-Containing Polyacrylate-tetraoctylammonium Bromide Films. <i>Macromolecules</i> , 2007 , 40, 5060-5066	5.5	21

27	Tunability of Fluorescence Property of a Terbium-Complex-Containing Polymer via Incorporation of a Transition-Metal Complex. <i>Macromolecular Rapid Communications</i> , 2006 , 27, 937-942	4.8	6
26	Synthesis and fluorescence property of terbium complex with novel schiff-base macromolecular ligand. <i>European Polymer Journal</i> , 2006 , 42, 1670-1675	5.2	71
25	Novel Structure Change in Nonequimolar Complexes of Linear Poly(ethylenimine) and Octadecanoic Acid: Effects of Composition. <i>Macromolecules</i> , 2005 , 38, 5675-5680	5.5	16
24	Mesomorphous Structure and Macroscopic Patterns Formed by Polymer and Surfactant from Organic Solutions. <i>Macromolecules</i> , 2005 , 38, 9266-9274	5.5	8
23	Energy and electron transfers in photosensitive chitosan. <i>Journal of the American Chemical Society</i> , 2005 , 127, 2048-9	16.4	58
22	Preparation of colloidal crystals with polyhedral building blocks through post-polymerization. <i>Colloid and Polymer Science</i> , 2004 , 282, 651-655	2.4	6
21	Photo-induced birefringence and all-optical switching effect in azobenzene-grafted polyurethanes. <i>Optical Materials</i> , 2004 , 27, 585-590	3.3	21
20	Fluorescence study for the electrostatic interaction and aggregation in dilute polar solution of polyelectrolytes. <i>Macromolecular Symposia</i> , 2003 , 192, 251-264	0.8	4
19	Binding on strong polyelectrolytes of mixed ionic and nonionic surfactants below their critical micelle concentration observed by fluorescence. <i>Colloid and Polymer Science</i> , 2003 , 282, 141-148	2.4	5
18	Thermooptical properties of poly(methyl methacrylate)-based azobenzene composites. <i>Journal of Applied Polymer Science</i> , 2003 , 89, 2374-2377	2.9	2
17	All-Optical Switching Effect in Novel Chiral Biazobenzene Polymer Films. <i>Macromolecules</i> , 2003 , 36, 9292-9294	2.5	29
16	Preparation of Highly Charged, Monodisperse Nanospheres. <i>Macromolecular Chemistry and Physics</i> , 2002 , 203, 673-677	2.6	14
15	Fluorescence study of chromophore labeled strong polyelectrolyte bound with oppositely charged surfactant. <i>Colloid and Polymer Science</i> , 2002 , 280, 814-821	2.4	8
14	Fabrication of Inverse Opal via Ordered Highly Charged Colloidal Spheres. <i>Langmuir</i> , 2002 , 18, 9116-9120	2.4	49
13	Phase Separation and Network Formation in Poly(vinyl methyl ether)/Water Solutions. <i>Polymer Journal</i> , 2001 , 33, 399-403	2.7	12
12	Phase stability of the photoconductive polysiloxane-based photorefractive composites. <i>Polymer-Plastics Technology and Engineering</i> , 2001 , 40, 627-634		
11	Phase separation in poly(N-isopropyl acrylamide)/water solutions. II. Salt effects on cloud-point curves and gelation. <i>Journal of Polymer Science, Part B: Polymer Physics</i> , 2001 , 39, 901-907	2.6	20
10	Temperature dependence of the photorefractive property of PMMA-based composite. <i>European Polymer Journal</i> , 2001 , 37, 459-462	5.2	6

9	Thermal reversible gelation during phase separation of poly(N-isopropyl acrylamide)/water solution. <i>Science in China Series B: Chemistry</i> , 2000 , 43, 428-434		5
8	Molecular chain properties of poly (N-isopropyl acrylamide). <i>Science in China Series B: Chemistry</i> , 1999 , 42, 290-297		18
7	Effects of glass-transition temperature on properties of photorefractive polymer composite. <i>Journal of Polymer Science, Part B: Polymer Physics</i> , 1999 , 37, 3302-3306	2.6	4
6	Inverse Molecular Weight Dependence of Cloud Points for Aqueous Poly(N-isopropylacrylamide) Solutions. <i>Macromolecules</i> , 1999 , 32, 4488-4490	5.5	185
5	Electrorheological effect and conductive properties of carboxymethylchitosan suspensions. <i>Angewandte Makromolekulare Chemie</i> , 1998 , 260, 21-24		
4	Diffusion-controlled deposition of polyaniline onto poly(methyl methacrylate) substrates. <i>Polymer International</i> , 1998 , 47, 335-339	3.3	3
3	The conductive properties of the electrorheological suspensions based on dihydroxypropyl chitosan particles. <i>Journal of Applied Polymer Science</i> , 1998 , 67, 2077-2082	2.9	10
2	Phase Separation in Poly(N-isopropyl acrylamide)/Water Solutions I. Cloud Point Curves and Microgelation. <i>Polymer Journal</i> , 1998 , 30, 284-288	2.7	45
1	The Electrorheological Properties of Polypyrrole Suspensions. <i>Polymer Journal</i> , 1998 , 30, 451-454	2.7	10