## Erika Ramos-Tovar

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5562263/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Molecular Mechanisms That Link Oxidative Stress, Inflammation, and Fibrosis in the Liver. Antioxidants, 2020, 9, 1279.	2.2	131
2	ÂRedox state and methods to evaluate oxidative stress in liver damage: From bench to bedside. Annals of Hepatology, 2016, 15, 160-73.	0.6	81
3	Naringenin prevents experimental liver fibrosis by blocking TGFβ-Smad3 and JNK-Smad3 pathways. World Journal of Gastroenterology, 2017, 23, 4354.	1.4	62
4	Free radicals, antioxidants, nuclear factorâ€E2â€related factorâ€2 and liver damage. Journal of Applied Toxicology, 2020, 40, 151-168.	1.4	59
5	Fructose and the Liver. International Journal of Molecular Sciences, 2021, 22, 6969.	1.8	56
6	Stevia Prevents Acute and Chronic Liver Injury Induced by Carbon Tetrachloride by Blocking Oxidative Stress through Nrf2 Upregulation. Oxidative Medicine and Cellular Longevity, 2018, 2018, 1-12.	1.9	47
7	Antioxidant and immunomodulatory activity induced by stevioside in liver damage: In vivo, in vitro and in silico assays. Life Sciences, 2019, 224, 187-196.	2.0	38
8	Naringenin attenuates the progression of liver fibrosis via inactivation of hepatic stellate cells and profibrogenic pathways. European Journal of Pharmacology, 2019, 865, 172730.	1.7	32
9	Quercetin reverses experimental cirrhosis by immunomodulation of the proinflammatory and profibrotic processes. Fundamental and Clinical Pharmacology, 2017, 31, 610-624.	1.0	29
10	<i>Stevia rebaudiana</i> tea prevents experimental cirrhosis via regulation of NFâ€₽̂B, Nrf2, transforming growth factor beta, Smad7, and hepatic stellate cell activation. Phytotherapy Research, 2018, 32, 2568-2576.	2.8	22
11	Experimental Models of Liver Damage Mediated by Oxidative Stress. , 2017, , 529-546.		16
12	An aqueous extract of Stevia rebaudiana variety Morita II prevents liver damage in a rat model of cirrhosis that mimics the human disease. Annals of Hepatology, 2019, 18, 472-479.	0.6	16
13	Caffeine mitigates experimental nonalcoholic steatohepatitis and the progression of thioacetamide-induced liver fibrosis by blocking the MAPK and TGF-β/Smad3 signaling pathways. Annals of Hepatology, 2022, 27, 100671.	0.6	15
14	Stevioside inhibits experimental fibrosis by downâ€regulating profibrotic Smad pathways and blocking hepatic stellate cell activation. Basic and Clinical Pharmacology and Toxicology, 2019, 124, 670-680.	1.2	14
15	Curcumin downregulates Smad pathways and reduces hepatic stellate cells activation in experimental fibrosis. Annals of Hepatology, 2020, 19, 497-506.	0.6	14
16	Phytotherapy for the Liver. , 2019, , 101-121.		13
17	Cirrhosis induced by thioacetamide is prevented by stevia. Molecular mechanisms. Journal of Functional Foods, 2019, 52, 552-564.	1.6	11
18	Stevia prevents experimental cirrhosis by reducing hepatic myofibroblasts and modulating molecular profibrotic pathways. Hepatology Research, 2019, 49, 212-223.	1.8	10

#	Article	IF	CITATIONS
19	Stevia as a Putative Hepatoprotector. , 2017, , 715-727.		9
20	Does Nutrition Matter in Liver Disease?. , 2017, , 743-759.		2
21	CHAPTER 9. Hepatoprotective Effect of Coffee. , 2019, , 211-233.		1
22	Understanding the cellular and molecular mechanisms of hepatic fibrosis is essential for basic and clinical researchers. Annals of Hepatology, 2022, 27, 100732.	0.6	1