

# Klaus Rajewsky

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/5570245/klaus-rajewsky-publications-by-year.pdf>

**Version:** 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

140  
papers

26,457  
citations

68  
h-index

149  
g-index

149  
ext. papers

28,819  
ext. citations

16.3  
avg, IF

6.79  
L-index

#	Paper	IF	Citations
140	The magnitude of germinal center reactions is restricted by a fixed number of preexisting niches. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2021</b> , 118,	11.5	1
139	Protocol for Efficient CRISPR/Cas9/AAV-Mediated Homologous Recombination in Mouse Hematopoietic Stem and Progenitor Cells. <i>STAR Protocols</i> , <b>2020</b> , 1, 100028	1.4	3
138	Functional interplay of Epstein-Barr virus oncoproteins in a mouse model of B cell lymphomagenesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2020</b> , 117, 14421-14432	11.5	9
137	Immunoglobulin expression in the endoplasmic reticulum shapes the metabolic fitness of B lymphocytes. <i>Life Science Alliance</i> , <b>2020</b> , 3,	5.8	5
136	CRISPR-Cas9-Mediated ELANE Mutation Correction in Hematopoietic Stem and Progenitor Cells to Treat Severe Congenital Neutropenia. <i>Molecular Therapy</i> , <b>2020</b> , 28, 2621-2634	11.7	10
135	B-Cell-Specific Myd88 L252P Expression Causes a Premalignant Gammopathy Resembling IgM MGUS. <i>Frontiers in Immunology</i> , <b>2020</b> , 11, 602868	8.4	2
134	Efficient CRISPR/Cas9-Mediated Gene Knockin in Mouse Hematopoietic Stem and Progenitor Cells. <i>Cell Reports</i> , <b>2019</b> , 28, 3510-3522.e5	10.6	6
133	sgRNA Sequence Motifs Blocking Efficient CRISPR/Cas9-Mediated Gene Editing. <i>Cell Reports</i> , <b>2019</b> , 26, 1098-1103.e3	10.6	49
132	TET enzymes control antibody production and shape the mutational landscape in germinal centre B cells. <i>FEBS Journal</i> , <b>2019</b> , 286, 3566-3581	5.7	23
131	B-1a cells acquire their unique characteristics by bypassing the pre-BCR selection stage. <i>Nature Communications</i> , <b>2019</b> , 10, 4768	17.4	19
130	Context-specific regulation of cell survival by a miRNA-controlled BIM rheostat. <i>Genes and Development</i> , <b>2019</b> , 33, 1673-1687	12.6	7
129	BCR-dependent lineage plasticity in mature B cells. <i>Science</i> , <b>2019</b> , 363, 748-753	33.3	46
128	A novel allele for inducible Cre expression in germinal center B cells. <i>European Journal of Immunology</i> , <b>2019</b> , 49, 192-194	6.1	3
127	Signaling by the Epstein-Barr virus LMP1 protein induces potent cytotoxic CD4 and CD8 T cell responses. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2018</b> , 115, E686-E695	11.5	22
126	A novel conditional Aire allele enables cell-specific ablation of the immune tolerance regulator Aire. <i>European Journal of Immunology</i> , <b>2018</b> , 48, 546-548	6.1	4
125	The c-Myc/miR17-92/PTEN Axis Tunes PI3K Activity to Control Expression of Recombination Activating Genes in Early B Cell Development. <i>Frontiers in Immunology</i> , <b>2018</b> , 9, 2715	8.4	15
124	CRISPR/Cas9-Mediated In Vitro Mutagenesis in GC-Like B Cells. <i>Methods in Molecular Biology</i> , <b>2017</b> , 1623, 135-145	1.4	2

123	miRNAs Are Essential for the Regulation of the PI3K/AKT/FOXO Pathway and Receptor Editing during BCell Maturation. <i>Cell Reports</i> , <b>2016</b> , 17, 2271-2285	10.6	24
122	Mouse model for acute Epstein-Barr virus infection. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2016</b> , 113, 13821-13826	11.5	17
121	Efficient generation of Rosa26 knock-in mice using CRISPR/Cas9 in C57BL/6 zygotes. <i>BMC Biotechnology</i> , <b>2016</b> , 16, 4	3.5	133
120	Survival of IgD-deficient Mature B Cells Requires BAFF-R Function. <i>Journal of Immunology</i> , <b>2016</b> , 196, 2348-60	5.3	10
119	A c-Myc/miR17-92/Pten Axis Controls PI3K-Mediated Positive and Negative Selection in B Cell Development and Reconstitutes CD19 Deficiency. <i>Cell Reports</i> , <b>2016</b> , 16, 419-431	10.6	32
118	Canonical NF- $\kappa$ B signaling is uniquely required for the long-term persistence of functional mature B cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2016</b> , 113, 5065-70	11.5	15
117	Efficient CRISPR-mediated mutagenesis in primary immune cells using CrispRGold and a C57BL/6 Cas9 transgenic mouse line. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2016</b> , 113, 12514-12519	11.5	65
116	The DNA deamination model of somatic antibody diversification. <i>Journal of Immunology</i> , <b>2015</b> , 194, 2041-3	5.3	53
115	The Herzenberg lecture: how to make a B-1 cell?. <i>Annals of the New York Academy of Sciences</i> , <b>2015</b> , 1362, 6-7	6.5	7
114	Increasing the efficiency of homology-directed repair for CRISPR-Cas9-induced precise gene editing in mammalian cells. <i>Nature Biotechnology</i> , <b>2015</b> , 33, 543-8	44.5	771
113	An oncogenic role for alternative NF- $\kappa$ B signaling in DLBCL revealed upon deregulated BCL6 expression. <i>Cell Reports</i> , <b>2015</b> , 11, 715-26	10.6	57
112	The B-cell antigen receptor integrates adaptive and innate immune signals. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2015</b> , 112, 12145-50	11.5	22
111	Ectopic lymphoid structures function as microniches for tumor progenitor cells in hepatocellular carcinoma. <i>Nature Immunology</i> , <b>2015</b> , 16, 1235-44	19.1	178
110	Gab1 and Mapk Signaling Are Essential in the Hair Cycle and Hair Follicle Stem Cell Quiescence. <i>Cell Reports</i> , <b>2015</b> , 13, 561-572	10.6	28
109	Shp2/MAPK signaling controls goblet/paneth cell fate decisions in the intestine. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2014</b> , 111, 3472-7	11.5	83
108	STAT3 activation in Th17 and Th22 cells controls IL-22-mediated epithelial host defense during infectious colitis. <i>Journal of Immunology</i> , <b>2014</b> , 193, 3779-91	5.3	59
107	Years in Cologne. <i>Annual Review of Immunology</i> , <b>2013</b> , 31, 1-29	34.7	7
106	MicroRNA-17~92 plays a causative role in lymphomagenesis by coordinating multiple oncogenic pathways. <i>EMBO Journal</i> , <b>2013</b> , 32, 2377-91	13	106

105	Synergy between PI3K signaling and MYC in Burkitt lymphomagenesis. <i>Cancer Cell</i> , <b>2012</b> , 22, 167-79	24.3	212
104	The cell-cycle regulator c-Myc is essential for the formation and maintenance of germinal centers. <i>Nature Immunology</i> , <b>2012</b> , 13, 1092-100	19.1	258
103	Efficient generation of B lymphocytes by recognition of self-antigens. <i>European Journal of Immunology</i> , <b>2011</b> , 41, 2397-403	6.1	26
102	High-resolution description of antibody heavy-chain repertoires in humans. <i>PLoS ONE</i> , <b>2011</b> , 6, e22365	3.7	158
101	Constitutive canonical NF- $\kappa$ B activation cooperates with disruption of BLIMP1 in the pathogenesis of activated B cell-like diffuse large cell lymphoma. <i>Cancer Cell</i> , <b>2010</b> , 18, 580-9	24.3	146
100	MicroRNA control in the immune system: basic principles. <i>Cell</i> , <b>2009</b> , 136, 26-36	56.2	833
99	PI3 kinase signals BCR-dependent mature B cell survival. <i>Cell</i> , <b>2009</b> , 139, 573-86	56.2	482
98	Hyperactivable NFAT1 Ameliorates Autoimmune Encephalitis In Vivo.. <i>Blood</i> , <b>2009</b> , 114, 711-711	2.2	
97	Lymphoproliferative disease and autoimmunity in mice with increased miR-17-92 expression in lymphocytes. <i>Nature Immunology</i> , <b>2008</b> , 9, 405-14	19.1	1043
96	MicroRNA-155 suppresses activation-induced cytidine deaminase-mediated Myc-Igh translocation. <i>Immunity</i> , <b>2008</b> , 28, 630-8	32.3	391
95	Dicer ablation affects antibody diversity and cell survival in the B lymphocyte lineage. <i>Cell</i> , <b>2008</b> , 132, 860-74	56.2	486
94	Evidence for NFAT1 as a Tumor Suppressor in T-ALL. <i>Blood</i> , <b>2008</b> , 112, 3804-3804	2.2	
93	From a dream to reality. <i>European Journal of Immunology</i> , <b>2007</b> , 37 Suppl 1, S134-7	6.1	8
92	Joining the Institute of Genetics Early On as an Immunologist <b>2007</b> , 204-211		
91	Expression of Hyperactivable NFAT1 from the ROSA26 Locus Leads to Detrimental Effects during Embryonic Development.. <i>Blood</i> , <b>2007</b> , 110, 2296-2296	2.2	1
90	Tracking germinal center B cells expressing germ-line immunoglobulin gamma1 transcripts by conditional gene targeting. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2006</b> , 103, 7396-401	11.5	152
89	Canonical NF-kappaB activity, dispensable for B cell development, replaces BAFF-receptor signals and promotes B cell proliferation upon activation. <i>Immunity</i> , <b>2006</b> , 24, 729-739	32.3	258
88	Transcription factor IRF4 controls plasma cell differentiation and class-switch recombination. <i>Nature Immunology</i> , <b>2006</b> , 7, 773-82	19.1	526

87	IRF-4/MUM-1 Expression Is a Critical Switch in the Generation of Plasma Cells Versus Memory B-Cells.. <i>Blood</i> , <b>2005</b> , 106, 337-337	2.2	3
86	B cell receptor signal strength determines B cell fate. <i>Nature Immunology</i> , <b>2004</b> , 5, 317-27	19.1	456
85	Survival of resting mature B lymphocytes depends on BCR signaling via the Igalpha/beta heterodimer. <i>Cell</i> , <b>2004</b> , 117, 787-800	56.2	449
84	Rapid generation of inducible mouse mutants. <i>Nucleic Acids Research</i> , <b>2003</b> , 31, e12	20.1	243
83	Ability of the hydrophobic FGF and basic TAT peptides to promote cellular uptake of recombinant Cre recombinase: a tool for efficient genetic engineering of mammalian genomes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2002</b> , 99, 4489-94	11.5	275
82	Regulation of immunoglobulin light chain gene rearrangements during early B cell development in the human. <i>European Journal of Immunology</i> , <b>2001</b> , 31, 3631-3637	6.1	75
81	Receptor revision plays no major role in shaping the receptor repertoire of human memory B cells after the onset of somatic hypermutation. <i>European Journal of Immunology</i> , <b>2001</b> , 31, 3638-48	6.1	32
80	Molecular single-cell analysis of Hodgkin- and Reed-Sternberg cells harboring unmutated immunoglobulin variable region genes. <i>Laboratory Investigation</i> , <b>2001</b> , 81, 289-95	5.9	24
79	Survival and clonal expansion of mutating "forbidden" (immunoglobulin receptor-deficient) epstein-barr virus-infected b cells in angioimmunoblastic t cell lymphoma. <i>Journal of Experimental Medicine</i> , <b>2001</b> , 194, 927-40	16.6	93
78	Interference with immunoglobulin (Ig)alpha immunoreceptor tyrosine-based activation motif (ITAM) phosphorylation modulates or blocks B cell development, depending on the availability of an Igbeta cytoplasmic tail. <i>Journal of Experimental Medicine</i> , <b>2001</b> , 194, 455-69	16.6	105
77	B cell development is arrested at the immature B cell stage in mice carrying a mutation in the cytoplasmic domain of immunoglobulin beta. <i>Journal of Experimental Medicine</i> , <b>2001</b> , 193, 13-23	16.6	94
76	Indirect and direct evidence for DNA double-strand breaks in hypermutating immunoglobulin genes. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , <b>2001</b> , 356, 119-25	5.8	14
75	Homeostasis of peripheral B cells in the absence of B cell influx from the bone marrow. <i>Journal of Experimental Medicine</i> , <b>2001</b> , 194, 1151-64	16.6	240
74	Regulation of immunoglobulin light chain gene rearrangements during early B cell development in the human <b>2001</b> , 31, 3631		6
73	CD5-positive B cells in healthy elderly humans are a polyclonal B cell population. <i>European Journal of Immunology</i> , <b>2000</b> , 30, 2918-23	6.1	22
72	Memory B-cell persistence is independent of persisting immunizing antigen. <i>Nature</i> , <b>2000</b> , 407, 636-42	50.4	265
71	Clonal deleterious mutations in the I kappa B alpha gene in the malignant cells in Hodgkin's lymphoma. <i>Journal of Experimental Medicine</i> , <b>2000</b> , 191, 395-402	16.6	233
70	Clonal expansions of CD8(+) T cells dominate the T cell infiltrate in active multiple sclerosis lesions as shown by micromanipulation and single cell polymerase chain reaction. <i>Journal of Experimental Medicine</i> , <b>2000</b> , 192, 393-404	16.6	722

69	Molecular Analysis of Single B Cells From T-Cell-Rich B-Cell Lymphoma Shows the Derivation of the Tumor Cells From Mutating Germinal Center B Cells and Exemplifies Means by Which Immunoglobulin Genes Are Modified in Germinal Center B Cells. <i>Blood</i> , <b>1999</b> , 93, 2679-2687	2.2	87
68	Rearrangement and expression of immunoglobulin light chain genes can precede heavy chain expression during normal B cell development in mice. <i>Journal of Experimental Medicine</i> , <b>1999</b> , 189, 75-88	16.6	65
67	B cell antigen receptor specificity and surface density together determine B-1 versus B-2 cell development. <i>Journal of Experimental Medicine</i> , <b>1999</b> , 190, 471-7	16.6	182
66	Secondary V(D)J recombination in B-1 cells. <i>Nature</i> , <b>1999</b> , 397, 355-9	50.4	62
65	Mutation of the p53 Gene Is Not a Typical Feature of Hodgkin and Reed-Sternberg Cells in Hodgkin's Disease. <i>Blood</i> , <b>1999</b> , 94, 1755-1760	2.2	3
64	Somatic hypermutation in normal and transformed human B cells. <i>Immunological Reviews</i> , <b>1998</b> , 162, 261-80	11.3	294
63	The origin of Hodgkin and Reed/Sternberg cells in Hodgkin's disease. <i>Annual Review of Immunology</i> , <b>1998</b> , 16, 471-93	34.7	258
62	Temporally and spatially regulated somatic mutagenesis in mice. <i>Nucleic Acids Research</i> , <b>1998</b> , 26, 1427-32	20.1	157
61	Human immunoglobulin (Ig)M+IgD+ peripheral blood B cells expressing the CD27 cell surface antigen carry somatically mutated variable region genes: CD27 as a general marker for somatically mutated (memory) B cells. <i>Journal of Experimental Medicine</i> , <b>1998</b> , 188, 1679-89	16.6	903
60	The roles of gamma 1 heavy chain membrane expression and cytoplasmic tail in IgG1 responses. <i>Science</i> , <b>1997</b> , 276, 412-5	33.3	125
59	B cell development under the condition of allelic inclusion. <i>Immunity</i> , <b>1997</b> , 6, 225-33	32.3	189
58	Receptor editing in a transgenic mouse model: site, efficiency, and role in B cell tolerance and antibody diversification. <i>Immunity</i> , <b>1997</b> , 7, 765-75	32.3	246
57	In vivo ablation of surface immunoglobulin on mature B cells by inducible gene targeting results in rapid cell death. <i>Cell</i> , <b>1997</b> , 90, 1073-83	56.2	946
56	Evidence for a Large Compartment of IgM-Expressing Memory B Cells in Humans. <i>Blood</i> , <b>1997</b> , 89, 1288-1298	12.9	267
55	Diffuse large cell lymphomas are derived from mature B cells carrying V region genes with a high load of somatic mutation and evidence of selection for antibody expression. <i>European Journal of Immunology</i> , <b>1997</b> , 27, 1398-405	6.1	118
54	Molecular Ig gene analysis reveals that monocytoid B cell lymphoma is a malignancy of mature B cells carrying somatically mutated V region genes and suggests that rearrangement of the kappa-deleting element (resulting in deletion of the Ig kappa enhancers) abolishes somatic hypermutation in the human. <i>European Journal of Immunology</i> , <b>1996</b> , 26, 1794-800	6.1	56
53	Somatic hypermutation occurs in B cells of terminal deoxynucleotidyl transferase-, CD23-, interleukin-4-, IgD- and CD30-deficient mouse mutants. <i>European Journal of Immunology</i> , <b>1996</b> , 26, 1966-9	6.1	16
52	Immunity to viruses in B cell-deficient mice: influence of antibodies on virus persistence and on T cell memory. <i>European Journal of Immunology</i> , <b>1996</b> , 26, 2257-62	6.1	87

51	Requirement of mammalian DNA polymerase-beta in base-excision repair. <i>Nature</i> , <b>1996</b> , 379, 183-6	50.4	751
50	Clonal selection and learning in the antibody system. <i>Nature</i> , <b>1996</b> , 381, 751-8	50.4	1403
49	Critical role for beta7 integrins in formation of the gut-associated lymphoid tissue. <i>Nature</i> , <b>1996</b> , 382, 366-70	50.4	478
48	Defective antigen receptor-mediated proliferation of B and T cells in the absence of Vav. <i>Nature</i> , <b>1995</b> , 374, 467-70	50.4	376
47	Impairment of T-cell-dependent B-cell responses and B-1 cell development in CD19-deficient mice. <i>Nature</i> , <b>1995</b> , 376, 352-5	50.4	587
46	A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. <i>Nucleic Acids Research</i> , <b>1995</b> , 23, 5080-1	20.1	1021
45	Rearrangement of upstream DH and VH genes to a rearranged immunoglobulin variable region gene inserted into the DQ52-JH region of the immunoglobulin heavy chain locus. <i>European Journal of Immunology</i> , <b>1995</b> , 25, 1888-96	6.1	58
44	Development and proliferation of lymphocytes in mice deficient for both interleukins-2 and -4. <i>European Journal of Immunology</i> , <b>1994</b> , 24, 281-4	6.1	131
43	Lymphocyte populations and immune responses in CD5-deficient mice. <i>European Journal of Immunology</i> , <b>1994</b> , 24, 1678-84	6.1	78
42	Early B-cell development in the mouse: insights from mutations introduced by gene targeting. <i>Immunological Reviews</i> , <b>1994</b> , 137, 135-53	11.3	115
41	Inactivation of the N-CAM gene in mice results in size reduction of the olfactory bulb and deficits in spatial learning. <i>Nature</i> , <b>1994</b> , 367, 455-9	50.4	893
40	Human IgM+IgD+ B cells, the major B cell subset in the peripheral blood, express V kappa genes with no or little somatic mutation throughout life. <i>European Journal of Immunology</i> , <b>1993</b> , 23, 3272-7	6.1	143
39	A critical role of lambda 5 protein in B cell development. <i>Cell</i> , <b>1992</b> , 69, 823-31	56.2	544
38	Targeted disruption of mu chain membrane exon causes loss of heavy-chain allelic exclusion. <i>Nature</i> , <b>1992</b> , 356, 154-6	50.4	332
37	An explanation for the defect in secretion of IgM Mott cells and their predominant occurrence in the Ly-1 B cell compartment. <i>European Journal of Immunology</i> , <b>1992</b> , 22, 531-9	6.1	16
36	A B cell-deficient mouse by targeted disruption of the membrane exon of the immunoglobulin mu chain gene. <i>Nature</i> , <b>1991</b> , 350, 423-6	50.4	1540
35	Intraclonal generation of antibody mutants in germinal centres. <i>Nature</i> , <b>1991</b> , 354, 389-92	50.4	913
34	Dividing cells in bone marrow and spleen incorporate bromodeoxyuridine with high efficiency. <i>European Journal of Immunology</i> , <b>1991</b> , 21, 235-8	6.1	34

33	Major histocompatibility complex class II hyperexpression on B cells in interleukin 4-transgenic mice does not lead to B cell proliferation and hypergammaglobulinemia. <i>European Journal of Immunology</i> , <b>1991</b> , 21, 921-5	6.1	37
32	Generation of long-lived B cells in germ-free mice. <i>European Journal of Immunology</i> , <b>1991</b> , 21, 1779-82	6.1	14
31	Maintenance of B-cell memory by long-lived cells generated from proliferating precursors. <i>Nature</i> , <b>1990</b> , 346, 749-51	50.4	284
30	The half-lives of serum immunoglobulins in adult mice. <i>European Journal of Immunology</i> , <b>1988</b> , 18, 313-66.1		333
29	Expansion and functional activity of Ly-1+ B cells upon transfer of peritoneal cells into allotype-congenic, newborn mice. <i>European Journal of Immunology</i> , <b>1987</b> , 17, 521-8	6.1	279
28	A new V gene expressed in lambda-2 light chains of the mouse. <i>European Journal of Immunology</i> , <b>1987</b> , 17, 731-4	6.1	50
27	In vivo generation and function of B cells in the presence of a monoclonal anti-IgM antibody: implications for B cell tolerance. <i>European Journal of Immunology</i> , <b>1987</b> , 17, 981-90	6.1	25
26	Timing, genetic requirements and functional consequences of somatic hypermutation during B-cell development. <i>Immunological Reviews</i> , <b>1987</b> , 96, 5-22	11.3	204
25	The bulk of endogenously produced IgG2a is eliminated from the serum of adult C57BL/6 mice with a half-life of 6-8 days. <i>European Journal of Immunology</i> , <b>1986</b> , 16, 871-4	6.1	26
24	Induction of chronic idiotype suppression by ligands binding to the variable (not the constant) region of the idiotypic target. <i>European Journal of Immunology</i> , <b>1986</b> , 16, 1419-25	6.1	6
23	Drastic change in idiotypic but not antigen-binding specificity of an antibody by a single amino-acid substitution. <i>Nature</i> , <b>1985</b> , 315, 506-8	50.4	68
22	Structure of primary anti-(4-hydroxy-3-nitrophenyl)acetyl (NP) antibodies in normal and idiotypically suppressed C57BL/6 mice. <i>European Journal of Immunology</i> , <b>1985</b> , 15, 512-20	6.1	176
21	A self-Ia reactive T cell clone directly stimulates every hundredth B cell and helps antigen-specific B cell responses. <i>European Journal of Immunology</i> , <b>1985</b> , 15, 927-34	6.1	35
20	VH-gene expression in murine lipopolysaccharide blasts distributes over the nine known VH-gene groups and may be random. <i>European Journal of Immunology</i> , <b>1985</b> , 15, 1154-6	6.1	118
19	Mechanism of neonatally induced idiotype suppression and its relevance for the acquisition of self-tolerance. <i>Immunological Reviews</i> , <b>1984</b> , 79, 103-17	11.3	43
18	Heterogeneous and monoclonal helper T cells induce similar anti-(4-hydroxy-3-nitrophenyl)acetyl (NP) antibody populations in the primary adoptive response. I. Isotype distribution. <i>European Journal of Immunology</i> , <b>1984</b> , 14, 188-94	6.1	21
17	Heterogeneous and monoclonal helper T cells induce similar anti-(4-hydroxy-3-nitrophenyl)acetyl (NP) antibody populations in the primary adoptive response. II. Lambda light chain dominance and idiotope expression. <i>European Journal of Immunology</i> , <b>1984</b> , 14, 195-200	6.1	16
16	Specificity, duration and mechanism of idiotype suppression induced by neonatal injection of monoclonal anti-idiotope antibodies into mice. <i>European Journal of Immunology</i> , <b>1984</b> , 14, 656-67	6.1	44



15	The expression of a set of antibody variable regions in lipopolysaccharide-reactive B cells at various stages of ontogeny and its control by anti-idiotypic antibody. <i>European Journal of Immunology</i> , <b>1983</b> , 13, 318-25	6.1	42
14	The immune response against anti-idiotope antibodies II. The induction of antibodies bearing the target idiotope (Ab3 beta) depends on the frequency of the corresponding B cells. <i>European Journal of Immunology</i> , <b>1983</b> , 13, 726-32	6.1	16
13	Somatic H-2Kk variants reveal nonidentity of serological and cytotoxic T cell-defined Kk determinants. <i>European Journal of Immunology</i> , <b>1983</b> , 13, 846-51	6.1	12
12	The immune response against anti-idiotope antibodies. I. Induction of idiotope-bearing antibodies and analysis of the idiotope repertoire. <i>European Journal of Immunology</i> , <b>1982</b> , 12, 1040-6	6.1	63
11	Spontaneous immunoglobulin class switching in myeloma and hybridoma cell lines differs from physiological class switching. <i>Immunological Reviews</i> , <b>1982</b> , 67, 59-72	11.3	57
10	Control of idiotope expression by monoclonal anti-idiotope and idiotope-bearing antibody. <i>European Journal of Immunology</i> , <b>1981</b> , 11, 418-23	6.1	63
9	Lambda chain expression at different stages of ontogeny in C57BL/6, BALB/c and SJL mice. <i>European Journal of Immunology</i> , <b>1981</b> , 11, 618-25	6.1	71
8	Activation of mouse complement by monoclonal mouse antibodies. <i>European Journal of Immunology</i> , <b>1981</b> , 11, 1012-6	6.1	191
7	Control idiotope expression by monoclonal anti-idiotope antibodies. <i>Immunological Reviews</i> , <b>1980</b> , 52, 75-88	11.3	106
6	The effect of light chain gene expression on the inheritance of an idiootype associated with primary anti-(4-hydroxy-3-nitrophenyl)acetyl(NP) antibodies. <i>European Journal of Immunology</i> , <b>1979</b> , 9, 324-31	6.1	55
5	Isolated hapten-binding receptors of sensitized lymphocytes. IV. Expression of immunoglobulin variable regions in (4-hydroxy-3-nitrophenyl) acetyl (NP)-specific receptors isolated from murine B and T lymphocytes. <i>European Journal of Immunology</i> , <b>1979</b> , 9, 332-8	6.1	46
4	Isolated hapten-binding receptors of sensitized lymphocytes. V. Cellular origin of receptor molecules. <i>European Journal of Immunology</i> , <b>1979</b> , 9, 815-20	6.1	22
3	Analysis of the repertoire of anti-(4-hydroxy-3-nitrophenyl)acetyl (NP) antibodies in C 57 BL/6 mice by cell fusion. II. Characterization of idiotopes by monoclonal anti-idiotope antibodies. <i>European Journal of Immunology</i> , <b>1979</b> , 9, 1004-13	6.1	250
2	B-1a cells acquire their unique characteristics by bypassing the pre-BCR selection stage		1
1	Histone methyltransferase DOT1L controls state-specific identity during B cell differentiation		1