

Che-Hsiung Wu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/557425/publications.pdf>

Version: 2024-02-01

16
papers

592
citations

933264

10
h-index

940416

16
g-index

16
all docs

16
docs citations

16
times ranked

898
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Volume overload correlates with cardiovascular risk factors in patients with chronic kidney disease. <i>Kidney International</i> , 2014, 85, 703-709. | 2.6 | 194 |
| 2 | Case detection and diagnosis of primary aldosteronism – The consensus of Taiwan Society of Aldosteronism. <i>Journal of the Formosan Medical Association</i> , 2017, 116, 993-1005. | 0.8 | 85 |
| 3 | Prevalence and clinical correlates of somatic mutation in aldosterone producing adenoma-Taiwanese population. <i>Scientific Reports</i> , 2015, 5, 11396. | 1.6 | 78 |
| 4 | Administrative data on diagnosis and mineralocorticoid receptor antagonist prescription identified patients with primary aldosteronism in Taiwan. <i>Journal of Clinical Epidemiology</i> , 2014, 67, 1139-1149. | 2.4 | 54 |
| 5 | Urinary Î-glutathione S-transferase Predicts Advanced Acute Kidney Injury Following Cardiovascular Surgery. <i>Scientific Reports</i> , 2016, 6, 26335. | 1.6 | 40 |
| 6 | Association of Fluid Retention With Anemia and Clinical Outcomes Among Patients With Chronic Kidney Disease. <i>Journal of the American Heart Association</i> , 2015, 4, e001480. | 1.6 | 34 |
| 7 | Comparison of 24-h Urinary Aldosterone Level and Random Urinary Aldosterone-to-Creatinine Ratio in the Diagnosis of Primary Aldosteronism. <i>PLoS ONE</i> , 2013, 8, e67417. | 1.1 | 22 |
| 8 | Meglitinides increase the risk of hypoglycemia in diabetic patients with advanced chronic kidney disease: a nationwide, population-based study. <i>Oncotarget</i> , 2017, 8, 78086-78095. | 0.8 | 15 |
| 9 | Risk of liver injury after Î-glucosidase inhibitor therapy in advanced chronic kidney disease patients. <i>Scientific Reports</i> , 2016, 6, 18996. | 1.6 | 12 |
| 10 | Plasma Aldosterone After Seated Saline Infusion Test Outperforms Captopril Test at Predicting Clinical Outcomes After Adrenalectomy for Primary Aldosteronism. <i>American Journal of Hypertension</i> , 2019, 32, 1066-1074. | 1.0 | 12 |
| 11 | Nationwide epidemiology and prognosis of dialysis-requiring acute kidney injury (NEP study): Design and methods. <i>Nephrology</i> , 2016, 21, 758-764. | 0.7 | 11 |
| 12 | Risk of ischemic stroke in primary aldosteronism patients. <i>Clinica Chimica Acta</i> , 2015, 438, 86-89. | 0.5 | 9 |
| 13 | Novel Mutations Detection with Next-Generation Sequencing and Its Association with Clinical Outcome in Unilateral Primary Aldosteronism. <i>Biomedicines</i> , 2021, 9, 1167. | 1.4 | 8 |
| 14 | Improvement in Mortality and End-Stage Renal Disease in Patients With Type 2 Diabetes After Acute Kidney Injury Who Are Prescribed Dipeptidyl Peptidase-4 Inhibitors. <i>Mayo Clinic Proceedings</i> , 2018, 93, 1760-1774. | 1.4 | 7 |
| 15 | High plasma C-terminal FGF-23 levels predict poor outcomes in patients with chronic kidney disease superimposed with acute kidney injury. <i>Therapeutic Advances in Chronic Disease</i> , 2020, 11, 204062232096416. | 1.1 | 7 |
| 16 | Hemojuvelin Predicts Acute Kidney Injury and Poor Outcomes Following Cardiac Surgery. <i>Scientific Reports</i> , 2018, 8, 1938. | 1.6 | 4 |