

Che-Hsiung Wu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/557425/publications.pdf>

Version: 2024-02-01

16
papers

592
citations

932766

10
h-index

940134

16
g-index

16
all docs

16
docs citations

16
times ranked

898
citing authors

#	ARTICLE	IF	CITATIONS
1	Novel Mutations Detection with Next-Generation Sequencing and Its Association with Clinical Outcome in Unilateral Primary Aldosteronism. <i>Biomedicines</i> , 2021, 9, 1167.	1.4	8
2	High plasma C-terminal FGF-23 levels predict poor outcomes in patients with chronic kidney disease superimposed with acute kidney injury. <i>Therapeutic Advances in Chronic Disease</i> , 2020, 11, 204062232096416.	1.1	7
3	Plasma Aldosterone After Seated Saline Infusion Test Outperforms Captopril Test at Predicting Clinical Outcomes After Adrenalectomy for Primary Aldosteronism. <i>American Journal of Hypertension</i> , 2019, 32, 1066-1074.	1.0	12
4	Hemojuvelin Predicts Acute Kidney Injury and Poor Outcomes Following Cardiac Surgery. <i>Scientific Reports</i> , 2018, 8, 1938.	1.6	4
5	Improvement in Mortality and End-Stage Renal Disease in Patients With Type 2 Diabetes After Acute Kidney Injury Who Are Prescribed Dipeptidyl Peptidase-4 Inhibitors. <i>Mayo Clinic Proceedings</i> , 2018, 93, 1760-1774.	1.4	7
6	Case detection and diagnosis of primary aldosteronism – The consensus of Taiwan Society of Aldosteronism. <i>Journal of the Formosan Medical Association</i> , 2017, 116, 993-1005.	0.8	85
7	Meglitinides increase the risk of hypoglycemia in diabetic patients with advanced chronic kidney disease: a nationwide, population-based study. <i>Oncotarget</i> , 2017, 8, 78086-78095.	0.8	15
8	Nationwide epidemiology and prognosis of dialysis-requiring acute kidney injury (NEP-RAKI) study: Design and methods. <i>Nephrology</i> , 2016, 21, 758-764.	0.7	11
9	Urinary Î-glutathione S-transferase Predicts Advanced Acute Kidney Injury Following Cardiovascular Surgery. <i>Scientific Reports</i> , 2016, 6, 26335.	1.6	40
10	Risk of liver injury after Î-glucosidase inhibitor therapy in advanced chronic kidney disease patients. <i>Scientific Reports</i> , 2016, 6, 18996.	1.6	12
11	Prevalence and clinical correlates of somatic mutation in aldosterone producing adenoma-Taiwanese population. <i>Scientific Reports</i> , 2015, 5, 11396.	1.6	78
12	Association of Fluid Retention With Anemia and Clinical Outcomes Among Patients With Chronic Kidney Disease. <i>Journal of the American Heart Association</i> , 2015, 4, e001480.	1.6	34
13	Risk of ischemic stroke in primary aldosteronism patients. <i>Clinica Chimica Acta</i> , 2015, 438, 86-89.	0.5	9
14	Volume overload correlates with cardiovascular risk factors in patients with chronic kidney disease. <i>Kidney International</i> , 2014, 85, 703-709.	2.6	194
15	Administrative data on diagnosis and mineralocorticoid receptor antagonist prescription identified patients with primary aldosteronism in Taiwan. <i>Journal of Clinical Epidemiology</i> , 2014, 67, 1139-1149.	2.4	54
16	Comparison of 24-h Urinary Aldosterone Level and Random Urinary Aldosterone-to-Creatinine Ratio in the Diagnosis of Primary Aldosteronism. <i>PLoS ONE</i> , 2013, 8, e67417.	1.1	22