

# Radka Kořková

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5575758/publications.pdf>

Version: 2024-02-01

11  
papers

359  
citations

1478505

6  
h-index

1372567

10  
g-index

11  
all docs

11  
docs citations

11  
times ranked

502  
citing authors

#	ARTICLE	IF	CITATIONS
1	Aortic Valve Replacement Versus Conservative Treatment in Asymptomatic Severe Aortic Stenosis: The AVATAR Trial. <i>Circulation</i> , 2022, 145, 648-658.	1.6	130
2	Assessment of Asymptomatic Severe Aortic Regurgitation by Doppler-Derived Echo Indices: Comparison with Magnetic Resonance Quantification. <i>Journal of Clinical Medicine</i> , 2022, 11, 152.	2.4	3
3	New Imaging Markers of Clinical Outcome in Asymptomatic Patients with Severe Aortic Regurgitation. <i>Journal of Clinical Medicine</i> , 2019, 8, 1654.	2.4	4
4	Prognostic Implications of Magnetic Resonance-Derived Quantification in Asymptomatic Patients With Organic Mitral Regurgitation. <i>Circulation</i> , 2018, 137, 1349-1360.	1.6	104
5	Endocardial Fibroelastosis is Secondary to Hemodynamic Alterations in the Chick Embryonic Model of Hypoplastic Left Heart Syndrome. <i>Developmental Dynamics</i> , 2018, 247, 509-520.	1.8	24
6	Cardiac resynchronization therapy guided by cardiac magnetic resonance imaging: A prospective, single-centre randomized study (CMR-CRT). <i>International Journal of Cardiology</i> , 2018, 270, 325-330.	1.7	16
7	Native T1 Relaxation Time and Extracellular Volume Fraction as Accurate Markers of Diffuse Myocardial Fibrosis in Heart Valve Disease—Comparison With Targeted Left Ventricular Myocardial Biopsy. <i>Circulation Journal</i> , 2016, 80, 1202-1209.	1.6	46
8	Abstract 13596: Magnetic Resonance - Derived Pre-contrast T1 Relaxation Time is the Accurate Marker of Diffuse Myocardial Fibrosis in Severe Aortic Valve Disease: A Comparison With Left Ventricular Myocardial Biopsy. <i>Circulation</i> , 2015, 132, .	1.6	0
9	Adenylyl Cyclase Signaling in the Developing Chick Heart: The Deranging Effect of Antiarrhythmic Drugs. <i>BioMed Research International</i> , 2014, 2014, 1-6.	1.9	4
10	Three-Dimensional Echocardiography-Derived Vena Contracta Area at Rest and Its Increase During Exercise Predicts Clinical Outcome in Mild-Moderate Functional Mitral Regurgitation. <i>Circulation Journal</i> , 2014, 78, 2741-2749.	1.6	5
11	Heart rate changes mediate the embryotoxic effect of antiarrhythmic drugs in the chick embryo. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2013, 304, H895-H902.	3.2	23